


FORM PTO-1390 U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER <b>PF-0662 USN</b>
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		U.S. APPLICATION NO. (UNKNOWN) 37 CFR 1.5 TO BE ASSIGNED <b>09/889616</b>
INTERNATIONAL APPLICATION NO. PCT/US00/02237	INTERNATIONAL FILING DATE 28 January 2000	PRIORITY DATE CLAIMED 29 January 1999
TITLE OF INVENTION <b>CANCER-ASSOCIATED PROTEINS</b>		
APPLICANT(S) FOR DO/EO/US <b>INCYTE PHARMACEUTICALS, INC.; TANG, Y. Tom; LAL, Preeti; HILLMAN, Jennifer L.; YUE, Henry; AZIMZAI, Yalda; LU, Dyung Aina M.; BAUGHN, Mariah R.; TRAN, Bao; SHIH, Leo L.; AU-YOUNG, Janice</b>		
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:		
<ol style="list-style-type: none"> <li>1. <input checked="" type="checkbox"/> This is the <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371.</li> <li>2. <input type="checkbox"/> This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371.</li> <li>3. <input type="checkbox"/> This is an express request to promptly begin national examination procedures (35 U.S.C. 371 (f)).</li> <li>4. <input type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (PCT Article 31).</li> <li>5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2))             <ol style="list-style-type: none"> <li>a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau)</li> <li>b. <input type="checkbox"/> has been communicated by the International Bureau.</li> <li>c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).</li> </ol> </li> <li>6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).</li> <li>7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))             <ol style="list-style-type: none"> <li>a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau).</li> <li>b. <input type="checkbox"/> have been communicated by the International Bureau.</li> <li>c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.</li> <li>d. <input type="checkbox"/> have not been made and will not be made.</li> </ol> </li> <li>8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).</li> <li>9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).</li> <li>10. <input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).</li> </ol>		
Items 11 to 16 below concern document(s) or information included:		
<ol style="list-style-type: none"> <li>11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98.</li> <li>12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.27 and 3.31 is included.</li> <li>13. <input type="checkbox"/> A <b>FIRST</b> preliminary amendment.             <ol style="list-style-type: none"> <li><input type="checkbox"/> A <b>SECOND</b> or <b>SUBSEQUENT</b> preliminary amendment.</li> </ol> </li> <li>14. <input type="checkbox"/> A substitute specification.</li> <li>15. <input type="checkbox"/> A change of power of attorney and/or address letter.</li> <li>16. <input checked="" type="checkbox"/> Other items or information:             <ol style="list-style-type: none"> <li>1) Transmittal Letter (2 pp, in duplicate)</li> <li>2) Return Postcard</li> <li>3) Express Mail Label No.: <b>EL 856 154 129 US</b></li> <li>4) Request to Transfer</li> </ol> </li> </ol>		

JC18 Rec'd PCT/PTO 16 JUL 2001

U.S. APPLICATION NO. 09/889616 TO BE ASSIGNED		INTERNATIONAL APPLICATION NO. PCT/US00/02237		ATTORNEY'S DOCKET NUMBER PF-0662 USN	
17. <input checked="" type="checkbox"/> The following fees are submitted <b>BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):</b> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO .....\$1000.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO .....\$860.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO .....\$710.00 <input checked="" type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) .....\$690.00 <input type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) .....\$100.00					
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$690.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total Claims	23 =	3	X \$ 18.00	\$ 54.00	
Independent Claims	2 =	0	X \$ 80.00	\$	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$270.00	\$	
TOTAL OF ABOVE CALCULATIONS =				\$744.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$	
SUBTOTAL =				\$744.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	
TOTAL NATIONAL FEE =				\$744.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by the appropriate cover sheet (37 CFR 3.28, 3.31) \$40.00 per property				\$	
TOTAL FEES ENCLOSED =				\$744.00	
				Amount to be Refunded	\$
				Charged	\$
a. <input type="checkbox"/> A check in the amount of \$_____ to cover the above fees is enclosed. b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. 09-0108 in the amount of \$ 744.00 to cover the above fees c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 09-0108. A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO					
INCYTE GENOMICS, INC. 3160 Porter Drive Palo Alto, CA 94304			 SIGNATURE		
NAME: Diana Hamlet-Cox					
REGISTRATION NUMBER: 33,302					
DATE: 16 July 2001					

## NUCLEIC-ACID BINDING PROTEINS

## TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of nucleic-acid binding proteins and to the use of these sequences in the diagnosis, treatment, and prevention of reproductive, immune, and neurological disorders, and cell proliferative disorders including cancer.

5

## BACKGROUND OF THE INVENTION

Multicellular organisms are comprised of diverse cell types that differ dramatically both in structure and function. The identity of a cell is determined by its characteristic pattern of gene expression, and different cell types express overlapping but distinct sets of genes throughout development. Spatial and temporal regulation of gene expression is critical for the control of cell proliferation, cell differentiation, apoptosis, and other processes that contribute to organismal development. Furthermore, gene expression is regulated in response to extracellular signals that mediate cell-cell communication and coordinate the activities of different cell types. Appropriate gene regulation also ensures that cells function efficiently by expressing only those genes whose functions are required at a given time.

Transcriptional regulatory proteins are essential for the control of gene expression. Some of these proteins function as transcription factors that initiate, activate, repress, or terminate gene transcription. Transcription factors generally bind to promoter, enhancer, or upstream regulatory regions of a gene in a sequence-specific manner, although some factors bind regulatory elements within or downstream of the coding region. Transcription factors may bind to a specific region of DNA singly or as a complex with other accessory factors. (Reviewed in Lewin, B. (1990) Genes IV, Oxford University Press, New York, NY, pp. 554-570.)

The double helix structure and repeated sequences of DNA create topological and chemical features which can be recognized by transcription factors. These features include hydrogen bond donor and acceptor groups, hydrophobic patches, major and minor grooves, and regular repeated stretches of sequence which induce distinct bends in the helix. Typically, transcription factors recognize specific DNA sequence motifs of about 20 nucleotides in length. Multiple adjacent transcription factor-binding motifs may be required for gene regulation.

Many transcription factors incorporate DNA-binding structural motifs which comprise either  $\alpha$  helices or  $\beta$  sheets that bind to the major groove of DNA. Four well-characterized structural motifs are helix-turn-helix, zinc finger, leucine zipper, and helix-loop-helix. Proteins containing these motifs may act alone as monomers or form homo- or heterodimers that interact with DNA.

The helix-turn-helix motif consists of two  $\alpha$  helices connected at a fixed angle by a short

chain of amino acids. One of the helices binds to the major groove. Helix-turn-helix motifs are exemplified by the homeobox motif which is present in homeodomain proteins. These proteins are critical for specifying the anterior-posterior body axis during development and are conserved throughout the animal kingdom. The Antennapedia and Ultrabithorax proteins of Drosophila melanogaster are prototypical homeodomain proteins. (Pabo, C.O. and R.T. Sauer (1992) Ann. Rev. Biochem. 61:1053-1095.)

The zinc finger motif, which binds zinc ions, generally contains tandem repeats of about 30 amino acids consisting of periodically spaced cysteine and histidine residues. Examples of this sequence pattern include the C2H2-type and the C3HC4-type zinc fingers, and the PHD domain. (Lewin, supra ; Aasland, R., et al. (1995) Trends Biochem. Sci 20:56 - 59.) Zinc finger proteins each contain an  $\alpha$  helix and an antiparallel  $\beta$  sheet whose proximity and conformation are maintained by the zinc ion. Contact with DNA is made by the arginine preceding the  $\alpha$  helix and by the second, third, and sixth residues of the  $\alpha$  helix. Variants of the zinc finger motif include poorly defined cysteine-rich motifs which bind zinc or other metal ions. These motifs may not contain histidine residues and are generally nonrepetitive.

The leucine zipper motif comprises a stretch of amino acids rich in leucine which can form an amphipathic  $\alpha$  helix. This structure provides the basis for dimerization of two leucine zipper proteins. The region adjacent to the leucine zipper is usually basic, and upon protein dimerization, is optimally positioned for binding to the major groove. Proteins containing such motifs are generally referred to as bZIP transcription factors.

The helix-loop-helix motif (HLH) consists of a short  $\alpha$  helix connected by a loop to a longer  $\alpha$  helix. The loop is flexible and allows the two helices to fold back against each other and to bind to DNA. The transcription factor Myc contains a prototypical HLH motif.

Most transcription factors contain characteristic DNA binding motifs, and variations on the above motifs and new motifs have been and are currently being characterized. (Faissst, S. and S. Meyer (1992) Nucl. Acids Res. 20:3-26.)

Mutations in transcription factors contribute to oncogenesis. This is likely due to the role of transcription factors in the expression of genes involved in cell proliferation. For example, mutations in transcription factors encoded by proto-oncogenes, such as Fos, Jun, Myc, Rel, and Sp1, may be oncogenic due to increased stimulation of cell proliferation. Conversely, mutations in transcription factors encoded by tumor suppressor genes, such as p53, RB1, and WT1, may be oncogenic due to decreased inhibition of cell proliferation. (Latchman, D. (1995) Gene Regulation: A Eukaryotic Perspective, Chapman and Hall, London, UK, pp 242-255.)

Gene expression is also affected by chromatin-associated proteins. In the nucleus, DNA is

packaged into chromatin, the compact organization of which limits the accessibility of DNA to transcription factors and plays a key role in gene regulation. (Lewin, supra, pp. 409-410.) The compact structure of chromatin is determined and influenced by chromatin-associated proteins such as histones, high mobility group (HMG) proteins, helicases, and chromodomain proteins. There are five classes of histones, H1, H2A, H2B, H3, and H4, all of which are highly basic, low molecular weight proteins. The fundamental unit of chromatin, the nucleosome, consists of 200 base pairs of DNA associated with two copies each of H2A, H2B, H3, and H4. H1 links adjacent nucleosomes. HMG proteins are low molecular weight, non-histone proteins that may play a role in unwinding DNA and stabilizing single-stranded DNA. Helicases, which are DNA-dependent ATPases, unwind DNA, allowing access for transcription factors. Chromodomain proteins play a key role in the formation of highly-compacted, transcriptionally silent heterochromatin.

Much of the regulation of gene expression in eucaryotic cells occurs at the posttranscriptional level. Messenger RNAs (mRNA), which are produced in the cell nucleus from primary transcripts of protein-encoding genes, are processed and transported to the cytoplasm where the protein synthesis machinery is located. RNA-binding proteins are a group of proteins that participate in the processing, editing, transport, localization, and posttranscriptional regulation of mRNAs, and comprise the protein component of ribosomes as well. The RNA-binding activity of many of these proteins is mediated by a series of RNA-binding motifs identified within them. These domains include the RNP motif, the arginine-rich motif, the RGG box, and the KH motif. (Reviewed in Burd, C. G. and Dreyfuss, G. (1994) *Science* 265:615 - 621.) The RNP motif is the most widely found and best characterized of these motifs. The RNP motif is composed of 90-100 amino acids which form an RNA-binding domain and is found in one or more copies in proteins that bind pre-mRNA, mRNA, pre-ribosomal RNA, and small nuclear RNAs. The RNP motif is composed of two short sequences (RNP-1 and RNP-2) and a number of other mostly hydrophobic, conserved amino acids interspersed throughout the motif. (Burd, supra; ExPASy PROSITE document PDOC0030.)

Many neoplastic disorders in humans can be attributed to inappropriate gene expression. Malignant cell growth may result from either excessive expression of tumor promoting genes or insufficient expression of tumor suppressor genes. (Cleary, M.L. (1992) *Cancer Surv.* 15:89-104.) Chromosomal translocations may also produce chimeric loci which fuse the coding sequence of one gene with the regulatory regions of a second unrelated gene. Such an arrangement often results in inappropriate gene transcription. The Wilms tumor suppressor gene product, WT1, is a protein containing a DNA-binding domain consisting of four zinc fingers and a proline-glutamine rich region capable of regulating transcription. (ExPASy PROSITE document PR00049.) Deletions of the WT1 gene, or point mutations which destroy the DNA-binding activity of the protein are associated with development of the pediatric nephroblastoma, Wilms tumor, and Denys-Drash syndrome. (Rauscher,

F.J. (1993) FASEB J. 7:896-903.)

Certain proteins enriched in glutamine are associated with various neurological disorders including spinocerebellar ataxia, bipolar affective disorder, schizophrenia, and autism. (Margolis, R.L. et al. (1997) Human Genetics 100:114-122.) These proteins contain regions with as many as 15 or more consecutive glutamine residues and may function as transcription factors with a potential role in regulation of neurodevelopment or neuroplasticity.

The immune system responds to infection or trauma by activating a cascade of events that coordinate the progressive selection, amplification, and mobilization of cellular defense mechanisms. A complex and balanced program of gene activation and repression is involved in this process.

However, hyperactivity of the immune system as a result of improper or insufficient regulation of gene expression may result in considerable tissue or organ damage. This damage is well documented in immunological responses associated with arthritis, allergens, heart attack, stroke, and infections. (Harrison's Principles of Internal Medicine, 13/e, McGraw Hill, Inc. and Teton Data Systems Software, 1996.) In particular, a zinc finger protein termed Staf50 (for Stimulated trans-acting factor of 50 kDa) is a transcriptional regulator and is induced in various cell lines by interferon-I and -II. Staf50 appears to mediate the antiviral activity of interferon by down-regulating the viral transcription directed by the long terminal repeat promoter region of human immunodeficiency virus type-1 in transfected cells. (Tissot, C. (1995) J. Biol. Chem. 270:14891-14898.)

Furthermore, the generation of multicellular organisms is based upon the induction and coordination of cell differentiation at the appropriate stages of development. Central to this process is differential gene expression, which confers the distinct identities of cells and tissues throughout the body. Failure to regulate gene expression during development could result in developmental disorders.

The discovery of new nucleic-acid binding proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of reproductive, immune, and neurological disorders, and cell proliferative disorders including cancer.

## SUMMARY OF THE INVENTION

The invention features purified polypeptides, protnames, referred to collectively as "ABBR" and individually as "NuABP-1," "NuABP-2," "NuABP-3," "NuABP-4," "NuABP-5," "NuABP-6," "NuABP-7," "NuABP-8," "NuABP-9," "NuABP-10," "NuABP-11," "NuABP-12," "NuABP-13," "NuABP-14," "NuABP-15," "NuABP-16," "NuABP-17," "NuABP-18," "NuABP-19," "NuABP-20," "NuABP-21," "NuABP-22," "NuABP-23," "NuABP-24," "NuABP-25," "NuABP-26," "NuABP-27," "NuABP-28," "NuABP-29," "NuABP-30," "NuABP-31," "NuABP-32," "NuABP-33,"

"NuABP-34," "NuABP-35," "NuABP-36," "NuABP-37," "NuABP-38," "NuABP-39," "NuABP-40," "NuABP-41," "NuABP-42," "NuABP-43," "NuABP-44," "NuABP-45," "NuABP-46," "NuABP-47," "NuABP-48," "NuABP-49," "NuABP-50," "NuABP-51," "NuABP-52," "NuABP-53," "NuABP-54," and "NuABP-55." In one aspect, the invention provides an isolated polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-55.

The invention further provides an isolated polynucleotide encoding a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55. In one alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:56-110.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active  
 5 fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55.

The invention further provides an isolated polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:56-110, b) a naturally occurring  
 10 polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:56-110, c) a polynucleotide sequence complementary to a), or d) a polynucleotide sequence complementary to b). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a  
 15 sample, said target polynucleotide having a sequence of a polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:56-110, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:56-110, c) a polynucleotide sequence complementary to a), or d) a polynucleotide sequence complementary to b). The method comprises a)  
 20 hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe  
 25 comprises at least 30 contiguous nucleotides. In another alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a pharmaceutical composition comprising an effective amount of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an  
 30 amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, and a pharmaceutically acceptable excipient. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional NuABP,  
 35 comprising administering to a patient in need of such treatment the pharmaceutical composition.



The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional NuABP, comprising administering to a patient in need of such treatment the pharmaceutical composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional NuABP, comprising administering to a patient in need of such treatment the pharmaceutical composition.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:56-110, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

#### BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding NuABP.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of NuABP.

Table 3 shows selected fragments of each nucleic acid sequence: the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding NuABP were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze NuABP, along with applicable descriptions, references, and threshold parameters.

### DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

### DEFINITIONS

"NuABP" refers to the amino acid sequences of substantially purified NuABP obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of

NuABP. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of NuABP either by directly interacting with NuABP or by acting on components of the biological pathway in which NuABP participates.

An "allelic variant" is an alternative form of the gene encoding NuABP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding NuABP include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as NuABP or a polypeptide with at least one functional characteristic of NuABP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding NuABP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding NuABP. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent NuABP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of NuABP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophobicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of NuABP. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of NuABP either by directly interacting with NuABP or by acting on components of the biological pathway in which NuABP participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind NuABP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition containing a nucleic acid sequence which is complementary to the "sense" strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and to block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic NuABP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" and "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" bonds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity

between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding NuABP or fragments of NuABP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using the XL-PCR kit (Perkin-Elmer, Norwalk CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of one or more Incyte Clones and, in some cases, one or more public domain ESTs, using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that, when made, least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
25	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
30	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
35	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
40	Thr	Ser, Val

Trp  
Tyr  
Val

Phe, Tyr  
His, Phe, Trp  
Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "fragment" is a unique portion of NuABP or the polynucleotide encoding NuABP which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:56-110 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:56-110, for example, as distinct from any other sequence in the same genome. A fragment of SEQ ID NO:56-110 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:56-110 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:56-110 and the region of SEQ ID NO:56-110 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-55 is encoded by a fragment of SEQ ID NO:56-110. A

fragment of SEQ ID NO:1-55 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-55. For example, a fragment of SEQ ID NO:1-55 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-55. The precise length of a fragment of SEQ ID NO:1-55 and the region of SEQ ID NO:1-55 to which the  
 5 fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a  
 10 target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of  
 15 reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the  
 20 substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps  
 25 in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of  
 30 molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent  
 35 similarity" between aligned polynucleotide sequence pairs.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at

5 <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The

10 "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

*Reward for match: 1*

*Penalty for mismatch: -2*

*Open Gap: 5 and Extension Gap: 2 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 11*

*Filter: on*

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at

25 least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode

30 similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a

35 standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some



alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the hydrophobicity and acidity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) with blastp set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

*Open Gap: 11 and Extension Gap: 1 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 3*

*Filter: on*

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term "humanized antibody" refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific

hybridization is an indication that two nucleic acid sequences share a high degree of identity. Specific

hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for

annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Generally, such wash temperatures are selected to be about 5°C to 20°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating  $T_m$  and conditions for nucleic acid hybridization are well known and can be found in Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour.

Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g.,  $C_{0t}$  or  $R_{0t}$  analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide

sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect  
5 cellular and systemic defense systems.

The term "microarray" refers to an arrangement of distinct polynucleotides on a substrate.

The terms "element" and "array element" in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

The term "modulate" refers to a change in the activity of NuABP. For example, modulation  
10 may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of NuABP.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the  
15 antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences may be in close proximity or contiguous and,  
20 where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation,  
25 and may be pegylated to extend their lifespan in the cell.

"Probe" refers to nucleic acid sequences encoding NuABP, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

30 "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous  
35 nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also

be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

5       Methods for preparing and using probes and primers are described in the references, for example Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis et al., 1990, PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be  
10       derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

      Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to  
15       5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer  
20       selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The  
25       PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments  
30       identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

      A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence  
35       that is made by an artificial combination of two or more otherwise separated segments of sequence.

This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding NuABP, or fragments thereof, or NuABP itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

"Transformation" describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected

based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

## THE INVENTION

The invention is based on the discovery of new human nucleic-acid binding proteins (NuABP), the polynucleotides encoding NuABP, and the use of these compositions for the diagnosis, treatment, or prevention of reproductive, immune, and neurological disorders, and cell proliferative disorders including cancer.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding

NuABP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOS) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each NuABP were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their  
5 corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. The Incyte clones in column 5 were used to assemble the consensus nucleotide sequence of each NuABP and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each  
10 polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows identification or homologous sequences as identified by BLAST analysis; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence  
15 homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding NuABP. The first column of Table 3 lists the nucleotide SEQ ID NOS. Column 2 lists fragments of the nucleotide sequences of column 1. These  
20 fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:56-110 and to distinguish between SEQ ID NO:56-110 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express NuABP as a fraction of total tissues expressing NuABP. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing NuABP as a fraction of total tissues expressing NuABP. Of particular note is the expression of SEQ  
25 ID NO:83 and SEQ ID NO:110 in neurological tissue. About 53% of the cDNA libraries expressing SEQ ID NO:83 are derived from neurological tissue. Furthermore, SEQ ID NO:110 expression is detected exclusively in a cDNA library derived from brain tissue afflicted with Huntington's disease. Column 5 lists the vectors used to subclone each cDNA library.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries  
30 from which cDNA clones encoding NuABP were isolated. Column 1 references the nucleotide SEQ ID NOS, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

Fragments of the nucleotide sequences encoding NuABP are useful, for example, in hybridization or amplification technologies to identify SEQ ID NOS:56-110 and to distinguish  
35 between SEQ ID NOS:56-110 and related polynucleotide sequences. The polypeptides encoded by

these fragments are useful, for example, as immunogenic peptides.

The invention also encompasses NuABP variants. A preferred NuABP variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the NuABP amino acid sequence, and which contains at least one functional or structural characteristic of NuABP.

The invention also encompasses polynucleotides which encode NuABP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:56-110, which encodes NuABP.

The invention also encompasses a variant of a polynucleotide sequence encoding NuABP. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding NuABP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:56-110 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:56-110. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of NuABP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding NuABP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring NuABP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode NuABP and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring NuABP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding NuABP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding NuABP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.



The invention also encompasses production of DNA sequences which encode NuABP and NuABP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce

5 mutations into a sequence encoding NuABP or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:56-110 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

10

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler

15

20 (Perkin-Elmer). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Perkin-Elmer), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

25

The nucleic acid sequences encoding NuABP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic

30 DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent

35 to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al.

(1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060).

5 Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in  
10 length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T)  
15 library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate  
20 software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be  
25 present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode NuABP may be cloned in recombinant DNA molecules that direct expression of NuABP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a  
30 functionally equivalent amino acid sequence may be produced and used to express NuABP.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter NuABP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic  
35 oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-

mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In another embodiment, sequences encoding NuABP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; and Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.)

Alternatively, NuABP itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Perkin-Elmer). Additionally, the amino acid sequence of NuABP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY.)

In order to express a biologically active NuABP, the nucleotide sequences encoding NuABP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding NuABP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding NuABP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding NuABP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) Results Probl. Cell Differ. 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding NuABP and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques,

and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

5 A variety of expression vector/host systems may be utilized to contain and express sequences encoding NuABP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

10 In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding NuABP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding NuABP can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPT01 (Life Technologies). Ligation of sequences encoding NuABP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of NuABP are needed, e.g. for the production of antibodies, vectors which direct high level expression of NuABP may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

25 Yeast expression systems may be used for production of NuABP. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

30 Plant systems may also be used for expression of NuABP. Transcription of sequences encoding NuABP may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al.

(1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.)

These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

5 In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding NuABP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses NuABP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

10 Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of NuABP in cell lines is preferred. For example, sequences encoding NuABP can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *apr* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which

alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP: Clontech),  $\beta$  glucuronidase and its substrate  $\beta$ -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding NuABP is inserted within a marker gene sequence, transformed cells containing sequences encoding NuABP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding NuABP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding NuABP and that express NuABP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of NuABP using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on NuABP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding NuABP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding NuABP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase

such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding NuABP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode NuABP may be designed to contain signal sequences which direct secretion of NuABP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding NuABP may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric NuABP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of NuABP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the NuABP encoding sequence and the heterologous protein sequence, so that NuABP may be cleaved away from the heterologous moiety following purification.

Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled NuABP may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, <sup>35</sup>S-methionine.

Fragments of NuABP may be produced not only by recombinant means, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra, pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the ABI 431A peptide synthesizer (Perkin-Elmer). Various fragments of NuABP may be synthesized separately and then combined to produce the full length molecule.

#### THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of NuABP and nucleic-acid binding proteins. In addition, the expression of NuABP is closely associated with proliferative, neuronal, inflamed, and cancerous tissues and tissues of the reproductive system. Therefore, NuABP appears to play a role in reproductive, immune, and neurological disorders, and cell proliferative disorders including cancer. In the treatment of disorders associated with increased NuABP expression or activity, it is desirable to decrease the expression or activity of NuABP. In the treatment of disorders associated with decreased NuABP expression or activity, it is desirable to increase the expression or activity of NuABP.

Therefore, in one embodiment, NuABP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of NuABP. Examples of such disorders include, but are not limited to, a reproductive disorder such as disorders of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, disruptions of the estrous cycle, disruptions of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, endometrial and ovarian tumors, uterine fibroids, autoimmune disorders, ectopic pregnancies, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; disruptions of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia; an immune disorder such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, bursitis,



cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis. Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel syndrome,

5 episodic lymphopenia with lymphocytotoxins, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis,

10 Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, trauma, and hematopoietic cancer including lymphoma, leukemia, and myeloma; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy,

15 retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Strausler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis,

20 tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies,

25 myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, and Tourette's disorder; and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis,

30 primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

35 In another embodiment, a vector capable of expressing NuABP or a fragment or derivative

thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of NuABP including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified NuABP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of NuABP including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of NuABP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of NuABP including, but not limited to, those listed above.

In a further embodiment, an antagonist of NuABP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of NuABP. Examples of such disorders include, but are not limited to, those reproductive, immune, and neurological disorders, and cell proliferative disorders including cancer, described above. In one aspect, an antibody which specifically binds NuABP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express NuABP.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding NuABP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of NuABP including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of NuABP may be produced using methods which are generally known in the art. In particular, purified NuABP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind NuABP. Antibodies to NuABP may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with NuABP or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to

increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

5 It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to NuABP have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of NuABP amino  
10 acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to NuABP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma  
15 technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate  
20 antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce NuABP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be  
25 generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA  
30 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for NuABP may also be generated. For example, such fragments include, but are not limited to, F(ab')<sub>2</sub> fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and  
35 easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D.

et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between NuABP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering NuABP epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for NuABP. Affinity is expressed as an association constant,  $K_a$ , which is defined as the molar concentration of NuABP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The  $K_a$  determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple NuABP epitopes, represents the average affinity, or avidity, of the antibodies for NuABP. The  $K_a$  determined for a preparation of monoclonal antibodies, which are monospecific for a particular NuABP epitope, represents a true measure of affinity. High-affinity antibody preparations with  $K_a$  ranging from about  $10^9$  to  $10^{12}$  L/mole are preferred for use in immunoassays in which the NuABP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with  $K_a$  ranging from about  $10^6$  to  $10^7$  L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of NuABP, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume 1: A Practical Approach, IRL Press, Washington, DC; Liddell, J.E. and Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of NuABP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding NuABP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding NuABP may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding NuABP. Thus, complementary molecules or

fragments may be used to modulate NuABP activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding NuABP.

5 Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding NuABP. (See, e.g., Sambrook, supra; Ausubel, 1995, supra.)

10 Genes encoding NuABP can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding NuABP. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more  
15 with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding NuABP. Oligonucleotides derived from the transcription  
20 initiation site, e.g., between about positions -10 and +10 from the start site, may be employed. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber,  
25 B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme  
30 molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding NuABP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA,  
35 GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides,

corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding NuABP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of NuABP, antibodies to NuABP, and mimetics, agonists, antagonists, or inhibitors of NuABP. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered

to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in

aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily  
5 injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

10 For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

15 The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acids. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a  
20 pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of NuABP, such labeling would include amount, frequency, and method of administration.

25 Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of  
30 administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example NuABP or fragments thereof, antibodies of NuABP, and agonists, antagonists or inhibitors of NuABP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be  
35 determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such



WO 00/44900

PCT/US00/02237

as by calculating the  $ED_{50}$  (the dose therapeutically effective in 50% of the population) or  $LD_{50}$  (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the  $LD_{50}/ED_{50}$  ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the  $ED_{50}$  with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1  $\mu\text{g}$  to 100,000  $\mu\text{g}$ , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art.

Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

## DIAGNOSTICS

In another embodiment, antibodies which specifically bind NuABP may be used for the diagnosis of disorders characterized by expression of NuABP, or in assays to monitor patients being treated with NuABP or agonists, antagonists, or inhibitors of NuABP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for NuABP include methods which utilize the antibody and a label to detect NuABP in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring NuABP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of NuABP expression.

Normal or standard values for NuABP expression are established by combining body fluids or cell

extracts taken from normal mammalian subjects, for example, human subjects, with antibody to NuABP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of NuABP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values.

- 5 Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding NuABP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of NuABP may be correlated  
10 with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of NuABP, and to monitor regulation of NuABP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding NuABP or closely related molecules may be used to identify nucleic acid sequences which encode NuABP. The specificity of the probe, whether it is  
15 made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding NuABP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50%  
20 sequence identity to any of the NuABP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:56-110 or from genomic sequences including promoters, enhancers, and introns of the NuABP gene.

Means for producing specific hybridization probes for DNAs encoding NuABP include the cloning of polynucleotide sequences encoding NuABP or NuABP derivatives into vectors for the  
25 production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as  $^{32}\text{P}$  or  $^{35}\text{S}$ , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

30 Polynucleotide sequences encoding NuABP may be used for the diagnosis of disorders associated with expression of NuABP. Examples of such disorders include, but are not limited to, a reproductive disorder such as disorders of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, disruptions of the estrous cycle, disruptions of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, endometrial and ovarian  
35 tumors, uterine fibroids, autoimmune disorders, ectopic pregnancies, and teratogenesis; cancer of the

breast, fibrocystic breast disease, and galactorrhea; disruptions of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia; an immune disorder such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel syndrome, episodic lymphopenia with lymphocytotoxins, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, trauma, and hematopoietic cancer including lymphoma, leukemia, and myeloma; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, and Tourette's disorder; and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal

hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. The polynucleotide sequences encoding NuABP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered NuABP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding NuABP may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding NuABP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding NuABP in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of NuABP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding NuABP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the

development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

5 Additional diagnostic uses for oligonucleotides designed from the sequences encoding NuABP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding NuABP, or a fragment of a polynucleotide complementary to the polynucleotide encoding NuABP, and will be employed under optimized conditions for identification of a specific gene or  
10 condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

Methods which may also be used to quantify the expression of NuABP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplax, C.  
15 et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the  
20 polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

25 Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

30 In another embodiment of the invention, nucleic acid sequences encoding NuABP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single  
35 chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price,

C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the  
 5 Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding NuABP on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

10 In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides  
 15 valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be  
 20 used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, NuABP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a  
 25 solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between NuABP and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are  
 30 synthesized on a solid substrate. The test compounds are reacted with NuABP, or fragments thereof, and washed. Bound NuABP is then detected by methods well known in the art. Purified NuABP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

35 In another embodiment, one may use competitive drug screening assays in which neutralizing

antibodies capable of binding NuABP specifically compete with a test compound for binding NuABP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with NuABP.

In additional embodiments, the nucleotide sequences which encode NuABP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. 60/117,905 and U.S. Ser. No. 60/117,904, are hereby expressly incorporated by reference.

## EXAMPLES

### I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP

vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPT1 plasmid (Life Technologies), or pINCY (Incyte Pharmaceuticals, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 $\alpha$ , DH10B, or ElectroMAX DH10B from Life Technologies.

## II. Isolation of cDNA Clones

Plasmids were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

## III. Sequencing and Analysis

cDNA sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Perkin-Elmer) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the



5 disclosed in Example V.

10

20

35

NO:56-110. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

#### IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel, 1995, *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding NuABP occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

#### V. Extension of NuABP Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:56-110 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target

sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing  $Mg^{2+}$ ,  $(NH_4)_2SO_4$ , and  $\beta$ -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100  $\mu$ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5  $\mu$ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5  $\mu$ l to 10  $\mu$ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min;

Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing  
 5 primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

In like manner, the nucleotide sequences of SEQ ID NO:56-110 are used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for such extension, and an appropriate genomic library.

#### 10 VI. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:56-110 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06  
 15 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250  $\mu$ Ci of [ $\gamma$ -<sup>32</sup>P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10<sup>7</sup> counts per minute of the labeled probe is used in a typical membrane-based  
 20 hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature  
 25 under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

#### VII. Microarrays

A chemical coupling procedure and an ink jet device can be used to synthesize array  
 30 elements on the surface of a substrate. (See, e.g., Baldeschweiler, *supra*.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and  
 35 patterns of fluorescence. The degree of complementarity and the relative abundance of each probe

which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

### VIII. Complementary Polynucleotides

Sequences complementary to the NuABP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring NuABP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of NuABP. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the NuABP-encoding transcript.

### IX. Expression of NuABP

Expression and purification of NuABP is achieved using bacterial or virus-based expression systems. For expression of NuABP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*lac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express NuABP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of NuABP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding NuABP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to

infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, NuABP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from NuABP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified NuABP obtained by these methods can be used directly in the following activity assay.

#### **X. Demonstration of NuABP Activity**

NuABP activity is measured by its ability to stimulate transcription of a reporter gene (Liu, H.Y. et al. (1997) EMBO J. 16(17):5289-5298.) The assay entails the use of a well characterized reporter gene construct, LexA<sub>op</sub>-LacZ, that consists of LexA DNA transcriptional control elements (LexA<sub>op</sub>) fused to sequences encoding the E. coli LacZ enzyme. The methods for constructing and expressing fusions genes, introducing them into cells, and measuring LacZ enzyme activity, are well known to those skilled in the art. Sequences encoding NuABP are cloned into a plasmid that directs the synthesis of a fusion protein, LexA-NuABP, consisting of NuABP and a DNA binding domain derived from the LexA transcription factor. The resulting plasmid, encoding a LexA-NuABP fusion protein, is introduced into yeast cells along with a plasmid containing the LexA<sub>op</sub>-LacZ reporter gene. The amount of LacZ enzyme activity associated with LexA-NuABP transfected cells, relative to control cells, is proportional to the amount of transcription stimulated by the NuABP.

#### **XI. Functional Assays**

NuABP function is assessed by expressing the sequences encoding NuABP at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT (Life Technologies) and pCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10  $\mu$ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell

line, using either liposome formulations or electroporation. 1-2  $\mu$ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green

- 5 Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA
- 10 with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in
- 15 flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of NuABP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding NuABP and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected

- 20 cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding NuABP and other genes of interest can be analyzed by northern analysis or microarray techniques.

## **XII. Production of NuABP Specific Antibodies**

- 25 NuABP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the NuABP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is

- 30 synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St.

- 35 Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase

immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-NuABP activity by, for example, binding the peptide or NuABP to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

5 **XIII. Purification of Naturally Occurring NuABP Using Specific Antibodies**

Naturally occurring or recombinant NuABP is substantially purified by immunoaffinity chromatography using antibodies specific for NuABP. An immunoaffinity column is constructed by covalently coupling anti-NuABP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is  
10 blocked and washed according to the manufacturer's instructions.

Media containing NuABP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of NuABP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/NuABP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such  
15 as urea or thiocyanate ion), and NuABP is collected.

**XIV. Identification of Molecules Which Interact with NuABP**

NuABP, or biologically active fragments thereof, are labeled with <sup>125</sup>I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled NuABP, washed,  
20 and any wells with labeled NuABP complex are assayed. Data obtained using different concentrations of NuABP are used to calculate values for the number, affinity, and association of NuABP with the candidate molecules.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention.  
25 Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.



TABLE 1

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	56	025733	SPLNFET01	025733H1 and 025733X30TD2 (SPLNFET01), 1519809F1 and 1519809T1 (BLADTUT04), 1526288T6 (OCMCL5T01), 1595557X16C1 and 1595557X19C1 (BRATNOT14), 1903359F6 (OVARNOT07), 2417225F6 (HNT3AZT01), 4283542H1 (LIVDIR01)
2	57	079702	SYNORAB01	002190H1 (O93TNOT01), 079479F1 and 079702H1 (SYNORAB01), 125223T6 (LUNGNOT01), 371006R6 (LUNGNOT02), 2129460T6 (KIDNNOT05), 2999366H1 (LYNNOT06), 3031905F6 (TYRNOT05), 4949863H1 (SINTNOT25)
3	58	116208	KIDNNOT01	116208H1 and 116208R1 (KIDNNOT01), 2293058R6 (BRAINNOT1), 3731418H1 and 3731418T6 (SMCCNQ03)
4	59	179261	PLACNOB01	179261H1 (PLACNOB01), 3666231F6 and 3666231T6 (PANCNOT16)
5	60	259161	HNT2RAT01	259161H1 (HNT2RAT01), 1005021R6 (BRSTNOT03), 2634660H1 (COLNUT015), 2894333H1 (KIDNUT14), 2924845H1 (SININOT04), 3659440H1 (ENDPNOT02), SBMA02955F1, SBMA03577F1, SBMA01445F1, SBMA00985F1, SBMA01499F1
6	61	320087	EOSIHET02	016657F1 (HUELPEB01), 320087H1 (EOSIHET02), 824110R1 (PROSNOT06), 987467H1 (LVENLPT03), 1235752F1 (LUNGFT03), 1361280F1 (LUNGNOT12), 1389740H1 (EOSINOT01), 1534332F1 (SPLNNOT04), 1813754F6 (SKINBIT01), 4184915H1 (BRSTNOT31), 5306522H1 (MONOTXT02)
7	62	491271	HNT2AGT01	363816X3 and 363816X9 (PROSNOT01), 491271F1, 491271H1 and 491271T6 (HNT2AGT01), 967354R6, 967354X15 and 967354X27 (BRSTNOT05), 2733444H1 (OVARUT04)
8	63	585172	PROSNOT02	395188R6 (TWL2DT01), 585172H1 (PROSNOT02), 864269T1 (BRATUT03), 1417965F1 (KIDNNOT09)
9	64	615200	COLNUT02	615200H1 and 615200R6 (COLNUT02), 1213980R1 (BRSTTUT01), SBPA02731D1, SBPA00184D1
10	65	997067	KIDNUT01	125981X3 (LUNGNOT01), 997067H1, 997067R6 and 997067T6 (KIDNUT01), 1448201H1 (PLACNOT02), 1663447H1 (BRSTNOT09), 1889314H1 (BLATUT07), 1918706H1 (PROSNOT06), 2699956H1 (OVARUT10), 2702585H1 (FIBENOT10), 2900479H1 (DRGNOT01), 3595727T6 (FIBENOT11), 4109131H1 (BRANUT01)
11	66	144326 2	THYRNOT03	1257005F1 (MENITUT03), 1443262H1 (THYRNOT03), 1618906F6 (BRATUT12), 2474133T6 (THP1NOT03), 3594075H1 (FIBPNOT11), 4914442H1 (LIVFAT05)

TABLE 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
12	67	152164 8	BLADTUT04	292172H1 (TMR3DT01), 819925R6 (KERANOT02), 1353913H1 (LATRTUT02), 1521648H1 and 1522364F1 (BLADTUT04), 1963178H1 (BRSTNOT04), 2342505F6 (TESTTUT02), 4899970H1 (OVARDT01), 5043585H2 (PLACFER01)
13	68	168549 4	PROSNOT15	903980X13, 903980X14 and 903980X17 (COLNNOT07), 1685494H1 (PROSNOT15), 4164127T6 (BRSTNOT32)
14	69	173082 9	BRSTTUT08	116146R1 (KIDNNOT01), 836856R1 (PROSNOT07), 1730829H1, 1730829X11C1, 1730829X12C1 and 1730829X13C1 (BRSTTUT08), 1959889R6 (BRSTNOT04), 2188079H1 (PROSNOT26), 3384625H1 (ESGNOT04)
15	70	186464 1	PROSNOT19	1844372H1 (COLNNOT08), 1864641F6 and 1864641H1 (PROSNOT19), 3090702T6 (BRSTNOT19), 3411665H1 (BRSTTUS08), 5152366H1 (HEARFET03), 5166179H1 (MUSCDMT01)
16	71	244460 4	THPINOT03	1506659F1 (BRAITUT07), 1532034F1 (SPLNNOT04), 2444604H1 (THPINOT03)
17	72	244500 8	THPINOT03	605598X12 (BRSTTUT01), 628644H1 (KIDNNOT05), 732124R1 (LUNGNOT03), 819194R1 (KERANOT02), 1259467H1 (MENITUT03), 1363205F6 (LUNGNOT12), 1901312T6 (BLADTUT06), 2445008H1 (THPINOT03), 2681125H1 (SINIUCT01)
18	73	257246 2	HIPOAZT01	396323R6 (PITUNOT02), 863622H1 (BRAITUT03), 1848956F6 and 1848956T6 (LUNGFET03), 2345947H1 (TESTTUT02), 2396384F6 (THPIAZT01), 2572462H1 (HIPOAZT01), 2650980F6 (LUNGTUT12), 2814325H1 (OVARNOT10), 5076051H1 (COLCTUT03)
19	74	257289 2	HIPOAZT01	030596X15R1 (THP1NOB01), 539564X11 (LUNGNOT02), 1275514F1 and 1275514T6 (TESTTUT02), 2112383H1 (BRAITUT03), 2572892H1 (HIPOAZT01), 2986519H1 (CARGDT01)
20	75	278567 4	BRSTNOT13	261399H1 (HNT2ACT01), 1274739F1 (TESTTUT02), 2785674H1 (BRSTNOT13)
21	76	279747 9	NPOLNOT01	302614X13 (TESTNOT04), 2797479H1 (NPOLNOT01), SAIA02597F1, SAIA02739F1, SAIA02537F1
22	77	296064 0	ADRENOT09	027211R1, 027211X1 and 027211X3 (SPLNFET01), 1401538F6 (BRAITUT08), 2496984F6 (ADRENUT05), 2960640H1 (ADRENUT09), 3211036T6 (BLADNOT08)
23	78	345405 1	SPLNNOT11	279331R6 (LIVANOT02), 2515972T6 and 2516010T6 (LIVRTUT04), 2910726F6 (KIDNTUT15), 3454051H1 (SPLNNOT11)

TABLE 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
24	79	351064 0	CONCNOT01	556354H1 (MPHGLPT02), 581085H1 (BRAVXT05), 990636R6 (COLANNOT11), 1799185F6 and 1799185T6 (COLNNOT27), 3510640H1 (CONCNOT01), 4326648H1 (TLYMUT01), SBA01095F1
25	80	381508 3	TONSNOT03	2026951R6 and 2026951T6 (KERANOT02), 2300211R6 (BRSTNOT05), 2505283F6 (CONUTUT01), 3187267R6 (THYMNON04), 3815083H1 and 3815083T6 (TONSNOT03)
26	81	398845 7	LUNGNON03	609622R6 (COLNNOT01), 1710465F6 (PROSNOT16), 3988457H1 (LUNGNON03), SAA000089F1, SAA03055F1, SAQB00279F1
27	82	131890	BMARNOT02	131890H1 (BMARNOT02), 131890T6 (BMARNOT02), 132849R6 (BMARNOT02), 3357071F6 (PROSTUT16)
28	83	238642	SINTNOT02	238642H1 (SINTNOT02), 1620593F6 (BRAITUT13), 1620593H1 (BRAITUT13), 1620593T6 (BRAITUT13), 2534087F6 (BRAITUT18)
29	84	669862	CRBLNOT01	347231X7 (THYMNOT02), 669862H1 (CRBLNOT01), 2244458R6 (HIFONON02), 2244458T6 (HIFONON02), 2622610T6 (KERANOT02), 3536262H1 (KIDNNOT25), 4204212H1 (BRAITUT29)
30	85	100366 3	BRSTNOT03	850478T1 (NGANNOT01), 1003663H1 (BRSTNOT03), 1252179F2 (LUNGFET03), 1293336F1 (PGANNOT03), 1813002F6 (PROSTUT12), 2101974R6 (BRAITUT02)
31	86	143255 7	BEPINON01	1314545F6 (BLADUT02), 1432557H1 (BEPINON01), 1443311R1 (THYRNUT03), 1705738F6 (DUONOT02), 2182184F6 (SININOT01)
32	87	144177 0	THYRNUT03	035105R1 (HUVENOB01), 1441770H1 (THYRNUT03), 1500943F1 (SINTBST01), 2542840H1 (UTRSNCT11), 4533672H1 (OVARNOT12)
33	88	145668 4	COLNFET02	1456684F6 (COLNFET02), 1456684H1 (COLNFET02), 1456684T6 (COLNFET02), 1992143H1 (CORPNOT02), 2687476F6 (LUNGNUT02), 3139175F6 (SMCCNOT02), 4746319H1 (SMCRUNT01)
34	89	160291 6	BLADNOT03	3397976X305D2 (UTRSNCT16)
35	90	162881 6	COLNNOT23	999017R6 (KIDNNOT01), 1342490T1 (COLNUT03), 1421981F1 (KIDNNOT09), 1628816H1 (COLNNOT23), 2176832F6 (ENDCNOT03), 2451404F6 (ENDANOT01), 1968191H1 (BRSTNOT04), 1968191T6 (BRSTNOT04), 2752967R6 (THPIAZ508), 3281163T6 (STONFET02), 3748607H1 (UTRSNCT18)
36	91	196819 1	BRSTNOT04	
37	92	205206 1	LIVRFET02	003803X8 (HMCINOT01), 027044X1 (SPLNFET01), 027044X101 (SPLNFET01), 2052061H1 (LIVRFET02), 3931936F6 (PROSTUT09)

TABLE 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
38	93	205620 7	BEPINOT01	071525F1 (PLACNOB01), 162720R1 (ADENTN01), 270498H1 (HNT2NOT01), 1477853T1 (CORPNOT02), 1931058F6 (COLNUTUT03), 2056207H1 (BEPINOT01), 2056207X1I1 (BEPINOT01), 2231050F6 (PROSNOT16), 2420063X309D4 (SCORNON02), 3424630H1 (BRSTN01), 3873760F6 (HEARNOT06), 3873760T6 (HEARNOT06), SBOA00627D1, SCJA02192V1, SCJA02089V1
				399584R6 (PITUNOT02), 399584T6 (PITUNOT02), 1649058F6 (PROSTUT09), 1902809F6 (OVARNOT07), 2101803H1 (BRAITUT02), 2101803R6 (BRAITUT02), 3098623H1 (CERVNOT03)
				948628R1 (PANCNOT05), 1209447T1 (BRSTNOT02), 1814624F6 (PROSNOT20), 2112362H1 (BRAITUT03), 2945621H1 (BRAITUT23), 3285663H1 (HEAONOT05), 3526403H1 (ESOGTUN01), 5032729H1 (ENDIUNOT1), 5099417H1 (PROSTUS20)
				487994R6 (HNT2AGT01), 952855R1 (SCORNON01), 952855T1 (SCORNON01), 2117346H1 (BRSTTUT02), 2458342F6 (ENDANOT01), 2731585H1 (OYARTUT04), 3475780H1 (LUNGNOT27), 3538525F6 (SEMNUT04), 2119917H1 (BRSTTUT02), 2791421F6 (COLNUT16), 2794083F6 (COLNUT16), 5006921H1 (STOMNOT08)
				484031H1 (HNT2RAT01), 617559F1 (PGANNOT01), 617559R1 (PGANNOT01), 1575977F1 (LUNODNOT03), 2123456H1 (BRSTNOT07), 2958712H1 (ADRENUT09), 3764961H1 (BRSTNOT24)
44	99	214879 2	BRAINOT09	1732781F6 (BRSTTUT08), 2050885F6 (LIVREFET02), 2148792H1 (BRAINOT09), 2590822H1 (LUNGNOT22), 2972369T6 (HEAONOT02), SBOA00396D1, SBOA03678D1, SBOA02120D1, SBOA03269D1
				1720187X16C1 (BLADNOT06), 2751943H1 (THPIAZS08), 3492378H1 (ADRETUT07)
46	101	312891 3	LUNGUT12	2551859F6 (LUNGUT06), 3128913H1 (LUNGUT12), SBOA01861F1, SBOA02298F1, SBOA01013F1, SBOA02403F1, SBOA01362F1
				154741R6 (THPIPLB02), 155904R6 (THPIPLB02), 1233933T6 (LUNGFEFF03), 157816R1 (THPIPLB02), 979920H1 (TONGUT01), 1233933T6 (LUNGFEFF03), 1850777F6 (URETUT01), 2445017F6 (THPIPLB03), 3282941H1 (HEAONOT05), 3341633H1 (SPINNOT09), 3517140H1 (LUNGNON03)
48	103	328665 6	HEAONOT05	898123H1 (BRSTNOT05), 3286656H1 (HEAONOT05), 3641429T6 (LUNGNOT30), 3657668F6 (ENDPNOT02)

TABLE 1 (cont.)

Protein SEQ ID No:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
49	104	349080 2	EPIGNOT01	2238441H1 (PANCTUT02), 2700133F6 (OVARUT10), 2700133T6 (OVARUT10), 3490225H1 (EPIGNOT01), 3490802H1 (EPIGNOT01), 4822929H1 (PROSTUT17), 2130284H1 (KIDNNOT05), 3507366H1 (CONCNOT01), 3557087F6 (LUNGNOT31), 4241774H1 (SYNDIT01)
50	105	350736 6	CONCNOT01	
51	105	357306 0	BRONNOT01	3573060F6 (BRONNOT01), 3573060H1 (BRONNOT01), 3573060T6 (BRONNOT01), 3867263H1 (BRAITUT07), 5013346H1 (BRAXNOT03)
52	107	357366 1	BRONNOT01	3028034F6 (HEARFET02), 3152642H1 (ADREN004), 3573661F6 (BRONNOT01), 3573661H1 (BRONNOT01), 3577568F6 (BRONNOT01)
53	108	363342 2	LIVRN0T03	033412R6 (THPINOB01), 074123F1 (THPIPER01), 263241H1 (HNT2AGT01), 748567R1 (BRAITUT01), 1292088T1 (PGANNOT03), 1517449T1 (PANCTUT01), 3633422H1 (LIVRN0T03)
54	109	399337 7	LUNGNON03	3003233H1 (TYLNNOT06), 3993377H1 (LUNGNON03), 3993377T6 (LUNGNON03), 4251662F6 (BRADDIR01), SBSA02001V1
55	110	471793 6	BRAIHT02	4717936F6 (BRAIHT02), 4717936H1 (BRAIHT02), 4717936T6 (BRAIHT02)

TABLE 2

Polypeptide Seq ID No:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
1	754	S6 T15 S22 S26		L435 - P474 PFAM	DNA polymerase	PFAM, BLOCKS, MOTIFS
		S28 S30 S60 Y84				
		S98 S102 S103				
		S112 S121 S146				
2	593	T166 T183 T184		E102 - A157 Q201 - C252	PHD finger DNA binding protein (GI 3342452)	BLAST, PFAM, MOTIFS
		S231 Y253 S303				
		T304 S308 T327				
		T361 T393 T394				
3	534	T399 T448 S496		I17 - D57 PFAM	DNA polymerase	PFAM, BLOCKS, MOTIFS
		Y583 S586 T608				
		S635 S672 S673				
		S682 S691 S711				
4	255	S713 S751		Y58 - C86 F141 - H163 Y169 - H191 Y197 - H219 Y225 - H247 PFAM	C2H2-type zinc finger protein (GI 498721)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
		S32 Y62 T65 S83				
		S141 T159 T160				
		S185 S254 S288				
5	593	T311 S314 S333		I17 - D57 PFAM	DNA polymerase	PFAM, BLOCKS, MOTIFS
		T368 S380 S401				
		T410 S426 S452				
		S461 S479 T483				
6	534	T485 T576		Y58 - C86 F141 - H163 Y169 - H191 Y197 - H219 Y225 - H247 PFAM	C2H2-type zinc finger protein (GI 498721)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
		T39 S52 S301 T344				
		S373 T404 S425				
		S438 S439 S440				
7	534	S473 T480 S490		Y58 - C86 F141 - H163 Y169 - H191 Y197 - H219 Y225 - H247 PFAM	C2H2-type zinc finger protein (GI 498721)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
		T527				
		S10 S26 Y35 Y113				
		S149 T168 Y169				
8	255	T248		Y58 - C86 F141 - H163 Y169 - H191 Y197 - H219 Y225 - H247 PFAM	C2H2-type zinc finger protein (GI 498721)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS

TABLE 2 (cont.)

Polyypeptide Seq ID No:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
5	562	T29 T43 S76 S142 S165 S202 T214 Y302 S305 Y349 S385 S500 T526 S527		Y235 - D312 PFAM	DNA helicase (GI 531243; SEQ ID NO:113)	BLAST, PFAM, BLOCKS, MOTIFS
6	432	S33 S58 T166 T172 S197 T230 T261 S275 S286 S290 S298 S338 T362 S376 T407 T409 T419		E329 - A355 BLOCKS	CCAAT-box-binding transcription factor	BLOCKS, MOTIFS
7	799	T24 S33 T43 S73 T88 S91 S110 S147 T219 S262 S323 Y380 S532 S586 S756 T795		H250 - H291 Y324 - H346 Y253 - H374 Y380 - H402 PFAM	C2H2-type zinc finger protein (GI 498727)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
8	137	S3 T38 S74 S75 S118		R85 - L97 PFAM	BTB domain/C2H2-type zinc finger protein	PFAM, PRINTS, MOTIFS
9	230	T178 S187			sirtuin type 3 (GI 5225322)	BLAST, MOTIFS
10	446	T3 S28 S32 T52 T94 T96 S135 S143 T159 T165 S171 S433		H200 - H222 Y228 - H250 Y256 - H278 Y284 - H306	zinc finger protein ZFPI13 (GI 5640017)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
11	428	S72 S92 S101 S118 S120 S125 T245 T277 S289 S315 S317 Y326 S409			Skeletal muscle BOP2 (GI 5870834; SEQ ID NO:117)	BLAST, MOTIFS

TABLE 2 (cont.)

Polypeptide Seq ID No:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
12	590	S45 Y52 T60 S83 S90 T95 T116 T145 T233 T330 S391 S410 S411 T420 T439 T490 S521 S15 Y29 S30 S118 T173 T183 S203 S217 Y232 S235 T255 S352 S362 Y451			Methyl-CpG binding protein (GI 2239126)	BLAST, MOTIFS
13	479			Y232 - H254 H283 - H305 Y311 - H333 Y339 - H361 PFAM	SRE-ZBP (GI g36603)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
14	433	S92 S96 T250 S319 T322 T327 S335 T344		C380 - C421 PFAM	C3HC4-type zinc finger protein	PFAM, BLOCKS, MOTIFS
15	320	T6 S27 T125 T172 S229 S232 T239 S248 S259 S266 Y267 S291			Zinc finger factor (GI 3150148)	BLAST, MOTIFS
16	179	S11 T21 S46 S140			Single-stranded DNA binding protein (csdp) (GI 1562534)	BLAST, MOTIFS
17	494	T73 Y80 S104 Y116 T192 S289 S297 T329 T364 T376 S387		C13 - H41 BLOCKS	Zinc finger transcription factor (GI 2895870)	BLAST, BLOCKS, MOTIFS
18	401	S4 S82 S97 T166 S188 S249 S279 S289 S290 S294 S319 S368 S371 S372 S378 T392 S396		P5 - K81 A114 - S179 G186 - P262 PFAM	HPI-BP74 (GI 1480112)	PFAM, BLOCKS, PRINTS, MOTIFS



TABLE 2 (cont.)

Polyypeptide Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
19	264	S11 S25 S76 S82 S90 S92 S96 S119 T229		F154 - H176 C180 - H202 F208 - H230 Y236 - C259 PFAM	C2H2-type zinc finger protein (GI 429188)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
20	153	T23 S40 T44 S110 S120 T124		R42 - E141 PFAM	High mobility group-like nuclear protein (GI 2822179)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
21	243	S20 S21 S76 S100 S104 S160 T194 S196 S212 T222 Y229		Y90 - H112 H118 - H140 Y146 - H168 Y174 - H195 PFAM	C2H2-type zinc finger protein (GI 38015)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
22	485	T29 S34 S104 S147 T162 T248 S249 S256 S347 S452 S477		S309 - H331 H337 - H359 Y365 - H387 Y393 - H415 PFAM	BTB domain/C2H2-type zinc finger protein (GI 2843171)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
23	160	S118		C5 - F62 C80 - F137 PFAM	LIM domain protein/CRP2 (GI 487284)	BLAST, PFAM, BLOCKS, PROFILES SCAN, MOTIFS
24	511	S10 T36 S75 S90 S222 T245 T259 S399 S405 Y443 S500		Y171 - P223 Y267 - K294 BLOCKS	2'-5' oligoadenylate synthetase-related protein p56 (GI 4731857)	BLOCKS, MOTIFS
25	310	S24 S39 T69 Y104 S185 T282 T296			SIR2 family transcriptional regulatory protein (GI 2648874)	BLAST, MOTIFS

TABLE 2 (cont.)

Polypeptide Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
26	331	T29 S40 S74 S257 T270 S301		V112 - R134 BLOCKS	Histone protein	BLOCKS, MOTIFS
27	200	T43 S123 T129 S167 S183 S184			Zinc finger protein (g1373394)	MOTIFS BLAST
28	100	S44 Y25 Y98		Transcription anti-terminator; bglG family: E47-I100	transcription elongation factor (g4336506)	MOTIFS BLAST BLOCKS
29	528	S204 T487 S29 S34 T48 T227 S327 T367 T423 S483 Y39 Y44 Y112 Y163	N24 N52 N100 N481	C2H2 zinc fingers: Y191-R213 Y247-H269 Y275-H297 Y331-H353 Y387-H409 Y415-H437 Y443-H465 Y471-H493	Zinc finger protein (g498721)	MOTIFS BLAST BLOCKS PFAM
30	350	T264 S305	N33 N79	C3HC4 RING finger: C230-C271	C3HC4/RING zinc finger protein (g1321818)	MOTIFS BLAST BLOCKS PFAM PROFILES CAN
31	315	S51 T94 S121 S123 S142 S143 T184 S232 S252 T36 T46 S159 S163 S168			Similar to CCAAT/enhancer-binding protein (g1947129)	MOTIFS BLAST
32	120	S36 S56 T93 S104	N98 N103	bZIP transcription factor: P21-P85		MOTIFS PFAM BLOCKS

TABLE 2 (cont.)

Polypeptide Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
33	326	S59 S38 T207 S284 T319 T43 S80 T137 T155 T211 S238 T239	N209	C2H2 zinc fingers: C143-C171 Y169-H191 F197-H219	Zinc finger protein (g220443)	MOTIFS BLAST PFAM BLOCKS
34	106	S80 T89 T39 T53		Homeobox: R14-K70	CHOX M product; homeobox protein (g62701)	MOTIFS BLAST BLOCKS PFAM PROFLESCAN
35	209	S176 T180 S184 T193 S201 S4 T25 S49	N18	bZIP transcription factor: K115-E140	geminin (g3219357)	MOTIFS BLAST PFAM
36	212	S79 S107 T127 T202 S45 S56 S124 T152 T35	N92	HMG box: M1-Q36	Smad3-related protein (GI 4321968)	MOTIFS BLAST PFAM
37	359	T329 T50 S125 S224 S230 S235 S344 S31 S215 S312 V42	N45 N340	C2H2 zinc fingers: F278-C306 Y304-H328 F334-H356	BKLF; CACCC-box binding protein (g1244515)	MOTIFS BLAST BLOCKS PFAM
38	445	S68 T87 S153 S339 S405 S55 T105 S315 S422 V419	N337 N374 N388	C3HC4 RING fingers: C74-P120 N228-C235	ARI (RING finger) protein (g2058299)	MOTIFS BLAST BLOCKS PFAM
39	433	S283 T44 T57 T123 S136 T185 T220 S239 T268 S313 S330 T105 T109 S125 T216	N169 N206		skm-BOP2 zinc finger protein (g1809327)	MOTIFS BLAST

TABLE 2 (cont.)

Polypeptide Seq ID NO.	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
40	355	T72 T84 T184 S191 T244 S88 T162 T229 T294 S330	N308 N324		Sir2 family protein (g5353746)	MOTIFS BLAST
41	443	T76 T106 S148 T309 S12 T253 S299 S357 T373 T427 Y61 Y114	N366 N425	Myb-like DNA binding domain: D72-F118	ADA2 transcriptional adaptor protein (g170991)	MOTIFS BLAST BLOCKS PFAM PROFLESCAN
42	164	S37 T55 T64 S83 S22 Y146	N117	mut domain: V29-L70		MOTIFS BLOCKS PFAM
43	215	S79 S127 T12 T45		HMG box: M1-Q36	Sry-related protein (g211510)	MOTIFS BLAST PFAM
44	539	T15 S24 S66 S83 T97 T105 T109 S128 T149 S153 S198 T203 T225 T238 T296 S466 Y258	N465	C2H2 zinc fingers: H230-H252 Y258-H280 Y286-H308 Y314-H336 Y342-H364 F370-H392 Y398-H420 Y426-H448 Y482-H504 Y510-H532	C2H2 zinc finger protein (g5757625)	MOTIFS BLAST PFAM BLOCKS
45	182	T59 S112 S120 S100 S139 Y64			Transcriptional regulator (g2621798)	MOTIFS BLAST

TABLE 2 (cont.)

Polypeptide Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
46	534	S494 S31 S44 S117 T123 S185 S216 S476 S504 S176 S182 S211 T249 S293 S323 S409 T489 I76 Y285	N29 N39 N250 N351	C2H2 zinc fingers: Y285-H307 Y313-H335 Y341-H363 C369-H391 Y397-H419 Y425-H447 Y453-H475 Y481-H503 Y509-H531	Zinc finger protein (g1373394)	MOTIFS BLAST PFAM BLOCKS
		S5 S7 S40 S45 S46 S100 S144 S26 S107 T148 S185 Y38	N44 N177	Myc-type HLH domain: Q108-R160	Musculin (g3599519)	MOTIFS BLAST PFAM BLOCKS PROFILES SCAN
		T5 S87 S96 S115 T124 S22 T64 S185 S14 S48 T54 S118 T139 T161 T189 T217 Y256	N210 N214 N238 N260	C2H2 zinc fingers: F172-H194 Y200-H222 Y228-H250	KRAB zinc finger protein (g1049295)	MOTIFS BLAST
		S157 S42 T167 T222 T81 Y213	N40	C3HC4 RING finger: P126-L150	Repressor transcriptional factor (g1017722)	MOTIFS BLAST PFAM BLOCKS
50	236	S7 S8 S116 T127 S154 S191 T31 S41 T204	N2	Chromodomain: V113-E134	Ariadne-2 RING finger protein (g3445441)	MOTIFS BLAST BLOCKS
51	214				Nucleoplasmin (g833629)	MOTIFS BLAST BLOCKS

TABLE 2 (cont.)

Polypeptide Seq ID No:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
52	396	T148 T392 S118 T193 T201 S270 S294 S80 S112 S206 S260 T313 T355 S375 S387 S29 T58 S155 S239 T292 T379 S146 T271 S425	N2	C3HC4 RING finger: C26-C50	Midline 1/ cerebellar isoform 1 RING finger protein (g3462503)	MOTIFS BLAST PFAM BLOCKS PROFILERSCAN
53	486	S432 T502 S68 S195 T199 T226 S115 T379 T441 S282 S291 T327 T136 S391 S422 T481 Y257 Y274 T14 T42 Y48	N25 N66 N246 N364	ATP/GTP binding site (P-loop): A434-T441	5'-nucleotidase (g633071)	MOTIFS BLAST
54	555				Transcription termination factor I (TTF-I) interacting peptide 5 isoform (g2183083)	MOTIFS BLAST
55	61				Putative leucine-rich DNA-binding protein (g555991)	MOTIFS BLAST

TABLE 3

Nucleotide Seq ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
56	169-215	Reproductive (0.224) Nervous (0.198) Cardiovascular (0.112)	Cell Proliferative (0.725) Inflammation (0.190)	PBLUESCRIPT
57	551-595	Hematopoietic/Immune (0.240) Reproductive (0.180) Gastrointestinal (0.120)	Cell Proliferative (0.700) Inflammation (0.360)	PBLUESCRIPT
58	541-585	Nervous (0.286) Reproductive (0.286) Cardiovascular (0.214)	Cell Proliferative (0.643) Trauma (0.214)	PBLUESCRIPT
59	109-153	Reproductive (1.000)	Cell Proliferative (1.000) Inflammation (1.000)	PBLUESCRIPT
60	435-479	Hematopoietic/Immune (0.211) Gastrointestinal (0.183) Reproductive (0.183)	Cell Proliferative (0.620) Inflammation (0.338)	PBLUESCRIPT
61	1195-1239	Reproductive (0.248) Cardiovascular (0.174) Nervous (0.157)	Cell Proliferative (0.637) Inflammation (0.256)	PBLUESCRIPT
62	217-261	Reproductive (0.429) Nervous (0.238) Cardiovascular (0.095)	Cell Proliferative (0.667) Inflammation (0.143) Trauma (0.095)	PBLUESCRIPT
63	919-963	Reproductive (0.265) Nervous (0.235) Cardiovascular (0.088)	Cell Proliferative (0.618) Inflammation (0.206)	PSPORT1
64	823-876	Reproductive (0.382) Nervous (0.176) Gastrointestinal (0.118)	Cell Proliferative (0.794)	PSPORT1
65	380-424	Reproductive (0.346) Nervous (0.154) Gastrointestinal (0.135)	Cell Proliferative (0.750) Inflammation (0.231)	PSPORT1

TABLE 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
66	757-801	Nervous (0.222) Hematopoietic/Immune (0.167) Reproductive (0.167)	Cell Proliferative (0.778) Inflammation (0.222)	pINCY
67	812-856	Reproductive (0.246) Nervous (0.180) Gastrointestinal (0.148)	Cell Proliferative (0.639) Inflammation (0.246)	pINCY
68	326-370	Reproductive (0.500) Nervous (0.200) Gastrointestinal (0.150)	Cell Proliferative (0.700) Trauma (0.150)	pINCY
69	703-747	Reproductive (0.278) Hematopoietic/Immune (0.204) Nervous (0.148)	Cell Proliferative (0.777) Inflammation (0.222)	pINCY
70	759-803	Reproductive (0.261) Gastrointestinal (0.217) Nervous (0.174)	Cell Proliferative (0.565) Trauma (0.130)	pINCY
71	110-154	Nervous (0.250) Developmental (0.208) Gastrointestinal (0.167)	Cell Proliferative (0.583) Trauma (0.167)	pINCY
72	529-573	Reproductive (0.186) Gastrointestinal (0.168) Hematopoietic/Immune (0.138)	Cell Proliferative (0.700) Inflammation (0.251)	pINCY
73	1784-1828	Reproductive (0.286) Hematopoietic/Immune (0.190) Nervous (0.167)	Cell Proliferative (0.667) Inflammation (0.286)	PSPORT1
74	111-155	Reproductive (0.316) Nervous (0.211) Hematopoietic/Immune (0.158)	Cell Proliferative (0.632) Inflammation (0.211)	PSPORT1
75	543-587	Reproductive (0.258) Nervous (0.206) Gastrointestinal (0.134)	Cell Proliferative (0.608) Inflammation (0.196)	pINCY



TABLE 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
76	272-316	Reproductive (0.246) Nervous (0.180) Hematopoietic/Immune (0.148)	Cell Proliferative (0.606) Inflammation (0.279)	PINCY
77	227-271	Hematopoietic/Immune (0.222) Endocrine (0.167) Cardiovascular (0.111)	Cell Proliferative (0.666) Inflammation (0.333)	PINCY
78	487-531	Gastrointestinal (0.375) Reproductive (0.250)	Cell Proliferative (0.500) Inflammation (0.250)	PINCY
79	111-155	Gastrointestinal (0.280) Hematopoietic/Immune (0.240) Reproductive (0.120)	Cell Proliferative (0.640) Inflammation (0.440)	PINCY
80	595-639	Reproductive (0.211) Gastrointestinal (0.158) Urologic (0.158)	Cell Proliferative (0.684) Inflammation (0.263)	PINCY
81	425-469	Reproductive (0.222) Gastrointestinal (0.160) Nervous (0.148)	Cell Proliferative (0.568) Inflammation (0.259)	PSPORT1
82	774-818	Gastrointestinal (0.200) Hematopoietic/Immune (0.200) Nervous (0.200)	Cancer (0.600) Trauma (0.200) Inflammation (0.200)	PBLUESCRIPT
83	517-561	Nervous (0.526) Reproductive (0.132) Cardiovascular (0.105)	Cancer (0.342) Fetal (0.158) Inflammation (0.158)	PBLUESCRIPT
84	1944-1988	Nervous (0.250) Reproductive (0.188) Endocrine (0.125)	Cancer (0.438) Fetal (0.250) Trauma (0.250)	PSPORT1
85	1027-1071	Reproductive (0.219) Nervous (0.206) Hematopoietic/Immune (0.116)	Cancer (0.458) Inflammation (0.232) Fetal (0.181)	PSPORT1

TABLE 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
86	658-702	Reproductive (0.227) Hematopoietic/Immune (0.182) Gastrointestinal (0.167)	Cancer (0.424) Inflammation (0.318) Fetal (0.167)	PT7T3
87	488-532	Nervous (0.200) Reproductive (0.200) Musculoskeletal (0.120)	Cancer (0.320) Fetal (0.320) Inflammation (0.320)	pINCY
88	379-423	Cardiovascular (0.250) Nervous (0.250) Reproductive (0.250)	Cancer (0.417) Fetal (0.167) Neurological (0.167)	pINCY
89	632-676	Reproductive (0.417) Cardiovascular (0.167) Gastrointestinal (0.167)	Cancer (0.333) Fetal (0.167) Inflammation (0.167)	pINCY
90	258-302	Reproductive (0.294) Nervous (0.137) Hematopoietic/Immune (0.118)	Cancer (0.569) Fetal (0.431) Inflammation (0.176)	pINCY
91	433-477	Reproductive (0.750) Nervous (0.250)	Cancer (0.500) Inflammation (0.500)	PSPORT1
92	542-586	Gastrointestinal (0.273) Hematopoietic/Immune (0.273) Developmental (0.182)	Cancer (0.455) Inflammation (0.364) Fetal (0.182)	pINCY
93	218-262	Reproductive (0.272) Nervous (0.204) Cardiovascular (0.126)	Cancer (0.447) Inflammation (0.214) Fetal (0.155)	PSPORT1
94	541-585	Reproductive (0.273) Nervous (0.250) Cardiovascular (0.159)	Cancer (0.364) Fetal (0.205) Inflammation (0.205)	PSPORT1
95	111-155	Reproductive (0.250) Gastrointestinal (0.173) Nervous (0.154)	Cancer (0.481) Fetal (0.231) Inflammation (0.212)	PSPORT1
96	597-641	Reproductive (0.261) Cardiovascular (0.217) Nervous (0.130)	Cancer (0.391) Fetal (0.304) Inflammation (0.130)	PSPORT1

TABLE 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
97	434-478	Cardiovascular (0.222) Endocrine (0.222) Gastrointestinal (0.222)	Cancer (0.667) Fetal (0.111) Neurological (0.111)	PSPORT1
98	218-247 920-964	Reproductive (0.333) Gastrointestinal (0.129) Hematopoietic/Immune (0.118)	Cancer (0.559) Inflammation (0.204) Fetal (0.183)	pINCY
99	327-371	Gastrointestinal (0.211) Reproductive (0.211) Cardiovascular (0.158)	Cancer (0.421) Fetal (0.316) Inflammation (0.158)	pINCY
100	596-625	Reproductive (0.230) Nervous (0.164) Gastrointestinal (0.131)	Cancer (0.590) Inflammation (0.246) Fetal (0.082)	PSPORT
101	487-531	Cardiovascular (0.235) Reproductive (0.235) Hematopoietic/Immune (0.176)	Cancer (0.588) Inflammation (0.176) Trauma (0.118)	pINCY
102	218-247 542-586	Gastrointestinal (0.241) Hematopoietic/Immune (0.207) Cardiovascular (0.138)	Cancer (0.448) Fetal (0.276) Inflammation (0.276)	pINCY
103	219-263	Reproductive (0.500) Cardiovascular (0.250) Hematopoietic/Immune (0.250)	Cancer (0.500) Inflammation (0.250) Trauma (0.250)	pINCY
104	111-140 327-371	Hematopoietic/Immune (0.286) Nervous (0.238)	Cancer (0.333) Fetal (0.286) Inflammation (0.286)	pINCY
105	243-281	Musculoskeletal (0.286) Nervous (0.286)	Inflammation (0.429) Fetal (0.286) Cancer (0.286)	pINCY
106	271-315	Gastrointestinal (0.143) Nervous (0.800) Reproductive (0.100) Cardiovascular (0.100)	Cancer (0.400) Inflammation (0.200) Trauma (0.200)	pINCY

TABLE 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
107	489-533	Cardiovascular (0.364) Gastrointestinal (0.182) Reproductive (0.182)	Cancer (0.273) Trauma (0.273) Inflammation (0.182)	PINCY
108	156-200	Nervous (0.256) Reproductive (0.256) Hematopoietic/Immune (0.128)	Cancer (0.465) Fetal (0.291) Inflammation (0.186)	PINCY
109	1459-1503	Cardiovascular (0.250) Hematopoietic/Immune (0.250) Nervous (0.167)	Inflammation (0.417) Cancer (0.333) Trauma (0.167)	PSPORT1
110	164-208	Nervous (1.000)	Neurological (1.000)	PINCY

TABLE 4

Nucleotide SEQ ID NO:	Library	Library Comment
56	SPLNFETO 1	Library was constructed at Stratagene, using RNA isolated from a pool of fetal spleen tissue. Following vector packaging, 2x10 <sup>6</sup> primary clones were then amplified to stabilize the library for long-term storage. Amplification may significantly skew sequence abundances.
57	SYNORABO 1	Library was constructed using RNA isolated from the synovial membrane tissue of a 68-year-old Caucasian female with rheumatoid arthritis.
58	KIDNNOTO 1	Library was constructed using RNA isolated from the kidney tissue of a 64-year-old Caucasian female, who died from an intracranial bleed. Patient history included rheumatoid arthritis and tobacco use.
59	PLACNOBO 1	Library was constructed using RNA isolated from placenta.
60	HNTZRATO 1	Library was constructed at Stratagene (STR37231), using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated with retinoic acid for 24 hours.
61	EOSIHETO 2	Library was constructed using RNA isolated from peripheral blood cells apheresed from a 48-year-old Caucasian male. Patient history included hyperesinophilia. The cell population was determined to be greater than 77% eosinophils by Wright's staining.
62	HNTZAGTO 1	Library was constructed at Stratagene (STR37233), using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated with retinoic acid for 5 weeks and with mitotic inhibitors for two weeks and allowed to mature for an additional 4 weeks in conditioned medium.
63	PROSNOTO 2	Library was constructed using RNA isolated from the diseased prostate tissue removed from a 50-year-old Caucasian male during a retropubic prostatectomy. Pathology indicated adenofibromatous hyperplasia was present. Pathology for the associated tumor tissue indicated adenocarcinoma Gleason grade 3+3. Patient history included dysuria, carcinoma in situ of prostate, coronary atherosclerosis, and hyperlipidemia.
64	COLNUTTO 2	Library was constructed using RNA isolated from colon tumor tissue removed from a 75-year-old Caucasian male during a hemicolectomy. Pathology indicated invasive grade 3 adenocarcinoma arising in a tubulovillous adenoma, which was distal to the ileocecal valve in the cecum. The tumor penetrated deeply into the muscularis propria but not through it.

TABLE 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
65	KIDNTUT0 1	Library was constructed using RNA isolated from the kidney tumor tissue removed from an 8-month-old female during nephroureterectomy. Pathology indicated Wilms' tumor (nephroblastoma), which involved 90 percent of the renal parenchyma. Prior to surgery, the patient was receiving heparin anticoagulant therapy.
66	THYRN070 3	Library was constructed using RNA isolated from thyroid tissue removed from the left thyroid of a 28-year-old Caucasian female during a complete thyroidectomy. Pathology indicated a small nodule of adenomatous hyperplasia present in the left thyroid. Pathology for the associated tumor tissue indicated dominant follicular adenoma, forming a well-encapsulated mass in the left thyroid.
67	BLADTUT0 4	Library was constructed using RNA isolated from bladder tumor tissue removed from a 60-year-old Caucasian male during a radical cystectomy, prostatectomy, and vasectomy. Pathology indicated grade 3 transitional cell carcinoma in the left bladder wall. Carcinoma in-situ was identified in the dome and trigone. Patient history included tobacco use. Family history included type I diabetes, a malignant neoplasm of the stomach, atherosclerotic coronary artery disease, and an acute myocardial infarction.
68	PROSNOT1 5	Library was constructed using RNA isolated from diseased prostate tissue removed from a 66-year-old Caucasian male during radical prostatectomy and regional lymph node excision. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated an adenocarcinoma (Gleason grade 2+3). The patient presented with elevated prostate specific antigen (PSA). Family history included prostate cancer, secondary bone cancer, and benign hypertension.
69	BRSTTUT0 8	Library was constructed using RNA isolated from breast tumor tissue removed from a 45-year-old Caucasian female during unilateral extended simple mastectomy. Pathology indicated invasive nuclear grade 2-3 adenocarcinoma, ductal type, with 3 of 23 lymph nodes positive for metastatic disease. Greater than 50% of the tumor volume was in situ, both comedo and non-comedo types. Immunostains were positive for estrogen/progesterone receptors, and uninvolved tissue showed proliferative changes. The patient concurrently underwent a total abdominal hysterectomy. Patient history included valvuloplasty of mitral valve without replacement, rheumatic mitral insufficiency, and rheumatic heart disease. Family history included acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes.

TABLE 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
70	PROSN071 9	Library was constructed using RNA isolated from diseased prostate tissue removed from a 59-year-old Caucasian male during a radical prostatectomy with regional lymph node excision. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated an adenocarcinoma (Gleason grade 3+3). The patient presented with elevated prostate-specific antigen (PSA). Patient history included colon diverticuli, asbestosis, and thrombophlebitis. Previous surgeries included a partial colectomy. Family history included benign hypertension, multiple myeloma, hyperlipidemia and rheumatoid arthritis.
71	THP1N070 3	Library was constructed using RNA isolated from untreated THP-1 cells. THP-1 (ATCC TIB 202) is a human promonocyte line derived from the peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia (ref: Int. J. Cancer (1980) 26:171).
72	THP1N070 3	Library was constructed using RNA isolated from untreated THP-1 cells. THP-1 (ATCC TIB 202) is a human promonocyte line derived from the peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia (ref: Int. J. Cancer (1980) 26:171).
73	H1POAZT0 1	Library was constructed from RNA isolated from diseased hippocampus tissue removed from the brain of a 74-year-old Caucasian male who died from Alzheimer's disease.
74	H1POAZT0 1	Library was constructed from RNA isolated from diseased hippocampus tissue removed from the brain of a 74-year-old Caucasian male who died from Alzheimer's disease.
75	BRSTN071 3	Library was constructed using RNA isolated from breast tissue removed from the left medial lateral breast of a 36-year-old Caucasian female during bilateral simple mastectomy and total breast reconstruction. Pathology indicated benign breast tissue. Patient history included a breast neoplasm, depressive disorder, hyperlipidemia, chronic stomach ulcer, and an ectopic pregnancy. Family history included myocardial infarction, cerebrovascular disease, atherosclerotic coronary artery disease, hyperlipidemia, skin cancer, breast cancer, depressive disorder, esophageal cancer, bone cancer, Hodgkin's lymphoma, bladder cancer, and heart condition.
76	NPOLN070 1	Library was constructed using RNA isolated from nasal polyp tissue removed from a 78-year-old Caucasian male during a nasal polypectomy. Pathology indicated a nasal polyp and striking eosinophilia. Patient history included asthma and nasal polyps.

TABLE 4 (cont.)

Nucleotide Seq ID NO:	Library	Library Comment
77	ADRENOTO 9	Library was constructed using RNA isolated from left adrenal gland tissue removed from a 43-year-old Caucasian male during nephroureterectomy, regional lymph node excision, and unilateral left adrenalectomy. Pathology indicated no diagnostic abnormalities of the adrenal gland. Pathology for the associated tumor tissue indicated a grade 2 renal cell carcinoma mass in the posterior lower pole of the left kidney with invasion into the renal pelvis.
78	SPLNOT1 1	Library was constructed using RNA isolated from diseased spleen tissue removed from a 14-year-old Asian male during a total splenectomy. Pathology indicated changes consistent with idiopathic thrombocytopenic purpura. The patient presented with bruising.
79	CONCNOTO 1	Library was constructed using RNA isolated from chest wall soft tissue removed from a 63-year-old Caucasian male during a chest wall lesion destruction. Pathology indicated surgical margins were free of tumor. Pathology for the associated tumor tissue indicated invasive grade 3 adenocarcinoma, forming a mass that extended through the visceral pleura to involve parietal pleura. Patient history included MEN (multiple endocrine neoplasia) syndrome type I, abnormal secretion of gastrin, alcohol and tobacco abuse, calcium metabolism disease, chronic stomach ulcer with hemorrhage, lung cancer, and calculus of the kidney. Family history included prostate cancer, benign hypertension, stroke, atherosclerotic coronary artery disease, type II diabetes, hyperlipidemia, and cancer of an unspecified location.
80	TONSNOTO 3	Library was constructed using RNA isolated from diseased left tonsil tissue removed from a 6-year-old Caucasian male during adenotonsillectomy. Pathology indicated reactive lymphoid hyperplasia, bilaterally. Family history included benign hypertension, myocardial infarction, and atherosclerotic coronary artery disease.
81	LUNGNOTO 3	This normalized library was constructed from $2.56 \times 10^5$ independent clones from a lung tissue library. RNA was made from lung tissue removed from the left lobe of a 58-year-old Caucasian male during a segmental lung resection. Pathology for the associated tumor tissue indicated a metastatic grade 3 (of 4) osteosarcoma. Patient history included soft tissue cancer, secondary cancer of the lung, prostate cancer, and an acute duodenal ulcer with hemorrhage. Patient also received radiation therapy to the retroperitoneum. Family history included prostate cancer, breast cancer, and acute leukemia. The normalization and hybridization conditions were adapted from Soares et al., PNAS (1994) 91:9228; Swaroop et al., NAR (1991) 19:1954; and Ronaldo et al., Genome Research (1996) 6:791.



TABLE 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
82	BMANOTO 2	The library was constructed using Clontech RNA isolated from the bone marrow of 24 male and female Caucasian donors, 16 to 70 years old.
83	SINTNOTO 2	The library was constructed using RNA isolated from the small intestine of a 55-year-old Caucasian female, who died from a subarachnoid hemorrhage. Serologies were positive for cytomegalovirus (CMV).
84	CRELNOTO 1	The library was constructed using RNA isolated from the cerebellum tissue of a 69-year-old Caucasian male who died from chronic obstructive pulmonary disease. Patient history included myocardial infarction, hypertension, and osteoarthritis.
85	BRSTNOTO 3	The library was constructed using RNA isolated from diseased breast tissue removed from a 54-year-old Caucasian female during a bilateral radical mastectomy. Pathology for the associated tumor tissue indicated residual invasive grade 3 mammary ductal adenocarcinoma. Patient history included kidney infection and condyloma acuminatum. Family history included benign hypertension, hyperlipidemia and a malignant neoplasm of the colon.
86	BEPINONO 1	The normalized bronchial epithelium library was constructed from 5.12 million independent clones from a bronchial epithelium library. RNA was isolated from a bronchial epithelium primary cell line derived from a 54-year-old Caucasian male. The normalization and hybridization conditions were adapted from Soares et al., PNAS (1994) 91:9228, using a 24-hour reannealing hybridization period.
87	THYRNOTO 3	The library was constructed using RNA isolated from thyroid tissue removed from the left thyroid of a 28-year-old Caucasian female during a complete thyroidectomy. Pathology indicated a small nodule of adenomatous hyperplasia present in the left thyroid. Pathology for the associated tumor tissue indicated dominant follicular adenoma, forming a well-encapsulated mass in the left thyroid.
88	COLNFETO 2	The library was constructed using RNA isolated from the colon tissue of a Caucasian female fetus who died at 20 weeks' gestation.
89	BLADNOTO 3	The library was constructed using RNA isolated from bladder tissue removed from an 80-year-old Caucasian female during a radical cystectomy and lymph node excision. Pathology for the associated tumor tissue indicated grade 3 invasive transitional cell carcinoma. Patient history included malignant neoplasm of the uterus, atherosclerosis, and atrial fibrillation. Family history included acute renal failure, osteoarthritis, and atherosclerosis.

TABLE 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
90	COLNNOT2 3	The library was constructed using RNA isolated from diseased colon tissue removed from a 16-year-old Caucasian male during a total colectomy with abdominal/perineal resection. Pathology indicated gastritis and pancolitis consistent with the acute phase of ulcerative colitis. Inflammation was more severe in the transverse colon with inflammation confined to the mucosa. The ascending and sigmoid colon was mildly involved. Family history included irritable bowel syndrome.
91	BRSTNOTO 4	The library was constructed using RNA isolated from breast tissue removed from a 62-year-old East Indian female during a unilateral extended simple mastectomy. Pathology for the associated tumor tissue indicated an invasive grade 3 ductal carcinoma. Patient history included benign hypertension, hyperlipidemia, and hematuria. Family history included cerebrovascular and cardiovascular disease, hyperlipidemia, and liver cancer.
92	LIVRFETO 2	The library was constructed using RNA isolated from liver tissue removed from a Caucasian female fetus who died at 20 weeks' gestation. Family history included seven days of erythromycin treatment for bronchitis in the mother during the first trimester.
93	BEPINOTO 1	The library was constructed using RNA isolated from a bronchial epithelium primary cell line derived from a 54-year-old Caucasian male.
94	BRAITUTO 2	The library was constructed using RNA isolated from brain tumor tissue removed from the frontal lobe of a 58-year-old Caucasian male during excision of a cerebral meningial lesion. Pathology indicated a grade 2 metastatic hypernephroma. Patient history included a grade 2 renal cell carcinoma, insomnia, and chronic airway obstruction. Family history included a malignant neoplasm of the kidney.
95	BRAITUTO 3	The library was constructed using RNA isolated from brain tumor tissue removed from the left frontal lobe of a 17-year-old Caucasian female during excision of a cerebral meningial lesion. Pathology indicated a grade 4 fibrillary giant and small-cell astrocytoma. Family history included benign hypertension and cerebrovascular disease.
96	BRSTTUTO 2	The library was constructed using RNA isolated from breast tumor tissue removed from a 54-year-old Caucasian female during a bilateral radical mastectomy with reconstruction. Pathology indicated residual invasive grade 3 mammary ductal adenocarcinoma. The remaining breast parenchyma exhibited proliferative fibrocystic changes without atypia. One of 10 axillary lymph nodes had metastatic tumor. Patient history included kidney infection and condyloma acuminatum. Family history included benign hypertension, hyperlipidemia, and a malignant colon neoplasm.

TABLE 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
97	BRSTTUT0 2	The library was constructed using RNA isolated from breast tumor tissue removed from a 54-year-old Caucasian female during a bilateral radical mastectomy with reconstruction. Pathology indicated residual invasive grade 3 mammary ductal adenocarcinoma. The remaining breast parenchyma exhibited proliferative fibrocystic changes without atypia. One of 10 axillary lymph nodes had metastatic tumor. Patient history included kidney infection and condyloma acuminatum. Family history included benign hypertension, hyperlipidemia, and a malignant colon neoplasm.
98	BRSTN070 7	The library was constructed using RNA isolated from diseased breast tissue removed from a 43-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated mildly proliferative fibrocystic changes with epithelial hyperplasia, papillomatosis, and duct ectasia. Pathology for the associated tumor tissue indicated invasive grade 4, nuclear grade 3 mammary adenocarcinoma with extensive comedo necrosis. Family history included epilepsy, cardiovascular disease, and type II diabetes.
99	BRAIN070 9	The library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus who died at 23 weeks' gestation.
100	THP1AZ50 8	The library was constructed using RNA isolated from 5.76 million clones from a 5-aza-2'-deoxycytidine treated THP-1 cell library. The library was subjected to subtractive hybridization using 5 million clones from an untreated THP-1 cell library. Hybridization conditions were adapted from Swaroop et al., NAR (1991) 19:1954; and Bonaldo et al., Genome Research (1996) 6:791. THP-1 (ATCC TIB 202) is a human promonocyte cell line derived from peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia.
101	LUNG07U1 2	The library was constructed using RNA isolated from tumorous lung tissue removed from a 70-year-old Caucasian female during a lung lobectomy of the left upper lobe. Pathology indicated grade 3 (of 4) adenocarcinoma and vascular invasion. Patient history included tobacco abuse, depressive disorder, anxiety state, and skin cancer. Family history included cerebrovascular disease, congestive heart failure, colon cancer, depressive disorder, and primary liver.
102	HEA0N070 5	The library was constructed using RNA isolated from aortic tissue removed from a 17-year-old Hispanic female who died from a gunshot wound.
103	HEA0N070 5	The library was constructed using RNA isolated from aortic tissue removed from a 17-year-old Hispanic female who died from a gunshot wound.

TABLE 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
104	EPIGNOTO 1	The library was constructed using RNA isolated from epiglottic tissue removed from a 71-year-old male during laryngectomy with right parathyroid biopsy. Pathology for the associated tumor tissue indicated recurrent grade 1 papillary thyroid carcinoma.
105	CONCNOTO 1	The library was constructed using RNA isolated from chest wall soft tissue removed from a 63-year-old Caucasian male during a chest wall lesion destruction. Pathology for the associated tumor tissue indicated invasive grade 3 adenocarcinoma forming a mass that extended through the visceral pleura to involve parietal pleura. Patient history included multiple endocrine neoplasia syndrome type 1, abnormal secretion of gastrin, alcohol and tobacco abuse, calcium metabolism disease, chronic stomach ulcer with hemorrhage, lung cancer, and calculus of the kidney. Family history included prostate cancer, benign hypertension, stroke, atherosclerotic coronary artery disease, type II diabetes, hyperlipidemia, and an unspecified cancer.
106	BRONNOTO 1	The library was constructed using RNA isolated from bronchial tissue removed from a 15-year-old Caucasian male.
107	BRONNOTO 1	The library was constructed using RNA isolated from bronchial tissue removed from a 15-year-old Caucasian male.
108	LIVRNOTO 3	The library was constructed using RNA isolated from liver tissue removed from a Caucasian male fetus who died from Patau's syndrome (trisomy 13) at 20 weeks' gestation.
109	LUNGNOTO 3	The normalized library was constructed from 2.56 million independent clones from a lung tissue library. RNA was isolated from lung tissue removed from the left lobe a 58-year-old Caucasian male during a segmental lung resection. Pathology for the associated tumor tissue indicated a metastatic grade 3 (of 4) osteosarcoma. Patient history included soft tissue cancer, secondary cancer of the lung, prostate cancer, acute duodenal ulcer with hemorrhage, and radiation therapy to the retroperitoneum. Family history included prostate cancer, breast cancer, and acute leukemia. The normalization and hybridization conditions were adapted from Soares et al., PNAS (1994) 91:9228; Swaroop et al., NAR (1991) 19:1954; and Donaldo et al., Genome Research (1996) 6:791.
110	BRAIHCOTO 2	The library was constructed using RNA isolated from diseased choroid plexus tissue removed from the brain of a 57-year-old Caucasian male who died from a cerebrovascular accident. Patient history included Huntington's disease and emphysema.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, ifasta, tfasta, tfasta, and search.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.0E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater fasta E value= 1.0E-8 or less Full Length sequences: fasta score= 100 or greater
BLIMPS	A BLOCKS IMPROVED Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff, Nucl. Acid Res., 19:565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score=1000 or greater, Ratio of Score/Strength = 0.75 or larger, and, if applicable, Probability value= 1.0E-3 or less
IIMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribkov, M. et al. (1988) CABIOS 4:61-66; Gribkov, et al. (1989) Methods Enzymol. 183:146-159; Baiocchi, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Normalized quality scores GCG-specified "HIGH" value for that particular Prosite motif Generally, score=14-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program, including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Clavette, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Baiocchi et al. supg; Wisconsin Package Program Manual, version 9, page M51-59. Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising:

- a) an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3-5, SEQ ID NO:7-14, SEQ ID NO:16-31, SEQ ID NO:33-34, SEQ ID NO:36-40, SEQ ID NO:42-48, SEQ ID NO:50-55.
- b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3-5, SEQ ID NO:7-14, SEQ ID NO:16-31, SEQ ID NO:33-34, SEQ ID NO:36-40, SEQ ID NO:42-48, SEQ ID NO:50-55,
- c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3-5, SEQ ID NO:7-14, SEQ ID NO:16-31, SEQ ID NO:33-34, SEQ ID NO:36-40, SEQ ID NO:42-48, SEQ ID NO:50-55, or
- d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3-5, SEQ ID NO:7-14, SEQ ID NO:16-31, SEQ ID NO:33-34, SEQ ID NO:36-40, SEQ ID NO:42-48, SEQ ID NO:50-55.

2. An isolated polypeptide of claim 1, having an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3-5, SEQ ID NO:7-14, SEQ ID NO:16-31, SEQ ID NO:33-34, SEQ ID NO:36-40, SEQ ID NO:42-48, SEQ ID NO:50-55.

3. An isolated polynucleotide encoding a polypeptide of claim 1.

4. An isolated polynucleotide of claim 3, having a sequence selected from the group consisting of SEQ ID NO:56-110.

5. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

6. A cell transformed with a recombinant polynucleotide of claim 5.

7. A transgenic organism comprising a polynucleotide of claim 5.

8. A method for producing a polypeptide of claim 1, the method comprising:

- a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said

WO 00/44900

PCT/US00/02237

cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

b) recovering the polypeptide so expressed.

5

9. An isolated antibody which specifically binds to a polypeptide of claim 1.

10. An isolated polynucleotide comprising:

10

a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:56-110,

b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:56-110,

c) a polynucleotide sequence complementary to a), or

d) a polynucleotide sequence complementary to b).

15

11. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 10.

12. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 10, the method comprising:

20

a) hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and

25

b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

13. A method of claim 12, wherein the probe comprises at least 30 contiguous nucleotides.

30

14. A method of claim 12, wherein the probe comprises at least 60 contiguous nucleotides.

15. A pharmaceutical composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

35

16. A method of treating a disease or condition associated with decreased expression of



functional NuABP, comprising administering to a patient in need of such treatment the pharmaceutical composition of claim 15.

17. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

18. A pharmaceutical composition comprising an agonist compound identified by a method of claim 17 and a pharmaceutically acceptable excipient.

19. A method of treating a disease or condition associated with decreased expression of functional NuABP, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 18.

20. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

21. A pharmaceutical composition comprising an antagonist compound identified by a method of claim 20 and a pharmaceutically acceptable excipient.

22. A method for treating a disease or condition associated with overexpression of functional NuABP, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 21.

23. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 4, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
3 August 2000 (03.08.2000)

PCT

(10) International Publication Number  
WO 00/44900 A3(51) International Patent Classification<sup>7</sup>: C12N 15/12,  
C12Q 1/68, C07K 14/47, 16/18, G01N 33/68, A61K 38/17

(21) International Application Number: PCT/US00/02237

(22) International Filing Date: 28 January 2000 (28.01.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/117,905 29 January 1999 (29.01.1999) US  
60/117,904 29 January 1999 (29.01.1999) US(63) Related by continuation (CON) or continuation-in-part  
(CIP) to earlier applications:US 60/117,904 (CIP)  
Filed on 29 January 1999 (29.01.1999)  
US 60/117,905 (CIP)  
Filed on 29 January 1999 (29.01.1999)(71) Applicant (for all designated States except US): INCYTE  
PHARMACEUTICALS, INC. [US/US]; 3174 Porter  
Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): TANG, Y., Tom  
[CN/US]; 4230 Ranwick Court, San Jose, CA 95118  
(US). LAL, Pretti [IN/US]; 2382 Lass Drive, Santa Clara,  
CA 95054 (US). HILLMAN, Jennifer, L. [US/US]; 230  
Monrow Drive #12, Mountain View, CA 94040 (US).  
YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale,  
CA 94087 (US). AZIMZAI, Valda [US/US]; 2045 Rock  
Springs Drive, Hayward, CA 94545 (US). LU, Aina,  
M., D. [US/US]; 55 Park Belmont Place, San Jose, CA  
95136 (US). BAUGHN, Mariah, R. [US/US]; 14244Santiago Road, San Leandro, CA 94577 (US). TRAN,  
Bao [US/US]; 744 Kiely Boulevard, Santa Clara, CA  
95051 (US). SHIH, Leo, L. [US/US]; Apartment B., 1081  
Tanland Drive, Palo Alto, CA 94303 (US). AU-YOUNG,  
Janice, L. [US/US]; 233 Golden Eagle Lane, Brisbane,  
CA 94005 (US).(74) Agents: HAMLET-COX, Diana et al.; Incyte Pharma-  
ceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304  
(US).(81) Designated States (national): AE, AL, AM, AT, AU, AZ,  
BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE,  
ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,  
KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,  
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,  
SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ,  
VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

## Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

(88) Date of publication of the international search report:  
30 November 2000For two-letter codes and other abbreviations, refer to the "Guide-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette

(54) Title: NUCLEIC-ACID BINDING PROTEINS

(57) Abstract: The invention provides human nucleic-acid binding proteins (NuABP) and polynucleotides which identify and en-  
code NuABP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also  
provides methods for diagnosing, treating, or preventing disorders associated with expression of NuABP.

Docket No.: PF-0662 USN

**DECLARATION AND POWER OF ATTORNEY FOR  
UNITED STATES PATENT APPLICATION**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name, and

I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if more than one name is listed below) of the subject matter which is claimed and for which a United States patent is sought on the invention entitled

**NUCLEIC-ACID BINDING PROTEINS**

the specification of which:

   / is attached hereto.

   / was filed on \_\_\_\_\_ as application Serial No. \_\_\_\_\_ and if this box contains an X    /, was amended on \_\_\_\_\_.

  X   / was filed as Patent Cooperation Treaty international application No. PCT/US00/02237 on January 28, 2000, this box contains an X    /, was amended on under Patent Cooperation Treaty Article 19 on \_\_\_\_\_ 2001, and if this box contains an X    /, was amended on \_\_\_\_\_.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge my duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim the benefit under Title 35, United States Code, §119 or §365(a)-(b) of any foreign application(s) for patent or inventor's certificate indicated below and of any Patent Cooperation Treaty international application(s) designating at least one country other than the United States indicated below and have also identified below any foreign application(s) for patent or inventor's certificate and Patent Cooperation Treaty international application(s) designating at least one country other than the United States for the same subject matter and having a filing date before that of the application for said subject matter the priority of which is claimed:

**Docket No.: PF-0662 USN**

Country	Number	Filing Date	Priority Claimed
_____	_____	_____	// Yes // No
_____	_____	_____	// Yes // No

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

Application Serial No.	Filed	Status (Pending, Abandoned, Patented)
<u>60/117,905</u>	<u>January 29, 1999</u>	<u>Expired</u>
<u>60/117,904</u>	<u>January 29, 1999</u>	<u>Expired</u>

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in said prior application(s) in the manner required by the first paragraph of Title 35, United States Code §112, I acknowledge my duty to disclose material information as defined in Title 37 Code of Federal Regulations, §1.56(a) which occurred between the filing date(s) of the prior application(s) and the national or Patent Cooperation Treaty international filing date of this application:

Application Serial No.	Filed	Status (Pending, Abandoned, Patented)
_____	_____	_____

I hereby appoint the following:

Lucy J. Billings	Reg. No. <u>36,749</u>
Michael C. Cerrone	Reg. No. <u>39,132</u>
Diana Hamlet-Cox	Reg. No. <u>33,302</u>
Richard C. Ekstrom	Reg. No. <u>37,027</u>
Barrie D. Greene	Reg. No. <u>46,740</u>
Lynn E. Murry	Reg. No. <u>42,918</u>
Shirley A. Recipon	Reg. No. <u>47,016</u>
Susan K. Sather	Reg. No. <u>44,316</u>
Michelle M. Stempfen	Reg. No. <u>41,327</u>
David G. Streeter	Reg. No. <u>43,168</u>
Stephen Todd	Reg. No. <u>47,139</u>
P. Ben Wang	Reg. No. <u>41,420</u>

respectively and individually, as my patent attorneys and/or agents, with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith. Please address all communications to:

Docket No.: PF-0662 USN


**LEGAL DEPARTMENT**  
**INCYTE GENOMICS, INC.**  
**3160 PORTER DRIVE, PALO ALTO, CA 94304**

TEL: 650-855-0555


FAX: 650-849-8886 or 650-845-4166

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

1-00  
**First Joint Inventor:**

**Full name:**Y. Tom Tang**Signature:****Date:**Sept. 10, 2001**Citizenship**United States of America**Residence:**San Jose, California CA**P.O. Address:**4230 Ranwick Court  
San Jose, California 95118

2-00  
**Second Joint Inventor:**

**Full name:**Preeti Lal**Signature:****Date:**September 10, 2001**Citizenship**India**Residence:**Santa Clara, California CA**P.O. Address:**P.O. Box 5142  
Santa Clara, California 95056

Docket No.: PF-0662 USN

## Third Joint Inventor:

3-00  
Full name:Jennifer L. Hillman

Signature:

Jennifer L. Hillman

Date:

September 21, 2001

Citizenship

United States of America

Residence:

Mountain View, California CA

P.O. Address:

230 Monroe Drive, #17  
Mountain View, California 94040

## Fourth Joint Inventor:

4-00  
Full name:Henry Yue

Signature:

Henry Yue

Date:

September 24, 2001

Citizenship

United States of America

Residence:

Sunnyvale, California CA

P.O. Address:

826 Lois Avenue  
Sunnyvale, California 94087

## Fifth Joint Inventor:

5-00  
Full name:Yalda Azimzai

Signature:

Yalda Azimzai

Date:

September 13, 2001

Citizenship

United States of America

Residence:

Castro Valley, California CA

P.O. Address:

5518 Boulder Canyon Dr. Castro  
Valley, California 94552

Docket No.: PF-0662 USN

Sixth Joint Inventor:

6-00

Full name: Dyung Aina M. Lu  
 Signature: *Dyung Aina M. Lu*  
 Date: Sept 7, 2001  
 Citizenship: United States of America  
 Residence: San Jose, California CA  
 P.O. Address: 233 Coy Drive  
San Jose, California 95123

Seventh Joint Inventor:

7-00

Full name: Mariah R. Baughn  
 Signature: *Mariah R. Baughn*  
 Date: September 5, 2001  
 Citizenship: United States of America  
 Residence: San Leandro, California CA  
 P.O. Address: 14244 Santiago Road  
San Leandro, California 94577

Eighth Joint Inventor:

8-00

Full name: Bao Tran  
 Signature: \_\_\_\_\_  
 Date: \_\_\_\_\_, 2001  
 Citizenship: United States of America  
 Residence: Santa Clara, California CA  
 P.O. Address: 750 Salberg Avenue  
Santa Clara, California 95051

Docket No.: PF-0662 USN

Ninth Joint Inventor:

9-50  
Full name: Leo L. Shih  
Signature: \_\_\_\_\_  
Date: \_\_\_\_\_, 2001  
Citizenship: United States of America  
Residence: East Palo Alto, California CA  
P.O. Address: 908 O'Connor St.  
East Palo Alto, California 94303

Tenth Joint Inventor:

10-00  
Full name: Janice Au-Young  
Signature: Janice Au-Young  
Date: September 7, 2001  
Citizenship: United States of America  
Residence: Brisbane, California CA  
P.O. Address: 233 Golden Eagle Lane  
Brisbane, California 94005



Docket No.: PF-0662 USN

**DECLARATION AND POWER OF ATTORNEY FOR  
UNITED STATES PATENT APPLICATION**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name, and

I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if more than one name is listed below) of the subject matter which is claimed and for which a United States patent is sought on the invention entitled

**NUCLEIC-ACID BINDING PROTEINS**

the specification of which:

   / is attached hereto.

   / was filed on \_\_\_\_\_ as application Serial No. \_\_\_\_\_ and if this box contains an X    /, was amended on \_\_\_\_\_.

   / X / was filed as Patent Cooperation Treaty international application No. PCT/US00/02237 on January 28, 2000, this box contains an X    /, was amended on under Patent Cooperation Treaty Article 19 on \_\_\_\_\_ 2001, and if this box contains an X    /, was amended on \_\_\_\_\_.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge my duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim the benefit under Title 35, United States Code, §119 or §365(a)-(b) of any foreign application(s) for patent or inventor's certificate indicated below and of any Patent Cooperation Treaty international applications(s) designating at least one country other than the United States indicated below and have also identified below any foreign application(s) for patent or inventor's certificate and Patent Cooperation Treaty international application(s) designating at least one country other than the United States for the same subject matter and having a filing date before that of the application for said subject matter the priority of which is claimed:

**Docket No.: PF-0662 USN**

Country	Number	Filing Date	Priority Claimed
_____	_____	_____	// Yes // No
_____	_____	_____	// Yes // No

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

Application Serial No.	Filed	Status (Pending, Abandoned, Patented)
60/117,905	January 29, 1999	Expired
60/117,904	January 29, 1999	Expired

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in said prior application(s) in the manner required by the first paragraph of Title 35, United States Code §112, I acknowledge my duty to disclose material information as defined in Title 37 Code of Federal Regulations, §1.56(a) which occurred between the filing date(s) of the prior application(s) and the national or Patent Cooperation Treaty international filing date of this application:

Application Serial No.	Filed	Status (Pending, Abandoned, Patented)
_____	_____	_____

I hereby appoint the following:

Lucy J. Billings	Reg. No. 36,749
Michael C. Cerrone	Reg. No. 39,132
Diana Hamlet-Cox	Reg. No. 33,302
Richard C. Ekstrom	Reg. No. 37,027
Barrie D. Greene	Reg. No. 46,740
Lynn E. Murry	Reg. No. 42,918
Shirley A. Recipon	Reg. No. 47,016
Susan K. Sather	Reg. No. 44,316
Michelle M. Stempien	Reg. No. 41,327
David G. Streeter	Reg. No. 43,168

respectively and individually, as my patent attorneys and/or agents, with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith. Please address all communications to:

Docket No.: PF-0662 USN

LEGAL DEPARTMENT  
INCYTE GENOMICS, INC.  
3160 PORTER DRIVE, PALO ALTO, CA 94304

TEL: 650-855-0555 FAX: 650-849-8886 or 650-845-4166

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

<b>First Joint Inventor:</b>	<b>Full name:</b>	<u>Y. Tom Tang</u>
	<b>Signature:</b>	<u></u>
	<b>Date:</b>	<u>, 2001</u>
	<b>Citizenship</b>	<u>United States of America</u>
	<b>Residence:</b>	<u>San Jose, California</u>
	<b>P.O. Address:</b>	<u>4230 Ranwick Court</u> <u>San Jose, California 95118</u>

<b>Second Joint Inventor:</b>	<b>Full name:</b>	<u>Preeti Lal</u>
	<b>Signature:</b>	<u></u>
	<b>Date:</b>	<u>, 2001</u>
	<b>Citizenship</b>	<u>India</u>
	<b>Residence:</b>	<u>Santa Clara, California</u>
	<b>P.O. Address:</b>	<u>P.O. Box 5142</u> <u>Santa Clara, California 95056</u>

Docket No.: PF-0662 USN

**Third Joint Inventor:**

**Full name:** Jennifer L. Hillman  
**Signature:** \_\_\_\_\_  
**Date:** \_\_\_\_\_, 2001  
**Citizenship** United States of America  
**Residence:** Mountain View, California  
**P.O. Address:** 230 Monroe Drive, #17  
Mountain View, California 94040

**Fourth Joint Inventor:**

**Full name:** Henry Yue  
**Signature:** \_\_\_\_\_  
**Date:** \_\_\_\_\_, 2001  
**Citizenship** United States of America  
**Residence:** Sunnyvale, California  
**P.O. Address:** 826 Lois Avenue  
Sunnyvale, California 94087

**Fifth Joint Inventor:**

**Full name:** Yalda Azimzai  
**Signature:** \_\_\_\_\_  
**Date:** \_\_\_\_\_, 2001  
**Citizenship** United States of America  
**Residence:** Castro Valley, California  
**P.O. Address:** 5518 Boulder Canyon Dr. Castro  
Valley, California 94552

Docket No.: PF-0662 USN


## Sixth Joint Inventor:

Full name: Dyung Aina M. Lu  
Signature: \_\_\_\_\_  
Date: \_\_\_\_\_, 2001  
Citizenship: United States of America  
Residence: San Jose, California  
P.O. Address: 233 Coy Drive  
San Jose, California 95123

## Seventh Joint Inventor:

Full name: Mariah R. Baughn  
Signature: \_\_\_\_\_  
Date: \_\_\_\_\_, 2001  
Citizenship: United States of America  
Residence: San Leandro, California  
P.O. Address: 14244 Santiago Road  
San Leandro, California 94577

## Eighth Joint Inventor:

Full name: Bao Tran  
Signature:  \_\_\_\_\_  
Date: Oct 26, 2001  
Citizenship: United States of America  
Residence: Santa Clara, California  
P.O. Address: 750 Salberg Avenue  
Santa Clara, California 95051

Docket No.: PF-0662 USN

**Ninth Joint Inventor:**

**Full name:** Leo L. Shih  
**Signature:** \_\_\_\_\_  
**Date:** \_\_\_\_\_, 2001  
**Citizenship** United States of America  
**Residence:** East Palo Alto, California  
**P.O. Address:** 908 O'Connor St.  
East Palo Alto, California 94303

**Tenth Joint Inventor:**

**Full name:** Janice Au-Young  
**Signature:** \_\_\_\_\_  
**Date:** \_\_\_\_\_, 2001  
**Citizenship** United States of America  
**Residence:** Brisbane, California  
**P.O. Address:** 233 Golden Eagle Lane  
Brisbane, California 94005

Docket No.: PF-0662 USN

**DECLARATION AND POWER OF ATTORNEY FOR  
UNITED STATES PATENT APPLICATION**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name, and

I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if more than one name is listed below) of the subject matter which is claimed and for which a United States patent is sought on the invention entitled

**NUCLEIC-ACID BINDING PROTEINS**

the specification of which:

  /   is attached hereto.

  /   was filed on \_\_\_\_\_ as application Serial No. \_\_\_\_\_, and if this box contains an X   /  , was amended on \_\_\_\_\_.

  /   was filed as Patent Cooperation Treaty international application No. PCT/US00/02237 on January 28, 2000, this box contains an X   /  , was amended on under Patent Cooperation Treaty Article 19 on \_\_\_\_\_ 2001, and if this box contains an X   /  , was amended on \_\_\_\_\_.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge my duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim the benefit under Title 35, United States Code, §119 or §365(a)-(b) of any foreign application(s) for patent or inventor's certificate indicated below and of any Patent Cooperation Treaty international applications(s) designating at least one country other than the United States indicated below and have also identified below any foreign application(s) for patent or inventor's certificate and Patent Cooperation Treaty international application(s) designating at least one country other than the United States for the same subject matter and having a filing date before that of the application for said subject matter the priority of which is claimed:

## Docket No.: PF-0662 USN

Country	Number	Filing Date	Priority Claimed
_____	_____	_____	// Yes // No
_____	_____	_____	// Yes // No

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

Application Serial No.	Filed	Status (Pending, Abandoned, Patented)
60/117,905	January 29, 1999	Expired
60/117,904	January 29, 1999	Expired

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in said prior application(s) in the manner required by the first paragraph of Title 35, United States Code §112, I acknowledge my duty to disclose material information as defined in Title 37 Code of Federal Regulations, §1.56(a) which occurred between the filing date(s) of the prior application(s) and the national or Patent Cooperation Treaty international filing date of this application:

Application Serial No.	Filed	Status (Pending, Abandoned, Patented)
_____	_____	_____

I hereby appoint the following:

Lucy J. Billings	Reg. No. 36,749
Michael C. Cerrone	Reg. No. 39,132
Diana Hamlet-Cox	Reg. No. 33,302
Richard C. Ekstrom	Reg. No. 37,027
Barrie D. Greene	Reg. No. 46,740
Lynn E. Murry	Reg. No. 42,918
Shirley A. Recipon	Reg. No. 47,016
Susan K. Sather	Reg. No. 44,316
Michelle M. Stempien	Reg. No. 41,327
David G. Streeter	Reg. No. 43,168
P. Ben Wang	Reg. No. 41,420

respectively and individually, as my patent attorneys and/or agents, with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith. Please address all communications to:



Docket No.: PF-0662 USN

**LEGAL DEPARTMENT  
INCYTE GENOMICS, INC.  
3160 PORTER DRIVE, PALO ALTO, CA 94304**

**TEL: 650-855-0555 FAX: 650-849-8886 or 650-845-4166**

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

**First Joint Inventor:**

**Full name:** Y. Tom Tang  
**Signature:** \_\_\_\_\_  
**Date:** \_\_\_\_\_, 2001  
**Citizenship** United States of America  
**Residence:** San Jose, California  
**P.O. Address:** 4230 Ranwick Court  
San Jose, California 95118

**Second Joint Inventor:**

**Full name:** Preeti Lal  
**Signature:** \_\_\_\_\_  
**Date:** \_\_\_\_\_, 2001  
**Citizenship** India  
**Residence:** Santa Clara, California  
**P.O. Address:** P.O. Box 5142  
Santa Clara, California 95056

Docket No.: PF-0662 USN

**Third Joint Inventor:**

**Full name:** Jennifer L. Hillman  
**Signature:** \_\_\_\_\_  
**Date:** \_\_\_\_\_, 2001  
**Citizenship** United States of America  
**Residence:** Mountain View, California  
**P.O. Address:** 230 Monroe Drive, #17  
Mountain View, California 94040

**Fourth Joint Inventor:**

**Full name:** Henry Yue  
**Signature:** \_\_\_\_\_  
**Date:** \_\_\_\_\_, 2001  
**Citizenship** United States of America  
**Residence:** Sunnyvale, California  
**P.O. Address:** 826 Lois Avenue  
Sunnyvale, California 94087

**Fifth Joint Inventor:**

**Full name:** Yalda Azimzai  
**Signature:** \_\_\_\_\_  
**Date:** \_\_\_\_\_, 2001  
**Citizenship** United States of America  
**Residence:** Castro Valley, California  
**P.O. Address:** 5518 Boulder Canyon Dr. Castro  
Valley, California 94552

Docket No.: PF-0662 USN

## Sixth Joint Inventor:

Full name: Dyung Aina M. Lu  
Signature: \_\_\_\_\_  
Date: \_\_\_\_\_, 2001  
Citizenship: United States of America  
Residence: San Jose, California  
P.O. Address: 233 Coy Drive  
San Jose, California 95123

## Seventh Joint Inventor:


Full name: Mariah R. Baughn  
Signature: \_\_\_\_\_  
Date: \_\_\_\_\_, 2001  
Citizenship: United States of America  
Residence: San Leandro, California  
P.O. Address: 14244 Santiago Road  
San Leandro, California 94577

## Eighth Joint Inventor:

Full name: Bao Tran  
Signature: \_\_\_\_\_  
Date: \_\_\_\_\_, 2001  
Citizenship: United States of America  
Residence: Santa Clara, California  
P.O. Address: 750 Salberg Avenue  
Santa Clara, California 95051

Docket No.: PF-0662 USN

## Ninth Joint Inventor:

Full name: Leo L. Shih  
Signature:   
Date: October 27, 2001  
Citizenship: United States of America  
Residence: East Palo Alto, California  
P.O. Address: 908 O'Connor St.  
East Palo Alto, California 94303

## Tenth Joint Inventor:

Full name: Janice Au-Young  
Signature: \_\_\_\_\_  
Date: \_\_\_\_\_, 2001  
Citizenship: United States of America  
Residence: Brisbane, California  
P.O. Address: 233 Golden Eagle Lane  
Brisbane, California 94005

## SEQUENCE LISTING

&lt;110&gt; INCYTE PHARMACEUTICALS, INC.

TANG, Y. Tom  
 LAL, Preeti  
 HILLMAN, Jennifer L.  
 YUE, Henry  
 AZIMZAI, Yalda  
 LU, Aina M.D.  
 BAUGHN, Mariah R.  
 TRAN, Bao  
 SHIH, Leo L.  
 AU-YOUNG, Janice

&lt;120&gt; NUCLEIC ACID-BINDING PROTEINS

&lt;130&gt; PF-0662 PCT

&lt;140&gt; To Be Assigned

&lt;141&gt; Herewith

&lt;150&gt; 60/117,905; 60/117,904

&lt;151&gt; 1999-01-29; 1999-01-29

&lt;160&gt; 110

&lt;170&gt; PERL Program

&lt;210&gt; 1

&lt;211&gt; 754

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 025733CD1

&lt;400&gt; 1

```

Met Ala Ala Ala Gly Ser Arg Lys Arg Arg Leu Ala Glu Leu Thr
  1      5      10      15
Val Asp Glu Phe Leu Ala Ser Gly Phe Asp Ser Glu Ser Glu Ser
  20     25     30
Glu Ser Glu Asn Ser Pro Gln Ala Glu Thr Arg Glu Ala Arg Glu
  35     40     45
Ala Ala Arg Ser Pro Asp Lys Pro Gly Gly Ser Pro Ser Ala Ser
  50     55     60
Arg Arg Lys Gly Arg Ala Ser Glu His Lys Asp Gln Leu Ser Arg
  65     70     75
Leu Lys Asp Arg Asp Pro Glu Phe Tyr Lys Phe Leu Gln Glu Asn
  80     85     90
Asp Gln Ser Leu Leu Asn Phe Ser Asp Ser Asp Ser Ser Glu Glu
  95    100    105
Glu Glu Gly Pro Phe His Ser Leu Pro Asp Val Leu Glu Glu Ala
  110   115   120
Ser Glu Glu Glu Asp Gly Ala Glu Glu Gly Glu Asp Gly Asp Arg
  125   130   135
Val Pro Arg Gly Leu Lys Gly Lys Lys Asn Ser Val Pro Val Thr
  140   145   150
Val Ala Met Val Glu Arg Trp Lys Gln Ala Ala Lys Gln Arg Leu
  155   160   165

```

WO 00/44900

PCT/US00/02237

Thr	Pro	Lys	Leu	Phe	His	Glu	Val	Val	Gln	Ala	Phe	Arg	Ala	Ala	
				170					175					180	
Val	Ala	Thr	Thr	Arg	Gly	Asp	Gln	Glu	Ser	Ala	Glu	Ala	Asn	Lys	
				185					190					195	
Phe	Gln	Val	Thr	Asp	Ser	Ala	Ala	Phe	Asn	Ala	Leu	Val	Thr	Phe	
				200					205					210	
Cys	Ile	Arg	Asp	Leu	Ile	Gly	Cys	Leu	Gln	Lys	Leu	Leu	Phe	Gly	
				215					220					225	
Lys	Val	Ala	Lys	Asp	Ser	Ser	Arg	Met	Leu	Gln	Pro	Ser	Ser	Ser	
				230					235					240	
Pro	Leu	Trp	Gly	Lys	Leu	Arg	Val	Asp	Ile	Lys	Ala	Tyr	Leu	Gly	
				245					250					255	
Ser	Ala	Ile	Gln	Leu	Val	Ser	Cys	Leu	Ser	Glu	Thr	Thr	Val	Leu	
				260					265					270	
Ala	Ala	Val	Leu	Arg	His	Ile	Ser	Val	Leu	Val	Pro	Cys	Phe	Leu	
				275					280					285	
Thr	Phe	Pro	Lys	Gln	Cys	Arg	Met	Leu	Leu	Lys	Arg	Met	Val	Val	
				290					295					300	
Val	Trp	Ser	Thr	Gly	Glu	Glu	Ser	Leu	Arg	Val	Leu	Ala	Phe	Leu	
				305					310					315	
Val	Leu	Ser	Arg	Val	Cys	Arg	His	Lys	Lys	Asp	Thr	Phe	Leu	Gly	
				320					325					330	
Pro	Val	Leu	Lys	Gln	Met	Tyr	Ile	Thr	Tyr	Val	Arg	Asn	Cys	Lys	
				335					340					345	
Phe	Thr	Ser	Pro	Gly	Ala	Leu	Pro	Phe	Ile	Ser	Phe	Met	Gln	Trp	
				350					355					360	
Thr	Leu	Thr	Glu	Leu	Leu	Ala	Leu	Glu	Pro	Gly	Val	Ala	Tyr	Gln	
				365					370					375	
His	Ala	Phe	Leu	Tyr	Ile	Arg	Gln	Leu	Ala	Ile	His	Leu	Arg	Asn	
				380					385					390	
Ala	Met	Thr	Thr	Arg	Lys	Lys	Glu	Thr	Tyr	Gln	Ser	Val	Tyr	Asn	
				395					400					405	
Trp	Gln	Tyr	Val	His	Cys	Leu	Phe	Leu	Trp	Cys	Arg	Val	Leu	Ser	
				410					415					420	
Thr	Ala	Gly	Pro	Ser	Glu	Ala	Leu	Gln	Pro	Leu	Val	Tyr	Pro	Leu	
				425					430					435	
Ala	Gln	Val	Ile	Ile	Gly	Cys	Ile	Lys	Leu	Ile	Pro	Thr	Ala	Arg	
				440					445					450	
Phe	Tyr	Pro	Leu	Arg	Met	His	Cys	Ile	Arg	Ala	Leu	Thr	Leu	Leu	
				455					460					465	
Ser	Gly	Ser	Ser	Gly	Ala	Phe	Ile	Pro	Val	Leu	Pro	Phe	Ile	Leu	
				470					475					480	
Glu	Met	Phe	Gln	Gln	Val	Asp	Phe	Asn	Arg	Lys	Pro	Gly	Arg	Met	
				485					490					495	
Ser	Ser	Lys	Pro	Ile	Asn	Phe	Ser	Val	Ile	Leu	Lys	Leu	Ser	Asn	
				500					505					510	
Val	Asn	Leu	Gln	Glu	Lys	Ala	Tyr	Arg	Asp	Gly	Leu	Val	Glu	Gln	
				515					520					525	
Leu	Tyr	Asp	Leu	Thr	Leu	Glu	Tyr	Leu	His	Ser	Gln	Ala	His	Cys	
				530					535					540	
Ile	Gly	Phe	Pro	Glu	Leu	Val	Leu	Pro	Val	Val	Leu	Gln	Leu	Lys	
				545					550					555	
Ser	Phe	Leu	Arg	Glu	Cys	Lys	Val	Ala	Asn	Tyr	Cys	Arg	Gln	Val	
				560					565					570	
Gln	Gln	Leu	Leu	Gly	Lys	Val	Gln	Glu	Asn	Ser	Ala	Tyr	Ile	Cys	
				575					580					585	
Ser	Arg	Arg	Gln	Arg	Val	Ser	Phe	Gly	Val	Ser	Glu	Gln	Gln	Ala	
				590					595					600	
Val	Glu	Ala	Trp	Glu	Lys	Leu	Thr	Arg	Glu	Glu	Gly	Thr	Pro	Leu	
				605					610					615	

Thr	Leu	Tyr	Tyr	Ser	His	Trp	Arg	Lys	Leu	Arg	Asp	Arg	Glu	Ile	
				620					625					630	
Gln	Leu	Glu	Ile	Ser	Gly	Lys	Glu	Arg	Leu	Glu	Asp	Leu	Asn	Phe	
				635					640					645	
Pro	Glu	Ile	Lys	Arg	Arg	Lys	Met	Ala	Asp	Arg	Lys	Asp	Glu	Asp	
				650					655					660	
Arg	Lys	Gln	Phe	Lys	Asp	Leu	Phe	Asp	Leu	Asn	Ser	Ser	Glu	Glu	
				665					670					675	
Asp	Asp	Thr	Glu	Gly	Phe	Ser	Glu	Arg	Gly	Ile	Leu	Arg	Pro	Leu	
				680					685					690	
Ser	Thr	Arg	His	Gly	Val	Glu	Asp	Asp	Glu	Glu	Asp	Glu	Glu	Glu	
				695					700					705	
Gly	Glu	Glu	Asp	Ser	Ser	Asn	Ser	Glu	Gly	Glu	Trp	Ser	Trp	Asp	
				710					715					720	
Gly	Asp	Pro	Asp	Ala	Glu	Ala	Gly	Leu	Ala	Pro	Gly	Glu	Leu	Gln	
				725					730					735	
Gln	Leu	Ala	Gln	Gly	Pro	Glu	Asp	Glu	Leu	Glu	Asp	Leu	Gln	Leu	
				740					745					750	
Ser	Glu	Asp	Asp												

<210> 2  
 <211> 593  
 <212> PRT  
 <213> Homo sapiens  
  
 <220>  
 <221> misc-feature  
 <223> Incyte ID No.: 079702CD1

<400> 2  
 Met Arg Asp Ser Thr Gly Ala Gly Asn Ser Leu Val His Lys Arg  
 1 5 10 15  
 Ser Pro Leu Arg Arg Asn Gln Lys Thr Pro Thr Ser Leu Thr Lys  
 20 25 30  
 Leu Ser Leu Gln Asp Gly His Lys Ala Lys Lys Pro Ala Cys Lys  
 35 40 45  
 Phe Glu Glu Gly Gln Asp Val Leu Ala Arg Trp Ser Asp Gly Leu  
 50 55 60  
 Phe Tyr Leu Gly Thr Ile Lys Lys Ile Asn Ile Leu Lys Gln Ser  
 65 70 75  
 Cys Phe Ile Ile Phe Glu Asp Ser Ser Lys Ser Trp Val Leu Trp  
 80 85 90  
 Lys Asp Ile Gln Thr Gly Ala Thr Gly Ser Gly Glu Met Val Cys  
 95 100 105  
 Thr Ile Cys Gln Glu Glu Tyr Ser Glu Ala Pro Asn Glu Met Val  
 110 115 120  
 Ile Cys Asp Lys Cys Gly Gln Gly Tyr His Gln Leu Cys His Thr  
 125 130 135  
 Pro His Ile Asp Ser Ser Val Ile Asp Ser Asp Glu Lys Trp Leu  
 140 145 150  
 Cys Arg Gln Cys Val Phe Ala Thr Thr Thr Lys Arg Gly Gly Ala  
 155 160 165  
 Leu Lys Lys Gly Pro Asn Ala Lys Ala Leu Gln Val Met Lys Gln  
 170 175 180  
 Thr Leu Pro Tyr Ser Val Ala Asp Leu Glu Trp Asp Ala Gly His  
 185 190 195  
 Lys Thr Asn Val Gln Gln Cys Tyr Cys Tyr Cys Gly Gly Pro Gly  
 200 205 210  
 Asp Trp Tyr Leu Lys Met Leu Gln Cys Cys Lys Cys Lys Gln Trp

Phe His Glu Ala	215	220	225
Cys Val Gln Cys Leu	230	235	240
Gly Asp Arg Phe Tyr Thr Phe Ile Cys Ser Val Cys Ser Ser Gly	245	250	255
Pro Glu Tyr Leu Lys Arg Leu Pro Leu Gln Trp Val Asp Ile Ala	260	265	270
His Leu Cys Leu Tyr Asn Leu Ser Val Ile His Lys Lys Lys Tyr	275	280	285
Phe Asp Ser Glu Leu Glu Leu Met Thr Tyr Ile Asn Glu Asn Trp	290	295	300
Asp Arg Leu His Pro Gly Glu Leu Ala Asp Thr Pro Lys Ser Glu	305	310	315
Arg Tyr Glu His Val Leu Glu Ala Leu Asn Asp Tyr Lys Thr Met	320	325	330
Phe Met Ser Gly Lys Glu Ile Lys Lys Lys Lys His Leu Phe Gly	335	340	345
Leu Arg Ile Arg Val Pro Pro Val Pro Pro Asn Val Ala Phe Lys	350	355	360
Ala Glu Lys Glu Pro Glu Gly Thr Ser His Glu Phe Lys Ile Lys	365	370	375
Gly Arg Lys Ala Ser Lys Pro Ile Ser Asp Ser Arg Glu Val Ser	380	385	390
Asn Gly Ile Glu Lys Lys Lys Lys Lys Lys Ser Val Gly Arg Pro	395	400	405
Pro Gly Pro Tyr Thr Arg Lys Met Ile Gln Lys Thr Ala Glu Pro	410	415	420
Leu Leu Asp Lys Glu Ser Ile Ser Glu Asn Pro Thr Leu Asp Leu	425	430	435
Pro Cys Ser Ile Gly Arg Thr Glu Gly Thr Ala His Ser Ser Asn	440	445	450
Thr Ser Asp Val Asp Phe Thr Gly Ala Ser Ser Ala Lys Glu Thr	455	460	465
Thr Ser Ser Ser Ile Ser Arg His Tyr Gly Leu Ser Asp Ser Arg	470	475	480
Lys Arg Thr Arg Thr Gly Arg Ser Trp Pro Ala Ala Ile Pro His	485	490	495
Leu Arg Arg Arg Arg Gly Arg Leu Pro Arg Arg Ala Leu Gln Thr	500	505	510
Gln Asn Ser Glu Ile Val Lys Asp Asp Glu Gly Lys Glu Asp Tyr	515	520	525
Gln Phe Asp Glu Leu Asn Thr Glu Ile Leu Asn Asn Leu Ala Asp	530	535	540
Gln Glu Leu Gln Leu Asn His Leu Lys Asn Ser Ile Thr Ser Tyr	545	550	555
Phe Gly Ala Ala Gly Arg Ile Ala Cys Gly Glu Lys Tyr Arg Val	560	565	570
Leu Ala Arg Arg Val Thr Leu Asp Gly Lys Val Gln Tyr Leu Val	575	580	585
Glu Trp Glu Gly Ala Thr Ala Ser	590		

<210> 3  
 <211> 534  
 <212> PRT  
 <213> Homo sapiens  
 <220>



&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 116208CD1

&lt;400&gt; 3

```

Met Arg Ala Leu His 5 Leu Leu Lys Ser Gly Cys Ser Pro Ala Val
1 5 10 15
Gln Ile Lys Ile Arg 20 Glu Leu Tyr Arg Arg Arg Tyr Pro Arg Thr
20 25 30
Leu Glu Gly Leu Ser Asp Leu Ser Thr Ile Lys Ser Ser Val Phe
35 40 45
Ser Leu Asp Gly Gly Ser Ser Pro Val Glu Pro Asp Leu Ala Val
50 55 60
Ala Gly Ile His Ser Leu Pro Ser Thr Ser Val Thr Pro His Ser
65 70 75
Pro Ser Ser Pro Val Gly Ser Val Leu Leu Gln Asp Thr Lys Pro
80 85 90
Thr Phe Glu Met Gln Gln Pro Ser Pro Pro Ile Pro Pro Val His
95 100 105
Pro Asp Val Gln Leu Lys Asn Leu Pro Phe Tyr Asp Val Leu Asp
110 115 120
Val Leu Ile Lys Pro Thr Ser Leu Val Gln Ser Ser Ile Gln Arg
125 130 135
Phe Gln Glu Lys Phe Phe Ile Phe Ala Leu Thr Pro Gln Gln Val
140 145 150
Arg Glu Ile Cys Ile Ser Arg Asp Phe Leu Pro Gly Gly Arg Arg
155 160 165
Asp Tyr Thr Val Gln Val Gln Leu Arg Leu Cys Leu Ala Glu Thr
170 175 180
Ser Cys Pro Gln Glu Asp Asn Tyr Pro Asn Ser Leu Cys Ile Lys
185 190 195
Val Asn Gly Lys Leu Phe Pro Leu Pro Gly Tyr Ala Pro Pro Pro
200 205 210
Lys Asn Gly Ile Glu Gln Lys Arg Pro Gly Arg Pro Leu Asn Ile
215 220 225
Thr Ser Leu Val Arg Leu Ser Ser Ala Val Pro Asn Gln Ile Ser
230 235 240
Ile Ser Trp Ala Ser Glu Ile Gly Lys Asn Tyr Ser Met Ser Val
245 250 255
Tyr Leu Val Arg Gln Leu Thr Ser Ala Met Leu Leu Gln Arg Leu
260 265 270
Lys Met Lys Gly Ile Arg Asn Pro Asp His Ser Arg Ala Leu Ser
275 280 285
Lys Glu Lys Leu Thr Ala Asp Pro Asp Ser Glu Ile Ala Thr Thr
290 295 300
Ser Leu Arg Val Ser Leu Met Cys Pro Leu Gly Lys Met Arg Leu
305 310 315
Thr Ile Pro Cys Arg Ala Val Thr Cys Thr His Leu Gln Cys Phe
320 325 330
Asp Ala Ala Leu Tyr Leu Gln Met Asn Glu Lys Lys Pro Thr Trp
335 340 345
Ile Cys Pro Val Cys Asp Lys Lys Ala Ala Tyr Glu Ser Leu Ile
350 355 360
Leu Asp Gly Leu Phe Met Glu Ile Leu Asn Asp Cys Ser Asp Val
365 370 375
Asp Glu Ile Lys Phe Gln Glu Asp Gly Ser Trp Cys Pro Met Arg
380 385 390
Pro Lys Lys Glu Ala Met Lys Val Ser Ser Gln Pro Cys Thr Lys
395 400 405
Ile Glu Ser Ser Ser Val Leu Ser Lys Pro Cys Ser Val Thr Val
410 415 420
Ala Ser Glu Ala Ser Lys Lys Lys Val Asp Val Ile Asp Leu Thr
425 430 435

```

Ile Glu Ser Ser Asp Glu Glu Glu Asp Pro Pro Ala Lys Arg  
 440 445 450  
 Lys Cys Ile Phe Met Ser Glu Thr Gln Ser Ser Pro Thr Lys Gly  
 455 460 465  
 Val Leu Met Tyr Gln Pro Ser Ser Val Arg Val Pro Ser Val Thr  
 470 475 480  
 Ser Val Asp Pro Ala Ala Ile Pro Pro Ser Leu Thr Asp Tyr Ser  
 485 490 495  
 Val Pro Phe His His Thr Pro Ile Ser Ser Met Ser Ser Asp Leu  
 500 505 510  
 Pro Gly Glu Gln Arg Arg Asn Asp Ile Asn Asn Glu Leu Lys Leu  
 515 520 525  
 Gly Thr Ser Ser Asp Thr Val Gln Gln  
 530

&lt;210&gt; 4

&lt;211&gt; 255

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc feature

&lt;223&gt; Incyte ID No.: 179261CD1

&lt;400&gt; 4

Met Lys Ile Ile Ser Glu Asn Val Pro Ser Tyr Lys Thr His Glu  
 1 5 10 15  
 Ser Leu Thr Leu Pro Arg Arg Thr His Asp Ser Glu Lys Pro Tyr  
 20 25 30  
 Glu Tyr Lys Glu Tyr Glu Lys Val Phe Ser Cys Asp Leu Glu Phe  
 35 40 45  
 Asp Glu Tyr Gln Lys Ile His Thr Gly Gly Lys Asn Tyr Glu Cys  
 50 55 60  
 Asn Gln Cys Trp Lys Thr Phe Gly Ile Asp Asn Ser Ser Met Leu  
 65 70 75  
 Gln Leu Asn Ile His Thr Gly Val Lys Pro Cys Lys Tyr Met Glu  
 80 85 90  
 Tyr Gly Asn Thr Cys Ser Phe Tyr Lys Asp Phe Asn Val Tyr Gln  
 95 100 105  
 Lys Ile His Asn Glu Lys Phe Tyr Lys Cys Lys Glu Tyr Arg Arg  
 110 115 120  
 Thr Phe Glu Arg Val Gly Lys Val Thr Pro Leu Gln Arg Val His  
 125 130 135  
 Asp Gly Glu Lys His Phe Glu Cys Ser Phe Cys Gly Lys Ser Phe  
 140 145 150  
 Arg Val His Ala Gln Leu Thr Arg His Gln Lys Ile His Thr Asp  
 155 160 165  
 Glu Lys Thr Tyr Lys Cys Met Glu Cys Gly Lys Asp Phe Arg Phe  
 170 175 180  
 His Ser Gln Leu Thr Glu His Gln Arg Ile His Thr Gly Glu Lys  
 185 190 195  
 Pro Tyr Lys Cys Met His Cys Glu Lys Val Phe Arg Ile Ser Ser  
 200 205 210  
 Gln Leu Ile Glu His Gln Arg Ile His Thr Gly Glu Lys Pro Tyr  
 215 220 225  
 Ala Cys Lys Glu Cys Gly Lys Ala Phe Gly Val Cys Arg Glu Leu  
 230 235 240  
 Ala Arg His Gln Arg Ile His Thr Gly Lys Tyr Cys Gly Trp Ile  
 245 250 255

<210> 5  
 <211> 562  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc feature  
 <223> Incyte ID No.: 259161CD1

<400> 5  
 Met Ala Ser Val Ser Ala Leu Thr Glu Glu Leu Asp Ser Ile Thr  
 1 5 10 15  
 Ser Glu Leu His Ala Val Glu Ile Gln Ile Gln Glu Leu Thr Glu  
 20 25 30  
 Arg Gln Gln Glu Leu Ile Gln Lys Lys Lys Val Leu Thr Lys Lys  
 35 40 45  
 Ile Lys Gln Cys Leu Glu Asp Ser Asp Ala Gly Ala Ser Asn Glu  
 50 55 60  
 Tyr Asp Ser Ser Pro Ala Ala Trp Asn Lys Glu Asp Phe Pro Trp  
 65 70 75  
 Ser Gly Lys Val Lys Asp Ile Leu Gln Asn Val Phe Lys Leu Glu  
 80 85 90  
 Lys Phe Arg Pro Leu Gln Leu Glu Thr Ile Asn Val Thr Met Ala  
 95 100 105  
 Gly Lys Glu Val Phe Leu Val Met Pro Thr Gly Gly Gly Lys Ser  
 110 115 120  
 Leu Cys Tyr Gln Leu Pro Ala Leu Cys Ser Asp Gly Phe Thr Leu  
 125 130 135  
 Val Ile Cys Pro Leu Ile Ser Leu Met Glu Asp Gln Leu Met Val  
 140 145 150  
 Leu Lys Gln Leu Gly Ile Ser Ala Thr Met Leu Asn Ala Ser Ser  
 155 160 165  
 Ser Lys Glu His Val Lys Trp Val His Ala Glu Met Val Asn Lys  
 170 175 180  
 Asn Ser Glu Leu Lys Leu Ile Tyr Val Thr Pro Glu Lys Ile Ala  
 185 190 195  
 Lys Ser Lys Met Phe Met Ser Arg Leu Glu Lys Ala Tyr Glu Ala  
 200 205 210  
 Arg Arg Phe Thr Arg Ile Ala Val Asp Glu Val His Cys Cys Ser  
 215 220 225  
 Gln Trp Gly His Asp Phe Arg Pro Asp Tyr Lys Ala Leu Gly Ile  
 230 235 240  
 Leu Lys Arg Gln Phe Pro Asn Ala Ser Leu Ile Gly Leu Thr Ala  
 245 250 255  
 Thr Ala Thr Asn His Val Leu Thr Asp Ala Gln Lys Ile Leu Cys  
 260 265 270  
 Ile Glu Lys Cys Phe Thr Phe Thr Ala Ser Phe Asn Lys Pro Asp  
 275 280 285  
 Val Arg Phe Val Ile His His Ser Met Ser Lys Ser Met Glu Asn  
 290 295 300  
 Tyr Tyr Gln Glu Ser Gly Arg Ala Gly Arg Asp Asp Met Lys Ala  
 305 310 315  
 Asp Cys Ile Leu Tyr Tyr Gly Phe Gly Asp Ile Phe Arg Ile Ser  
 320 325 330  
 Ser Met Val Val Met Glu Asn Val Gly Gln Gln Lys Leu Tyr Glu  
 335 340 345  
 Met Val Ser Tyr Cys Gln Asn Ile Ser Lys Cys Arg Arg Val Leu  
 350 355 360  
 Met Ala Gln His Phe Asp Glu Val Trp Asn Ser Glu Ala Cys Asn

Lys Met Cys Asp	365	Asn Cys Cys Lys Asp	370	Ala Phe Glu Arg	375
	380		385		390
Asn Ile Thr Glu	395	Tyr Cys Arg Asp Leu	400	Ile Lys Ile Leu Lys	405
	410		415		420
Ala Glu Glu Leu	425	Asn Glu Lys Leu Thr	430	Pro Leu Lys Leu Ile	435
	440		445		450
Ser Trp Met Gly	455	Lys Gly Ala Ala Lys	460	Leu Arg Val Ala Gly	465
	470		475		480
Val Ala Pro Thr	485	Leu Pro Arg Glu Asp	490	Glu Lys Ile Ile Ala	495
	500		505		510
His Phe Leu Ile	515	Gln Gln Tyr Leu Lys	520	Glu Asp Tyr Ser Phe	525
	530		535		540
Ala Tyr Ala Thr	545	Ile Ser Tyr Leu Lys	550	Ile Gly Pro Lys Ala	555
Leu Leu Asn Asn		Glu Ala His Ala Ile		Thr Met Gln Val Thr	
Ser Thr Gln Asn		Ser Phe Arg Ala Glu		Ser Ser Gln Thr Cys	
Ser Glu Gln Gly		Asp Lys Lys Ile Gly		Gly Lys Lys Phe Gln	
Thr Ser Arg Arg		Arg Leu Gln Thr Cys		Phe Ser Asn Leu Val	
Arg Ile Gln Glu		Leu Arg Lys Glu Lys		Met Met Pro Asp	
Thr Val Thr Lys		Phe Ser Asn			

&lt;210&gt; 6

&lt;211&gt; 432

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 320087CD1

&lt;400&gt; 6

Met Ser Glu Leu Lys Asp Cys Pro Leu	Gln Phe His Asp Phe Lys
1	15
Ser Val Asp His Leu Lys Val Cys Pro	Arg Tyr Thr Ala Val Leu
20	30
Ala Arg Ser Glu Asp Asp Gly Ile Gly	Ile Glu Glu Leu Asp Thr
35	45
Leu Gln Leu Glu Leu Glu Thr Leu Leu	Ser Ser Ala Ser Arg Arg
50	60
Leu Arg Val Leu Glu Ala Glu Thr Gln	Ile Leu Thr Asp Trp Gln
65	75
Asp Lys Lys Gly Asp Arg Arg Phe Leu	Lys Leu Gly Arg Asp His
80	90
Glu Leu Gly Ala Pro Pro Lys His Gly	Lys Pro Lys Lys Gln Lys
95	105
Leu Glu Gly Lys Ala Gly His Gly Pro	Gly Pro Gly Pro Gly Arg
110	120
Pro Lys Ser Lys Asn Leu Gln Pro Lys	Ile Gln Glu Tyr Glu Phe
125	135
Thr Asp Asp Pro Ile Asp Val Pro Arg	Ile Pro Lys Asn Asp Ala
140	150

```

Pro Asn Arg Phe Trp Ala Ser Val Glu Pro Tyr Cys Ala Asp Ile
155 160 165
Thr Ser Glu Glu Val Arg Thr Leu Glu Glu Leu Leu Lys Pro Pro
170 175 180
Glu Asp Glu Ala Glu His Tyr Lys Ile Pro Pro Leu Gly Lys His
185 190 195
Tyr Ser Gln Arg Trp Ala Gln Glu Asp Leu Leu Glu Glu Gln Lys
200 205 210
Asp Gly Ala Arg Ala Ala Ala Val Ala Asp Lys Lys Lys Gly Leu
215 220 225
Met Gly Pro Leu Thr Glu Leu Asp Thr Lys Asp Val Asp Ala Leu
230 235 240
Leu Lys Lys Ser Glu Ala Gln His Glu Gln Pro Glu Asp Gly Cys
245 250 255
Pro Phe Gly Ala Leu Thr Gln Arg Leu Leu Gln Ala Leu Val Glu
260 265 270
Glu Asn Ile Ile Ser Pro Met Glu Asp Ser Pro Ile Pro Asp Met
275 280 285
Ser Gly Lys Glu Ser Gly Ala Asp Gly Ala Ser Thr Ser Pro Arg
290 295 300
Asn Gln Asn Lys Pro Phe Ser Val Pro His Thr Lys Ser Leu Glu
305 310 315
Ser Arg Ile Lys Glu Glu Leu Ile Ala Gln Gly Leu Leu Glu Ser
320 325 330
Glu Asp Arg Pro Ala Glu Asp Ser Glu Asp Glu Val Leu Ala Glu
335 340 345
Leu Arg Lys Arg Gln Ala Glu Leu Lys Ala Leu Ser Ala His Asn
350 355 360
Arg Thr Lys Lys His Asp Leu Leu Arg Leu Ala Lys Glu Glu Val
365 370 375
Ser Arg Gln Glu Leu Arg Gln Arg Val Arg Met Ala Asp Asn Glu
380 385 390
Val Met Asp Ala Phe Arg Lys Ile Met Ala Ala Arg Gln Lys Lys
395 400 405
Arg Thr Pro Thr Lys Lys Glu Lys Asp Gln Ala Trp Lys Thr Leu
410 415 420
Lys Glu Arg Glu Ser Ile Leu Lys Leu Leu Asp Gly
425 430

```

<210> 7  
 <211> 799  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> 491271CD1

```

<400> 7
Met Pro Ser Gln Asn Tyr Asp Leu Pro Gln Lys Lys Gln Glu Lys
1 5 10 15
Met Thr Lys Phe Gln Glu Ala Val Thr Phe Lys Asp Val Ala Val
20 25 30
Val Phe Ser Arg Glu Glu Leu Arg Leu Leu Asp Leu Thr Gln Arg
35 40 45
Lys Leu Tyr Arg Asp Val Met Val Glu Asn Phe Lys Asn Leu Val
50 55 60
Ala Val Gly His Leu Pro Phe Gln Pro Asp Met Val Ser Gln Leu
65 70 75

```

Glu Ala Glu Glu Lys Leu Trp Met Met Glu Thr Glu Thr Gln Arg  
 80 85 90  
 Ser Ser Lys His Gln Asn Lys Met Glu Thr Leu Gln Lys Phe Ala  
 95 100 105  
 Leu Lys Tyr Leu Ser Asn Gln Glu Leu Ser Cys Trp Gln Ile Trp  
 110 115 120  
 Lys Gln Val Ala Ser Glu Leu Thr Arg Cys Leu Gln Gly Lys Ser  
 125 130 135  
 Ser Gln Leu Leu Gln Gly Asp Ser Ile Gln Val Ser Glu Asn Glu  
 140 145 150  
 Asn Asn Ile Met Asn Pro Lys Gly Asp Ser Pro Ile Tyr Ile Glu  
 155 160 165  
 Asn Gln Glu Phe Pro Phe Trp Arg Thr Gln His Ser Cys Gly Asn  
 170 175 180  
 Thr Tyr Leu Ser Glu Ser Gln Ile Gln Ser Arg Gly Lys Gln Ile  
 185 190 195  
 Asp Val Lys Asn Asn Leu Gln Ile Arg Glu Asp Phe Val Lys Lys  
 200 205 210  
 Ser Pro Phe His Glu His Ile Lys Thr Asp Thr Glu Pro Lys Pro  
 215 220 225  
 Cys Lys Gly Asn Glu Tyr Gly Lys Ile Ile Ser Asp Gly Ser Asn  
 230 235 240  
 Gln Lys Leu Pro Leu Gly Glu Lys Pro His Pro Cys Gly Glu Cys  
 245 250 255  
 Gly Arg Gly Phe Ser Tyr Ser Pro Arg Leu Pro Leu His Pro Asn  
 260 265 270  
 Val His Thr Gly Glu Lys Cys Phe Ser Gln Ser Ser His Leu Arg  
 275 280 285  
 Thr His Gln Arg Ile His Pro Gly Glu Lys Leu Asn Arg Cys His  
 290 295 300  
 Glu Ser Gly Asp Cys Phe Asn Lys Ser Ser Phe His Ser Tyr Gln  
 305 310 315  
 Ser Asn His Thr Gly Glu Lys Ser Tyr Arg Cys Asp Ser Cys Gly  
 320 325 330  
 Lys Gly Phe Ser Ser Ser Thr Gly Leu Ile Ile His Tyr Arg Thr  
 335 340 345  
 His Thr Gly Glu Lys Pro Tyr Lys Cys Glu Glu Cys Gly Lys Cys  
 350 355 360  
 Phe Ser Gln Ser Ser Asn Phe Gln Cys His Gln Arg Val His Thr  
 365 370 375  
 Glu Glu Lys Pro Tyr Lys Cys Glu Glu Cys Gly Lys Gly Phe Gly  
 380 385 390  
 Trp Ser Val Asn Leu Arg Val His Gln Arg Val His Arg Gly Glu  
 395 400 405  
 Lys Pro Tyr Lys Cys Glu Glu Cys Gly Lys Gly Phe Thr Gln Ala  
 410 415 420  
 Ala His Phe His Ile His Gln Arg Val His Thr Gly Glu Lys Pro  
 425 430 435  
 Tyr Lys Cys Asp Val Cys Gly Lys Gly Phe Ser His Asn Ser Pro  
 440 445 450  
 Leu Ile Cys His Arg Arg Val His Thr Gly Glu Lys Pro Tyr Lys  
 455 460 465  
 Cys Glu Ala Cys Gly Lys Gly Phe Thr Arg Asn Thr Asp Leu His  
 470 475 480  
 Ile His Phe Arg Val His Thr Gly Glu Lys Pro Tyr Lys Cys Lys  
 485 490 495  
 Glu Cys Gly Lys Gly Phe Ser Gln Ala Ser Asn Leu Gln Val His  
 500 505 510  
 Gln Asn Val His Thr Gly Glu Lys Arg Phe Lys Cys Glu Thr Cys  
 515 520 525  
 Gly Lys Gly Phe Ser Gln Ser Ser Lys Leu Gln Thr His Gln Arg

Val His Thr Gly	530	535	540
Glu Lys Pro Tyr Arg	545	Cys Asp Val Cys Gly	Lys
Asp Phe Ser Tyr	560	550	555
Ser Ser Asn Leu Lys	565	Leu His Gln Val Ile	His
Thr Gly Glu Lys	575	565	570
Pro Tyr Lys Cys Glu	580	Cys Gly Lys Gly	Phe
Ser Trp Arg Ser	590	580	585
Asn Leu His Ala His	595	Gln Arg Val His Ser	Gly
Glu Lys Pro Tyr	605	595	600
Lys Cys Glu Gln Cys	610	Asp Lys Ser Phe Ser	Gln
Ala Ile Asp Phe	620	610	615
Arg Val His Gln Arg	625	Val His Thr Gly Glu	Lys
Pro Tyr Lys Cys	635	625	630
Gly Val Cys Gly Lys	640	Gly Phe Ser Gln Ser	Ser
Gly Leu Gln Ser	650	640	645
His Gln Arg Val His	655	Thr Gly Glu Lys Pro	Tyr
Lys Cys Asp Val	665	655	660
Cys Gly Lys Gly Phe	670	Arg Tyr Ser Ser Gln	Phe
Ile Tyr His Gln	680	670	675
Arg Gly His Thr Gly	685	Glu Lys Pro Tyr Lys	Cys
Glu Glu Cys Gly	695	685	690
Lys Gly Phe Gly Arg	700	Ser Leu Asn Leu Arg	His
His Gln Arg Val	710	700	705
His Thr Gly Glu Lys	715	Pro His Ile Cys Glu	Glu
Cys Gly Lys Ala	725	715	720
Phe Ser Leu Pro Ser	730	Leu Arg Val His Leu	Leu
Gly Val His Thr	740	730	735
Arg Glu Lys Leu Phe	745	Lys Cys Glu Glu Cys	Gly
Lys Gly Phe Ser	755	745	750
Gln Ser Ala Arg Leu	760	Glu Ala His Gln Arg	Val
His Thr Gly Glu Lys	770	760	765
Pro Tyr Lys Cys	775	Asp Ile Cys Asp Lys	Asp
Phe Arg His Arg	785	775	780
Ser Arg Leu Thr Tyr	790	His Gln Lys Val His	Thr
Gly Lys Lys Leu		790	795

&lt;210&gt; 8

&lt;211&gt; 137

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 585172CD1

&lt;400&gt; 8

Met Leu Ser Gly Arg Leu Val Leu Gly Leu Val Ser Met Ala Gly	1	5	10	15
Arg Val Cys Leu Cys Gln Gly Ser Ala Gly Ser Gly Ala Ile Gly	20	25	30	35
Pro Val Glu Ala Ala Ile Arg Thr Lys Leu Glu Glu Ala Leu Ser	40	45	50	55
Pro Glu Val Leu Glu Leu Arg Asn Glu Ser Gly Gly His Ala Val	60	65	70	75
Pro Pro Gly Ser Glu Thr His Phe Arg Val Ala Val Val Ser Ser	80	85	90	95
Arg Phe Glu Gly Leu Ser Pro Leu Gln Arg His Arg Leu Val His	100	105	110	115

Ala	Ala	Leu	Ala	80	Glu	Glu	Leu	Gly	Gly	85	Pro	Val	His	Ala	Leu	Ala	90
				95						100							105
Ile	Gln	Ala	Arg	Thr	Pro	Ala	Gln	Trp	Arg	Glu	Asn	Ser	Gln	Leu			
				110						115							120
Asp	Thr	Ser	Pro	Pro	Cys	Leu	Gly	Gly	Asn	Lys	Lys	Thr	Leu	Gly			
				125						130							135
Thr	Pro																

<210> 9  
 <211> 230  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 615200CD1

<400> 9  
 Met Val Gly Ala Gly Ile Ser Thr Pro Ser Gly Ile Pro Asp Phe  
 1 5 10 15  
 Arg Ser Pro Gly Ser Gly Leu Tyr Ser Asn Leu Gln Gln Tyr Asp  
 20 25 30  
 Leu Pro Tyr Pro Glu Ala Ile Phe Glu Leu Pro Phe Phe Phe His  
 35 40 45  
 Asn Pro Lys Pro Phe Phe Thr Leu Ala Lys Glu Leu Tyr Pro Gly  
 50 55 60  
 Asn Tyr Lys Pro Asn Ile Thr His Tyr Phe Leu Arg Leu Leu His  
 65 70 75  
 Asp Lys Gly Leu Leu Leu Arg Leu Tyr Thr Gln Asn Ile Asp Gly  
 80 85 90  
 Leu Glu Arg Val Ser Gly Ile Pro Ala Ser Lys Leu Val Glu Ala  
 95 100 105  
 His Gly Thr Phe Ala Ser Ala Thr Cys Thr Val Cys Gln Arg Pro  
 110 115 120  
 Phe Pro Gly Glu Asp Ile Arg Ala Asp Val Met Ala Asp Arg Val  
 125 130 135  
 Pro Arg Cys Pro Val Cys Thr Gly Val Val Lys Pro Asp Ile Val  
 140 145 150  
 Phe Phe Gly Glu Pro Leu Pro Gln Arg Phe Leu Leu His Val Val  
 155 160 165  
 Asp Phe Pro Met Ala Asp Leu Leu Leu Ile Leu Gly Thr Ser Leu  
 170 175 180  
 Glu Val Glu Pro Phe Ala Ser Leu Thr Glu Ala Val Arg Thr Gln  
 185 190 195  
 Phe Pro Asp Cys Ser Ser Thr Gly Thr Trp Trp Gly Pro Trp Leu  
 200 205 210  
 Gly Ile Leu Ala Ala Gly Thr Trp Pro Ser Trp Gly Thr Trp Phe  
 215 220 225  
 Thr Ala Trp Lys Ala  
 230

<210> 10  
 <211> 446  
 <212> PRT  
 <213> Homo sapiens



<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 997067CD1

<400> 10

Met Glu Thr Gln Ala	Asp Leu Val Ser	Gln Glu Pro Gln Ala Leu	15
1	5	10	15
Leu Asp Ser Ala Leu	Pro Ser Lys Val	Pro Ala Phe Ser Asp Lys	30
20	25	30	35
Asp Ser Leu Gly Asp	Glu Met Leu Ala	Ala Ala Leu Leu Lys Ala	45
35	40	45	50
Lys Ser Gln Glu Leu	Val Thr Phe Glu	Asp Val Ala Val Tyr Phe	60
50	55	60	65
Ile Arg Lys Glu Trp	Lys Arg Leu Glu	Pro Ala Gln Arg Asp Leu	75
65	70	75	80
Tyr Arg Asp Val Met	Leu Glu Asn Tyr	Gly Asn Val Phe Ser Leu	90
80	85	90	95
Asp Arg Glu Thr Arg	Thr Glu Asn Asp	Gln Glu Ile Ser Glu Asp	105
95	100	105	110
Thr Arg Ser His Gly	Val Leu Leu Gly	Arg Phe Gln Lys Asp Ile	120
110	115	120	125
Ser Gln Gly Leu Lys	Phe Lys Glu Ala	Tyr Glu Arg Glu Val Ser	135
125	130	135	140
Leu Lys Arg Pro Leu	Gly Asn Ser Pro	Gly Glu Arg Leu Asn Arg	150
140	145	150	155
Lys Met Pro Asp Phe	Gly Gln Val Thr	Val Glu Glu Lys Leu Thr	165
155	160	165	170
Pro Arg Gly Glu Arg	Ser Glu Lys Tyr	Asn Asp Phe Gly Asn Ser	180
170	175	180	185
Phe Thr Val Asn Ser	Asn Leu Ile Ser	His Gln Arg Leu Pro Val	195
185	190	195	200
Gly Asp Arg Pro His	Lys Cys Asp Glu	Cys Ser Lys Ser Phe Asn	210
200	205	210	215
Arg Thr Ser Asp Leu	Ile Gln His Gln	Ile His Thr Gly Glu	225
215	220	225	230
Lys Pro Tyr Glu Cys	Asn Glu Cys Gly	Lys Ala Phe Ser Gln Ser	240
230	235	240	245
Ser His Leu Ile Gln	His Gln Arg Ile	His Thr Gly Glu Lys Pro	255
245	250	255	260
Tyr Glu Cys Ser Asp	Cys Gly Lys Thr	Phe Ser Cys Ser Ser Ala	270
260	265	270	275
Leu Ile Leu His Arg	Arg Ile His Thr	Gly Glu Lys Pro Tyr Glu	285
275	280	285	290
Cys Asn Glu Cys Gly	Lys Thr Phe Ser	Trp Ser Ser Thr Leu Thr	300
290	295	300	305
His His Gln Arg Ile	His Thr Gly Glu	Lys Pro Tyr Ala Cys Asn	315
305	310	315	320
Glu Cys Gly Lys Ala	Phe Ser Arg Ser	Ser Thr Leu Ile His His	330
320	325	330	335
Gln Arg Ile His Thr	Gly Glu Lys Pro	Tyr Glu Cys Asn Glu Cys	345
335	340	345	350
Gly Lys Ala Phe Ser	Gln Ser Ser His	Leu Tyr Gln His Gln Arg	360
350	355	360	365
Ile His Thr Gly Glu	Lys Pro Tyr Glu	Cys Met Glu Cys Gly Gly	375
365	370	375	380
Lys Phe Thr Tyr Ser	Ser Gly Leu Ile	Gln His Gln Arg Ile His	390
380	385	390	395
Thr Gly Glu Asn Pro	Tyr Glu Cys Ser	Glu Cys Gly Lys Ala Phe	405
395	400	405	410
Arg Tyr Ser Ser Ala	Leu Val Arg His	Gln Arg Ile His Thr Gly	420
410	415	420	425
Glu Lys Pro Leu Asn	Gly Ile Gly Met	Ser Lys Ser Ser Leu Arg	

			425					430		435
Val	Thr	Thr	Glu	Leu	Asn	Ile	Arg	Glu	Ser	Thr
			440					445		

<210> 11  
 <211> 428  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID NO.: 1443262CD1

<400> 11  
 Met Glu Pro Leu Lys Val Glu Lys Phe Ala Thr Ala Asn Arg Gly  
 1 5 10 15  
 Asn Gly Leu Arg Ala Val Thr Pro Leu Arg Pro Gly Glu Leu Leu  
 20 25 30  
 Phe Arg Ser Asp Pro Leu Ala Tyr Thr Val Cys Lys Gly Ser Arg  
 35 40 45  
 Gly Val Val Cys Asp Arg Cys Leu Leu Gly Lys Glu Lys Leu Met  
 50 55 60  
 Arg Cys Ser Gln Cys Arg Val Ala Lys Tyr Cys Ser Ala Lys Cys  
 65 70 75  
 Gln Lys Lys Ala Trp Pro Asp His Lys Arg Glu Cys Lys Cys Leu  
 80 85 90  
 Lys Ser Cys Lys Pro Arg Tyr Pro Pro Asp Ser Val Arg Leu Leu  
 95 100 105  
 Gly Arg Val Val Phe Lys Leu Met Asp Gly Ala Pro Ser Glu Ser  
 110 115 120  
 Glu Lys Leu Tyr Ser Phe Tyr Asp Leu Glu Ser Asn Ile Asn Lys  
 125 130 135  
 Leu Thr Glu Asp Lys Lys Glu Gly Leu Arg Gln Leu Val Met Thr  
 140 145 150  
 Phe Gln His Phe Met Arg Glu Glu Ile Gln Asp Ala Ser Gln Leu  
 155 160 165  
 Pro Pro Ala Phe Asp Leu Phe Glu Ala Phe Ala Lys Val Ile Cys  
 170 175 180  
 Asn Ser Phe Thr Ile Cys Asn Ala Glu Met Gln Glu Val Gly Val  
 185 190 195  
 Gly Leu Tyr Pro Ser Ile Ser Leu Leu Asn His Ser Cys Asp Pro  
 200 205 210  
 Asn Cys Ser Ile Val Phe Asn Gly Pro His Leu Leu Leu Arg Ala  
 215 220 225  
 Val Arg Asp Ile Glu Val Gly Glu Glu Leu Thr Ile Cys Tyr Leu  
 230 235 240  
 Asp Met Leu Met Thr Ser Glu Glu Arg Arg Lys Gln Leu Arg Asp  
 245 250 255  
 Gln Tyr Cys Phe Glu Cys Asp Cys Phe Arg Cys Gln Thr Gln Asp  
 260 265 270  
 Lys Asp Ala Asp Met Leu Thr Gly Asp Glu Gln Val Trp Lys Glu  
 275 280 285  
 Val Gln Glu Ser Leu Lys Lys Ile Glu Glu Leu Lys Ala His Trp  
 290 295 300  
 Lys Trp Glu Gln Val Leu Ala Met Cys Gln Ala Ile Ile Ser Ser  
 305 310 315  
 Asn Ser Glu Arg Leu Pro Asp Ile Asn Ile Tyr Gln Leu Lys Val  
 320 325 330

WO 00/44900

PCT/US00/02237

Leu Asp Cys Ala Met Asp Ala Cys Ile Asn Leu Gly Leu Leu Glu  
 335 340 345  
 Glu Ala Leu Phe Tyr Gly Thr Arg Thr Met Glu Pro Tyr Arg Ile  
 350 355 360  
 Phe Phe Pro Gly Ser His Pro Val Arg Gly Val Gln Val Met Lys  
 365 370 375  
 Val Gly Lys Leu Gln Leu His Gln Gly Met Phe Pro Gln Ala Met  
 380 385 390  
 Lys Asn Leu Arg Leu Ala Phe Asp Ile Met Arg Val Thr His Gly  
 395 400 405  
 Arg Glu His Ser Leu Ile Glu Asp Leu Ile Leu Leu Leu Glu Glu  
 410 415 420  
 Cys Asp Ala Asn Ile Arg Ala Ser  
 425

&lt;210&gt; 12

&lt;211&gt; 590

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 1521648CD1

&lt;400&gt; 12

Met Ala Glu Asp Trp Leu Asp Cys Pro Ala Leu Gly Pro Gly Trp  
 1 5 10 15  
 Lys Arg Arg Glu Val Phe Arg Lys Ser Gly Ala Thr Cys Gly Arg  
 20 25 30  
 Ser Asp Thr Tyr Tyr Gln Ser Pro Thr Gly Asp Arg Ile Arg Ser  
 35 40 45  
 Lys Val Glu Leu Thr Arg Tyr Leu Gly Pro Ala Cys Asp Leu Thr  
 50 55 60  
 Leu Phe Asp Phe Lys Gln Gly Ile Leu Cys Tyr Pro Ala Pro Lys  
 65 70 75  
 Ala His Pro Val Ala Val Ala Ser Lys Lys Arg Lys Lys Pro Ser  
 80 85 90  
 Arg Pro Ala Lys Thr Arg Lys Arg Gln Val Gly Pro Gln Ser Gly  
 95 100 105  
 Glu Val Arg Lys Glu Ala Pro Arg Asp Glu Thr Lys Ala Asp Thr  
 110 115 120  
 Asp Thr Ala Pro Ala Ser Phe Pro Ala Pro Gly Cys Cys Glu Asn  
 125 130 135  
 Cys Gly Ile Ser Phe Ser Gly Asp Gly Thr Gln Arg Gln Arg Leu  
 140 145 150  
 Lys Thr Leu Cys Lys Asp Cys Arg Ala Gln Arg Ile Ala Phe Asn  
 155 160 165  
 Arg Glu Gln Arg Met Phe Lys Arg Val Gly Cys Gly Glu Cys Ala  
 170 175 180  
 Ala Cys Gln Val Thr Glu Asp Cys Gly Ala Cys Ser Thr Cys Leu  
 185 190 195  
 Leu Gln Leu Pro His Asp Val Ala Ser Gly Leu Phe Cys Lys Cys  
 200 205 210  
 Glu Arg Arg Arg Cys Leu Arg Ile Val Glu Arg Ser Arg Gly Cys  
 215 220 225  
 Gly Val Cys Arg Gly Cys Gln Thr Gln Glu Asp Cys Gly His Cys  
 230 235 240  
 Pro Ile Cys Leu Arg Pro Pro Arg Pro Gly Leu Arg Arg Gln Trp  
 245 250 255

WO 00/44900

PCT/US00/02237

Lys Cys Val Gln	Arg	Arg	Cys	Leu	Arg	Gly	Lys	His	Ala	Arg	Arg	
	260					265						270
Lys Gly Gly Cys	Asp	Ser	Lys	Met	Ala	Ala	Arg	Arg	Arg	Pro	Gly	
	275					280						285
Ala Gln Pro Leu	Pro	Pro	Pro	Pro	Pro	Ser	Gln	Ser	Pro	Glu	Pro	
	290					295						300
Thr Glu Pro His	Pro	Arg	Ala	Leu	Ala	Pro	Ser	Pro	Pro	Ala	Glu	
	305					310						315
Phe Ile Tyr Tyr	Cys	Val	Asp	Glu	Asp	Glu	Leu	Gln	Pro	Tyr	Thr	
	320					325						330
Asn Arg Arg Gln	Asn	Arg	Lys	Cys	Gly	Ala	Cys	Ala	Ala	Cys	Leu	
	335					340						345
Arg Arg Met Asp	Cys	Gly	Arg	Cys	Asp	Phe	Cys	Cys	Asp	Lys	Pro	
	350					355						360
Lys Phe Gly Gly	Ser	Asn	Gln	Lys	Arg	Gln	Lys	Cys	Arg	Trp	Arg	
	365					370						375
Gln Cys Leu Gln	Phe	Ala	Met	Lys	Arg	Leu	Leu	Pro	Ser	Val	Trp	
	380					385						390
Ser Glu Ser Glu	Asp	Gly	Ala	Gly	Ser	Pro	Pro	Pro	Tyr	Arg	Arg	
	395					400						405
Arg Lys Arg Pro	Ser	Ser	Ala	Arg	Arg	His	His	Leu	Gly	Pro	Thr	
	410					415						420
Leu Lys Pro Thr	Leu	Ala	Thr	Arg	Thr	Ala	Gln	Pro	Asp	His	Thr	
	425					430						435
Gln Ala Pro Thr	Lys	Gln	Glu	Ala	Gly	Gly	Gly	Phe	Val	Leu	Pro	
	440					445						450
Pro Pro Gly Thr	Asp	Leu	Val	Phe	Leu	Arg	Glu	Gly	Ala	Ser	Ser	
	455					460						465
Pro Val Gln Val	Pro	Gly	Pro	Val	Ala	Ala	Ser	Thr	Glu	Ala	Leu	
	470					475						480
Leu Gln Val Lys	Gln	Glu	Lys	Ala	Asp	Thr	Gln	Asp	Glu	Trp	Thr	
	485					490						495
Pro Gly Thr Ala	Val	Leu	Thr	Ser	Pro	Val	Leu	Val	Pro	Gly	Cys	
	500					505						510
Pro Ser Lys Ala	Val	Asp	Pro	Gly	Leu	Pro	Ser	Val	Lys	Gln	Glu	
	515					520						525
Pro Pro Asp Pro	Glu	Glu	Asp	Lys	Glu	Glu	Asn	Lys	Asp	Asp	Ser	
	530					535						540
Ala Ser Lys Leu	Ala	Pro	Glu	Glu	Glu	Ala	Gly	Gly	Ala	Gly	Thr	
	545					550						555
Pro Val Ile Thr	Glu	Ile	Phe	Ser	Leu	Gly	Gly	Thr	Arg	Phe	Arg	
	560					565						570
Asp Thr Ala Val	Trp	Leu	Pro	Arg	Ser	Lys	Asp	Leu	Lys	Lys	Pro	
	575					580						585
Gly Ala Arg Lys	Gln											
	590											

<210> 13  
 <211> 479  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 1685494CD1

<400> 13  
 Met Ala Thr Ala Leu Val Ser Ala His Ser Leu Ala Pro Leu Ser

1	5	10	15
Leu Lys Lys Glu Gly	Leu Arg Val Val Arg	Glu Asp His Tyr Ser	
20	25	30	
Thr Trp Glu Gln Gly	Phe Lys Leu Gln Gly	Asn Ser Lys Gly Leu	
35	40	45	
Gly Gln Glu Pro Leu	Cys Lys Gln Phe Arg	Gln Leu Arg Tyr Glu	
50	55	60	
Glu Thr Thr Gly Pro	Arg Glu Ala Leu Ser	Arg Leu Arg Glu Leu	
65	70	75	
Cys Gln Gln Trp Leu	Gln Pro Glu Thr His	Thr Lys Glu Gln Ile	
80	85	90	
Leu Glu Leu Leu Val	Leu Glu Gln Phe Leu	Ile Ile Leu Pro Lys	
95	100	105	
Glu Leu Gln Ala Arg	Val Gln Glu His His	Pro Glu Ser Arg Glu	
110	115	120	
Asp Val Val Val Val	Leu Glu Asp Leu Gln	Leu Asp Leu Gly Val	
125	130	135	
Thr Gly Gln Gln Val	Asp Pro Asp Gln Pro	Lys Lys Gln Lys Ile	
140	145	150	
Leu Val Glu Glu Met	Ala Pro Leu Lys Gly	Val Gln Glu Gln Gln	
155	160	165	
Val Arg His Glu Cys	Glu Val Thr Lys Pro	Glu Lys Glu Lys Gly	
170	175	180	
Glu Glu Thr Arg Ile	Glu Asn Gly Lys Leu	Ile Val Val Thr Asp	
185	190	195	
Ser Cys Gly Arg Val	Glu Ser Ser Gly Lys	Ile Ser Glu Pro Met	
200	205	210	
Glu Ala His Asn Glu	Gly Ser Asn Leu Glu	Arg His Gln Ala Lys	
215	220	225	
Pro Lys Glu Lys Ile	Glu Tyr Lys Cys Ser	Glu Arg Glu Gln Arg	
230	235	240	
Phe Ile Gln His Leu	Asp Leu Ile Glu His	Ala Ser Thr His Thr	
245	250	255	
Gly Lys Lys Leu Cys	Glu Ser Asp Val Cys	Gln Ser Ser Ser Leu	
260	265	270	
Thr Gly His Lys Lys	Val Leu Ser Arg Glu	Lys Gly His Gln Cys	
275	280	285	
His Glu Cys Gly Lys	Ala Phe Gln Arg Ser	Ser His Leu Val Arg	
290	295	300	
His Gln Lys Ile His	Leu Glu Gly Glu Lys	Pro Tyr Gln Cys Asn Glu	
305	310	315	
Cys Gly Lys Val Phe	Ser Ser Gln Asn Ala Gly	Leu Leu Glu His Leu	
320	325	330	
Arg Ile His Thr Gly	Glu Lys Pro Tyr Leu	Cys Ile His Cys Gly	
335	340	345	
Lys Asn Phe Arg Arg	Ser Ser His Leu Asn	Arg His Gln Arg Ile	
350	355	360	
His Ser Gln Glu Glu	Pro Cys Glu Cys Lys	Glu Cys Gly Lys Thr	
365	370	375	
Phe Ser Gln Ala Leu	Leu Leu Thr His His	Gln Arg Ile His Ser	
380	385	390	
His Ser Lys Ser His	Gln Cys Asn Glu Cys	Gly Lys Ala Phe Ser	
395	400	405	
Leu Thr Ser Asp Leu	Ile Arg His His Arg	Ile His Thr Gly Glu	
410	415	420	
Lys Pro Phe Lys Cys	Asn Ile Cys Gln Lys	Ala Phe Arg Leu Asn	
425	430	435	
Ser His Leu Ala Gln	His Val Arg Ile His	Asn Glu Glu Lys Pro	
440	445	450	
Tyr Gln Cys Ser Glu	Cys Gly Glu Ala Phe	Arg Gln Arg Ser Gly	
455	460	465	
Leu Phe Gln His Gln	Arg Tyr His His Lys	Asp Lys Leu Ala	

470

475

<210> 14  
 <211> 433  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 1730829CD1

<400> 14  
 Met Glu Ala Val Tyr Leu Val Val Asn Gly Leu Gly Leu Val Leu  
 1 5 10 15  
 Asp Val Leu Thr Leu Val Leu Asp Leu Asn Phe Leu Leu Val Ser  
 20 25 30  
 Ser Leu Leu Ala Ser Leu Ala Trp Leu Leu Ala Phe Val Tyr Asn  
 35 40 45  
 Leu Pro His Thr Val Leu Thr Ser Leu Leu His Leu Gly Arg Gly  
 50 55 60  
 Val Leu Leu Ser Leu Leu Ala Leu Ile Glu Ala Val Val Arg Phe  
 65 70 75  
 Thr Cys Gly Gly Leu Gln Ala Leu Cys Thr Leu Leu Tyr Ser Cys  
 80 85 90  
 Cys Ser Gly Leu Glu Ser Leu Lys Leu Leu Gly His Leu Ala Ser  
 95 100 105  
 His Gly Ala Leu Arg Ser Arg Glu Ile Leu His Arg Gly Val Leu  
 110 115 120  
 Asn Val Val Ser Ser Gly His Ala Leu Leu Arg Gln Ala Cys Asp  
 125 130 135  
 Ile Cys Ala Ile Ala Met Ser Leu Val Ala Tyr Val Ile Asn Ser  
 140 145 150  
 Leu Val Asn Ile Cys Leu Ile Gly Thr Gln Asn Leu Phe Ser Leu  
 155 160 165  
 Val Leu Ala Leu Trp Asp Ala Val Thr Gly Pro Leu Trp Arg Met  
 170 175 180  
 Thr Asp Val Val Ala Ala Phe Leu Ala His Ile Ser Ser Ser Ala  
 185 190 195  
 Val Ala Met Ala Ile Leu Leu Trp Thr Pro Cys Gln Leu Ala Leu  
 200 205 210  
 Glu Leu Leu Ala Ser Ala Ala Arg Leu Leu Ala Ser Phe Val Leu  
 215 220 225  
 Val Asn Leu Thr Gly Leu Val Leu Leu Ala Cys Val Leu Ala Val  
 230 235 240  
 Thr Val Thr Val Leu His Pro Asp Phe Thr Leu Arg Leu Ala Thr  
 245 250 255  
 Gln Ala Leu Ser Gln Leu His Ala Arg Pro Ser Tyr His Arg Leu  
 260 265 270  
 Arg Glu Asp Val Met Arg Leu Ser Arg Leu Ala Leu Gly Ser Glu  
 275 280 285  
 Ala Trp Arg Arg Val Trp Ser Arg Ser Leu Gln Leu Ala Ser Trp  
 290 295 300  
 Pro Asn Arg Gly Gly Ala Pro Gly Ala Pro Gln Gly Asp Pro Met  
 305 310 315  
 Arg Val Phe Ser Val Arg Thr Arg Arg Gln Asp Thr Leu Pro Glu  
 320 325 330  
 Ala Gly Arg Arg Ser Glu Ala Glu Glu Glu Ala Arg Thr Ile  
 335 340 345  
 Arg Val Thr Pro Val Arg Gly Arg Glu Arg Leu Asn Glu Glu Glu

18/91



	275		280		285
Glu Gly Arg Lys	Asp Ser Asp Thr Glu	Ser Ser Asp Leu Phe	Thr		
	290		295		300
Asn Leu Asn Leu	Gly Arg Thr Tyr Ala	Ser Gly Tyr Ala His	Tyr		
	305		310		315
Glu Glu Gln Glu	Asn				
	320				

<210> 16  
 <211> 179  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 2444604CD1

<400> 16  
 Met Ala Ala Gly Phe Phe Gln Pro Phe Met Ser Pro Arg Phe Pro  
 1 5 10 15  
 Gly Gly Pro Arg Pro Thr Leu Arg Met Pro Ser Gln Pro Pro Ala  
 20 25 30  
 Cys Leu Pro Gly Ser Gln Pro Leu Leu Pro Gly Ala Met Glu Pro  
 35 40 45  
 Ser Pro Arg Ala Gln Gly His Pro Ser Met Gly Gly Pro Met Gln  
 50 55 60  
 Arg Val Thr Pro Pro Arg Gly Met Ala Ser Val Gly Pro Gln Ser  
 65 70 75  
 Tyr Gly Gly Gly Met Arg Pro Pro Asn Ser Leu Ala Gly Pro  
 80 85 90  
 Gly Leu Pro Ala Met Asn Met Gly Pro Gly Val Arg Gly Pro Trp  
 95 100 105  
 Ala Ser Pro Ser Gly Asn Ser Ile Pro Tyr Ser Ser Ser Ser Pro  
 110 115 120  
 Gly Ser Tyr Thr Gly Pro Pro Gly Gly Gly Gly Pro Pro Gly Thr  
 125 130 135  
 Pro Ile Met Pro Ser Pro Gly Asp Ser Thr Asn Ser Ser Glu Asn  
 140 145 150  
 Met Tyr Thr Ile Met Asn Pro Ile Gly Gln Ala Gly Arg Ala  
 155 160 165  
 Asn Phe Pro Leu Gly Pro Gly Pro Glu Gly Pro Trp Pro Pro  
 170 175

<210> 17  
 <211> 494  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 2445008CD1

<400> 17  
 Met Gly Arg Lys Lys Lys Lys Gln Leu Lys Pro Trp Cys Trp Tyr  
 1 5 10 15  
 Cys Asn Arg Asp Phe Asp Asp Glu Lys Ile Leu Ile Gln His Gln



				20						25				30
Lys	Ala	Lys	His	Phe	Lys	Cys	His	Ile	Cys	His	Lys	Lys	Leu	Tyr
				35					40					45
Thr	Gly	Pro	Gly	Leu	Ala	Ile	His	Cys	Met	Gln	Val	His	Lys	Glu
				50					55					60
Thr	Ile	Asp	Ala	Val	Pro	Asn	Ala	Ile	Pro	Gly	Arg	Thr	Asp	Ile
				65					70					75
Glu	Leu	Glu	Ile	Tyr	Gly	Met	Glu	Gly	Ile	Pro	Glu	Lys	Asp	Met
				80					85					90
Asp	Glu	Arg	Arg	Arg	Leu	Leu	Glu	Gln	Lys	Thr	Gln	Glu	Ser	Gln
				95					100					105
Lys	Lys	Lys	Gln	Gln	Asp	Asp	Ser	Asp	Glu	Tyr	Asp	Asp	Asp	Asp
				110					115					120
Ser	Ala	Ala	Ser	Thr	Ser	Phe	Gln	Pro	Gln	Pro	Val	Gln	Pro	Gln
				125					130					135
Gln	Gly	Tyr	Ile	Pro	Pro	Met	Ala	Gln	Pro	Gly	Leu	Pro	Pro	Val
				140					145					150
Pro	Gly	Ala	Pro	Gly	Met	Pro	Pro	Gly	Ile	Pro	Pro	Leu	Met	Pro
				155					160					165
Gly	Val	Pro	Pro	Leu	Met	Pro	Gly	Met	Pro	Pro	Val	Met	Pro	Gly
				170					175					180
Met	Pro	Pro	Gly	Leu	His	His	Gln	Arg	Lys	Tyr	Thr	Gln	Ser	Phe
				185					190					195
Cys	Gly	Glu	Asn	Ile	Met	Met	Pro	Met	Gly	Gly	Met	Met	Pro	Pro
				200					205					210
Gly	Pro	Gly	Ile	Pro	Pro	Leu	Met	Pro	Gly	Met	Pro	Pro	Gly	Met
				215					220					225
Pro	Pro	Pro	Val	Pro	Arg	Pro	Gly	Ile	Pro	Pro	Met	Thr	Gln	Ala
				230					235					240
Gln	Ala	Val	Ser	Ala	Pro	Gly	Ile	Leu	Asn	Arg	Pro	Pro	Ala	Pro
				245					250					255
Thr	Ala	Thr	Val	Pro	Ala	Pro	Gln	Pro	Pro	Val	Thr	Lys	Pro	Leu
				260					265					270
Phe	Pro	Ser	Ala	Gly	Gln	Met	Gly	Thr	Pro	Val	Thr	Ser	Ser	Ser
				275					280					285
Thr	Ala	Ser	Ser	Asn	Ser	Glu	Ser	Leu	Ser	Ala	Ser	Ser	Lys	Ala
				290					295					300
Leu	Phe	Pro	Ser	Thr	Ala	Gln	Ala	Gln	Ala	Ala	Val	Gln	Gly	Pro
				305					310					315
Val	Gly	Thr	Asp	Phe	Lys	Pro	Leu	Asn	Ser	Thr	Pro	Ala	Thr	Thr
				320					325					330
Thr	Glu	Pro	Pro	Lys	Pro	Thr	Phe	Pro	Ala	Tyr	Thr	Gln	Ser	Thr
				335					340					345
Ala	Ser	Thr	Thr	Ser	Thr	Thr	Asn	Ser	Thr	Ala	Ala	Lys	Pro	Ala
				350					355					360
Ala	Ser	Ile	Thr	Ser	Lys	Pro	Ala	Thr	Leu	Thr	Thr	Thr	Ser	Ala
				365					370					375
Thr	Ser	Lys	Leu	Ile	His	Pro	Asp	Glu	Asp	Ile	Ser	Leu	Glu	Glu
				380					385					390
Arg	Arg	Ala	Gln	Leu	Pro	Lys	Tyr	Gln	Arg	Asn	Leu	Pro	Arg	Pro
				395					400					405
Gly	Gln	Ala	Pro	Ile	Gly	Asn	Pro	Pro	Val	Gly	Pro	Ile	Gly	Gly
				410					415					420
Met	Met	Pro	Pro	Gln	Pro	Gly	Ile	Pro	Gln	Gln	Gln	Gly	Met	Arg
				425					430					435
Pro	Pro	Met	Pro	Pro	His	Gly	Gln	Tyr	Gly	Gly	His	His	Gln	Gly
				440					445					450
Met	Pro	Gly	Tyr	Leu	Pro	Gly	Ala	Met	Pro	Pro	Tyr	Gly	Gln	Gly
				455					460					465
Pro	Pro	Met	Val	Pro	Pro	Tyr	Gln	Gly	Gly	Pro	Pro	Arg	Pro	Pro
				470					475					480

Met Gly Met Arg Pro Pro Val Met Ser Gln Gly Gly Arg Tyr  
485 490

<210> 18  
<211> 401  
<212> PRT  
<213> Homo sapiens

<220>  
<221> misc-feature  
<223> Incyte ID No.: 2572462CD1

<400> 18  
Met Ala Ser Ser Pro Arg Pro Lys Met Asp Ala Ile Leu Thr Glu  
1 5 10 15  
Ala Ile Lys Ala Cys Phe Gln Lys Ser Gly Ala Ser Val Val Ala  
20 25 30  
Ile Arg Lys Tyr Ile Ile His Lys Tyr Pro Ser Leu Glu Leu Glu  
35 40 45  
Arg Arg Gly Tyr Leu Leu Lys Gln Ala Leu Lys Arg Glu Leu Asn  
50 55 60  
Arg Gly Val Ile Lys Gln Val Lys Gly Lys Gly Ala Ser Gly Ser  
65 70 75  
Phe Val Val Val Gln Lys Ser Arg Lys Thr Pro Gln Lys Ser Arg  
80 85 90  
Asn Arg Lys Asn Arg Ser Ser Ala Val Asp Pro Glu Pro Gln Val  
95 100 105  
Lys Leu Glu Asp Val Leu Pro Leu Ala Phe Thr Arg Leu Cys Glu  
110 115 120  
Pro Lys Glu Ala Ser Tyr Ser Leu Ile Arg Lys Tyr Val Ser Gln  
125 130 135  
Tyr Tyr Pro Lys Leu Arg Val Asp Ile Arg Pro Gln Leu Leu Lys  
140 145 150  
Asn Ala Leu Gln Arg Ala Val Glu Arg Gly Gln Leu Glu Gln Ile  
155 160 165  
Thr Gly Lys Gly Ala Ser Gly Thr Phe Gln Leu Lys Lys Ser Gly  
170 175 180  
Glu Lys Pro Leu Leu Gly Gly Ser Leu Met Glu Tyr Ala Ile Leu  
185 190 195  
Ser Ala Ile Ala Ala Met Asn Glu Pro Lys Thr Cys Ser Thr Thr  
200 205 210  
Ala Leu Lys Lys Tyr Val Leu Glu Asn His Pro Gly Thr Asn Ser  
215 220 225  
Asn Tyr Gln Met His Leu Leu Lys Lys Thr Leu Gln Lys Cys Glu  
230 235 240  
Lys Asn Gly Trp Met Glu Gln Ile Ser Gly Lys Gly Phe Ser Gly  
245 250 255  
Thr Phe Gln Leu Cys Phe Pro Tyr Tyr Pro Ser Pro Gly Val Leu  
260 265 270  
Phe Pro Lys Lys Glu Pro Asp Asp Ser Arg Asp Glu Asp Glu Asp  
275 280 285  
Glu Asp Glu Ser Ser Glu Glu Asp Ser Glu Asp Glu Glu Pro Pro  
290 295 300  
Pro Lys Arg Arg Leu Gln Lys Lys Thr Pro Ala Lys Ser Pro Gly  
305 310 315  
Lys Ala Ala Ser Val Lys Gln Arg Gly Ser Lys Pro Ala Pro Lys  
320 325 330  
Val Ser Ala Ala Gln Arg Gly Lys Ala Arg Pro Leu Pro Lys Lys  
335 340 345

Ala	Pro	Pro	Lys	Ala	Lys	Thr	Pro	Ala	Lys	Lys	Thr	Arg	Pro	Ser	
			350						355					360	
Ser	Thr	Val	Ile	Lys	Lys	Pro	Ser	Gly	Gly	Ser	Ser	Lys	Lys	Pro	
			365						370					375	
Ala	Thr	Ser	Ala	Arg	Lys	Glu	Val	Lys	Leu	Pro	Gly	Lys	Gly	Lys	
			380						385					390	
Ser	Thr	Met	Lys	Lys	Ser	Phe	Arg	Val	Lys	Lys					
			395						400						

&lt;210&gt; 19

&lt;211&gt; 264

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 2572892CD1

&lt;400&gt; 19

Met	Pro	Arg	Ser	Phe	Leu	Val	Arg	Lys	Pro	Ser	Asp	Pro	Asn	Arg	
1				5					10					15	
Lys	Pro	Asn	Tyr	Ser	Glu	Leu	Gln	Asp	Ser	Asn	Pro	Glu	Phe	Thr	
			20						25					30	
Phe	Gln	Gln	Pro	Tyr	Asp	Gln	Ala	His	Leu	Leu	Ala	Ala	Ile	Pro	
			35						40					45	
Pro	Pro	Glu	Ile	Leu	Asn	Pro	Thr	Ala	Ser	Leu	Pro	Met	Leu	Ile	
			50						55					60	
Trp	Asp	Ser	Val	Leu	Ala	Pro	Gln	Ala	Gln	Pro	Ile	Ala	Trp	Ala	
			65						70					75	
Ser	Leu	Arg	Leu	Gln	Glu	Ser	Pro	Arg	Val	Ala	Glu	Leu	Thr	Ser	
			80						85					90	
Leu	Ser	Asp	Glu	Asp	Ser	Gly	Lys	Gly	Ser	Gln	Pro	Pro	Ser	Pro	
			95						100					105	
Pro	Ser	Pro	Ala	Pro	Ser	Ser	Phe	Ser	Ser	Thr	Ser	Ala	Ser	Ser	
			110						115					120	
Leu	Glu	Ala	Glu	Ala	Tyr	Ala	Ala	Phe	Pro	Gly	Leu	Gly	Gln	Val	
			125						130					135	
Pro	Lys	Gln	Leu	Ala	Gln	Leu	Ser	Glu	Ala	Lys	Asp	Leu	Gln	Ala	
			140						145					150	
Arg	Lys	Ala	Phe	Asn	Cys	Lys	Tyr	Cys	Asn	Lys	Glu	Tyr	Leu	Ser	
			155						160					165	
Leu	Gly	Ala	Leu	Lys	Met	His	Ile	Arg	Ser	His	Thr	Leu	Pro	Cys	
			170						175					180	
Val	Cys	Gly	Thr	Cys	Gly	Lys	Ala	Phe	Ser	Arg	Pro	Trp	Leu	Leu	
			185						190					195	
Gln	Gly	His	Val	Arg	Thr	His	Thr	Gly	Glu	Lys	Pro	Phe	Ser	Cys	
			200						205					210	
Pro	His	Cys	Ser	Arg	Ala	Phe	Ala	Asp	Arg	Ser	Asn	Leu	Arg	Ala	
			215						220					225	
His	Leu	Gln	Thr	His	Ser	Asp	Val	Lys	Lys	Tyr	Gln	Cys	Gln	Ala	
			230						235					240	
Cys	Ala	Arg	Thr	Phe	Ser	Arg	Met	Ser	Leu	Leu	His	Lys	His	Gln	
			245						250					255	
Glu	Ser	Gly	Cys	Ser	Gly	Cys	Pro	Arg							
			260												

<210> 20  
 <211> 153  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <221> misc-feature  
 <223> Incyte ID No.: 2785674CD1

<400> 20  
 Met Thr Lys Ile Lys Ala Asp Pro Asp Gly Pro Glu Ala Gln Ala  
 1 5 10 15  
 Glu Ala Cys Ser Gly Glu Arg Thr Tyr Gln Glu Leu Leu Val Asn  
 20 25 30  
 Gln Asn Pro Ile Ala Gln Pro Leu Ala Ser Arg Arg Leu Thr Arg  
 35 40 45  
 Lys Leu Tyr Lys Cys Ile Lys Lys Ala Val Lys Gln Lys Gln Ile  
 50 55 60  
 Arg Arg Gly Val Lys Glu Val Gln Lys Phe Val Asn Lys Gly Glu  
 65 70 75  
 Lys Gly Ile Met Val Leu Ala Gly Asp Thr Leu Pro Ile Glu Val  
 80 85 90  
 Tyr Cys His Leu Pro Val Met Cys Glu Asp Arg Asn Leu Pro Tyr  
 95 100 105  
 Val Tyr Ile Pro Ser Lys Thr Asp Leu Gly Ala Ala Ala Gly Ser  
 110 115 120  
 Lys Arg Pro Thr Cys Val Ile Met Val Lys Pro His Glu Glu Tyr  
 125 130 135  
 Gln Glu Ala Tyr Asp Glu Cys Leu Glu Glu Val Gln Ser Leu Pro  
 140 145 150  
 Leu Pro Leu

<210> 21  
 <211> 243  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <221> misc-feature  
 <223> Incyte ID No.: 2797479CD1

<400> 21  
 Met Gly Asp Asp Ile Ser Thr His Ile Ala Pro Lys Glu Leu Arg  
 1 5 10 15  
 His Lys His Pro Ser Ser Val Asp Glu Val Ala Gln Val Val Lys  
 20 25 30  
 Gln Leu Arg Ile Ile Leu Gln Gln Gln Val Arg Pro Gly Glu Ser  
 35 40 45  
 Thr Val Leu Ala Leu Arg Pro Asn Val Gln Gln Ile Glu Ala Pro  
 50 55 60  
 Asp Val Ser Arg Asp Pro Arg Val Leu Gly Phe Asp Phe Pro Gly  
 65 70 75  
 Ser Ala Arg Ile His Glu Gly Thr His Thr Leu Glu Lys Pro Tyr  
 80 85 90  
 Glu Cys Lys Gln Cys Gly Lys Leu Leu Ser His Arg Ser Ser Phe  
 95 100 105  
 Arg Arg His Met Met Ala His Thr Gly Asp Gly Pro His Lys Cys  
 110 115 120  
 Thr Val Cys Gly Lys Ala Phe Asp Ser Pro Ser Val Phe Gln Arg

	125		130		135
His Glu Arg Thr	His Thr Gly Glu Lys	Pro Tyr Glu Cys Lys	Gln		
	140		145		150
Cys Gly Lys Ala	Phe Arg Thr Ser Ser	Ser Leu Arg Lys His	Glu		
	155		160		165
Thr Thr His Thr	Gly Glu Gln Pro Tyr	Lys Cys Lys Cys Gly Lys			
	170		175		180
Ala Phe Ser Asp	Leu Phe Ser Phe Gln Ser	His Glu Thr Thr His			
	185		190		195
Ser Glu Glu Glu	Pro Tyr Glu Cys Lys	Glu Cys Gly Lys Ala Phe			
	200		205		210
Ser Ser Phe Lys	Tyr Phe Cys Arg His	Glu Arg Thr His Ser Glu			
	215		220		225
Glu Lys Ser Tyr	Glu Cys Gln Ile Cys	Gly Lys Leu Ser Val Val			
	230		235		240
Ser Val Thr					

&lt;210&gt; 22

&lt;211&gt; 485

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 2960640CD1

&lt;400&gt; 22

Met Arg Asp Asn Arg	Ala Val Ser Leu Cys	Gln Gln Glu Trp Met	
1	5	10	15
Cys Pro Gly Pro Ala	Gln Arg Ala Leu Tyr	Arg Gly Ala Thr Gln	
	20	25	30
Arg Lys Asp Ser His	Val Ser Leu Ala Thr	Gly Val Pro Trp Gly	
	35	40	45
Tyr Glu Glu Thr Lys	Thr Leu Leu Ala Ile	Leu Ser Ser Ser Gln	
	50	55	60
Phe Tyr Gly Lys Leu	Gln Thr Cys Gln Asn	Ser Gln Ile Tyr	
	65	70	75
Arg Ala Met Ala Glu	Gly Leu Trp Glu Gln	Gly Phe Leu Arg Thr	
	80	85	90
Pro Glu Gln Cys Arg	Thr Lys Phe Lys Ser	Leu Gln Leu Ser Tyr	
	95	100	105
Arg Lys Val Arg Arg	Gly Arg Val Pro Glu	Pro Cys Ile Phe Tyr	
	110	115	120
Glu Glu Met Asn Ala	Leu Ser Gly Ser Trp	Ala Ser Ala Pro Pro	
	125	130	135
Met Ala Ser Asp Ala	Val Pro Gly Gln Glu	Gly Ser Asp Ile Glu	
	140	145	150
Ala Gly Glu Leu Asn	His Gln Asn Gly Glu	Pro Thr Glu Val Glu	
	155	160	165
Asp Gly Thr Val Asp	Gly Ala Asp Arg Asp	Glu Lys Asp Phe Arg	
	170	175	180
Asn Pro Gly Gln Glu	Val Arg Lys Leu Asp	Leu Pro Val Leu Phe	
	185	190	195
Pro Asn Arg Leu Gly	Phe Glu Phe Lys Asn	Glu Ile Lys Lys Glu	
	200	205	210
Asn Leu Lys Trp Asp	Asp Ser Glu Glu Val	Glu Ile Asn Lys Ala	
	215	220	225
Leu Gln Arg Lys Ser	Arg Gly Val Tyr Trp	His Ser Glu Leu Gln	

Lys Gly Leu Glu	230	235	240
Ser Glu Pro Thr Ser	245	250	255
Ser Pro Gly Glu Ser	260	265	270
Ser His Gln Ser Phe	275	280	285
Leu Cys Gly Lys Asn	290	295	300
Pro Ala Leu Lys Leu	305	310	315
Lys Thr Phe Ser Arg	320	325	330
His Thr Gly Glu Lys	335	340	345
Phe Ser Glu Arg Ser	350	355	360
Gly Glu Arg Pro Tyr	365	370	375
Gln Ser Ser Ser Leu	380	385	390
Lys Pro Tyr Gln Cys	395	400	405
Ser Gln Phe Ser Ala	410	415	420
Tyr Lys Cys Ala Val	425	430	435
Phe Ser Ala His Arg	440	445	450
Cys Ser His Cys Glu	455	460	465
Arg His Gln Thr Val	470	475	480
Gly Arg Asp Ala Leu	485		

&lt;210&gt; 23

&lt;211&gt; 160

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 3454051CD1

&lt;400&gt; 23

Met Ser Trp Thr Cys	Pro Arg Cys Gln Gln	Pro Val Phe Phe Ala
1	5	10
Glu Lys Val Ser Ser	Leu Gly Lys Asn Trp	His Arg Phe Cys Leu
20	25	30
Lys Cys Glu Arg Cys	His Ser Ile Leu Ser	Pro Gly Gly His Ala
35	40	45
Glu His Asn Gly Arg	Pro Tyr Cys His Lys	Pro Cys Tyr Gly Ala
50	55	60
Leu Phe Gly Pro Arg	Gly Pro Pro His Met	Lys Thr Phe Thr Gly
65	70	75
Glu Thr Ser Leu Cys	Pro Gly Cys Gly Glu	Pro Val Tyr Phe Ala
80	85	90

Glu	Lys	Val	Met	Ser	Leu	Gly	Arg	Asn	Trp	His	Arg	Pro	Cys	Leu	
				95					100					105	
Arg	Cys	Gln	Arg	Cys	His	Lys	Thr	Leu	Thr	Ala	Gly	Ser	His	Ala	
				110					115					120	
Glu	His	Asp	Gly	Val	Pro	Tyr	Cys	His	Val	Pro	Cys	Tyr	Gly	Tyr	
				125					130					135	
Leu	Phe	Gly	Pro	Lys	Gly	Val	Asn	Ile	Gly	Asp	Val	Gly	Cys	Tyr	
				140					145					150	
Ile	Tyr	Asp	Pro	Val	Lys	Ile	Lys	Phe	Lys						
				155					160						

&lt;210&gt; 24

&lt;211&gt; 511

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt;

&lt;223&gt; Incyte ID No.: 3510640CD1

&lt;400&gt; 24

Met	Gln	Glu	Leu	Tyr	Ser	Thr	Pro	Ala	Ser	Arg	Leu	Asp	Ser	Phe	
1				5					10					15	
Val	Ala	Gln	Trp	Leu	Gln	Pro	His	Arg	Glu	Trp	Lys	Glu	Glu	Val	
				20					25					30	
Leu	Asp	Ala	Val	Arg	Thr	Val	Glu	Glu	Phe	Leu	Arg	Gln	Glu	His	
				35					40					45	
Phe	Gln	Gly	Lys	Arg	Gly	Leu	Asp	Gln	Asp	Val	Arg	Val	Leu	Lys	
				50					55					60	
Val	Val	Lys	Val	Gly	Ser	Phe	Gly	Asn	Gly	Thr	Val	Leu	Arg	Ser	
				65					70					75	
Thr	Arg	Glu	Val	Glu	Leu	Val	Ala	Phe	Leu	Ser	Cys	Phe	His	Ser	
				80					85					90	
Phe	Gln	Glu	Ala	Ala	Lys	His	His	Lys	Asp	Val	Leu	Arg	Leu	Ile	
				95					100					105	
Trp	Lys	Thr	Met	Trp	Gln	Ser	Gln	Asp	Leu	Leu	Asp	Leu	Gly	Leu	
				110					115					120	
Glu	Asp	Leu	Arg	Met	Glu	Gln	Arg	Val	Pro	Asp	Ala	Leu	Val	Phe	
				125					130					135	
Thr	Ile	Gln	Thr	Arg	Gly	Thr	Ala	Glu	Pro	Ile	Thr	Val	Thr	Ile	
				140					145					150	
Val	Pro	Ala	Tyr	Arg	Ala	Leu	Gly	Pro	Ser	Leu	Pro	Asn	Ser	Gln	
				155					160					165	
Pro	Pro	Pro	Glu	Val	Tyr	Val	Ser	Leu	Ile	Lys	Ala	Cys	Gly	Gly	
				170					175					180	
Pro	Gly	Asn	Phe	Cys	Pro	Phe	Phe	Ser	Glu	Leu	Gln	Arg	Asn	Phe	
				185					190					195	
Val	Lys	His	Arg	Pro	Thr	Lys	Leu	Lys	Ser	Leu	Leu	Arg	Leu	Val	
				200					205					210	
Lys	His	Trp	Tyr	Gln	Gln	Tyr	Val	Lys	Ala	Arg	Ser	Pro	Arg	Ala	
				215					220					225	
Asn	Leu	Pro	Pro	Leu	Tyr	Ala	Leu	Glu	Leu	Leu	Thr	Ile	Tyr	Ala	
				230					235					240	
Trp	Glu	Met	Gly	Thr	Glu	Glu	Asp	Glu	Asn	Phe	Met	Leu	Asp	Glu	
				245					250					255	
Gly	Phe	Thr	Thr	Val	Met	Asp	Leu	Leu	Leu	Glu	Tyr	Glu	Val	Ile	
				260					265					270	
Cys	Ile	Tyr	Trp	Thr	Lys	Tyr	Tyr	Thr	Leu	His	Asn	Ala	Ile	Ile	

Glu Asp Cys Val	275	Lys Gln Leu Lys	280	Lys Glu Arg Pro Ile	285
Leu Asp Pro Ala	290	Pro Thr Leu Asn	295	Val Ala Glu Gly Tyr	300
Trp Asp Ile Val	305	Gln Arg Ala Ser	310	Gln Cys Leu Lys Gln	315
Cys Cys Tyr Asp	320	Asn Arg Glu Asn Pro	325	Ile Ser Ser Trp Asn	330
Lys Arg Ala Arg	335	Asp Ile His Leu Thr	340	Val Glu Gln Arg Gly	345
Pro Asp Phe Asn	350	Leu Ile Val Asn Pro	355	Tyr Glu Pro Ile Arg	360
Val Lys Glu Lys	365	Ile Arg Arg Thr Arg	370	Gly Tyr Ser Gly Leu	375
Arg Leu Ser Phe	380	Gln Val Pro Gly Ser	385	Glu Arg Gln Leu Leu	390
Ser Arg Cys ser	395	Leu Ala Lys Tyr Gly	400	Ile Phe Ser His Thr	405
Ile Tyr Leu Leu	410	Glu Thr Ile Pro Ser	415	Glu Ile Gln Val Phe	420
Lys Asn Pro Asp	425	Gly Gly Ser Tyr Ala	430	Tyr Ala Ile Asn Pro	435
Ser Phe Ile Leu	440	Gly Leu Lys Gln Gln	445	Ile Glu Asp Gln Gln	450
	455		460		465
Leu Pro Lys Lys	470	Gln Gln Gln Leu Glu	475	Phe Gln Gly Gln Val	480
Gln Asp Trp Leu	485	Gly Leu Gly Ile Tyr	490	Gly Ile Gln Asp Ser	495
Thr Leu Ile Leu	500	Ser Lys Lys Lys Gly	505	Glu Ala Leu Phe Pro	510
Ser					

&lt;210&gt; 25

&lt;211&gt; 310

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt; -

&lt;223&gt; Incyte ID No.: 3815083CD1

&lt;400&gt; 25

Met Arg Pro Leu	1	Gln Ile Val Pro	5	Ser Arg Leu Ile Ser	9	Gln Leu	13
Tyr Cys Gly Leu	1	Lys Pro Pro Ala	5	Ser Thr Arg Asn Gln	9	Ile Cys	13
Leu Lys Met Ala	20	Arg Pro Ser Ser	20	Ser Met Ala Asp Phe	24	Arg Lys	28
Phe Phe Ala Lys	35	Ala Lys His Ile	35	Val Ile Ile Ser Gly	39	Ala Gly	43
Val Ser Ala Glu	50	Ser Gly Val Pro	50	Thr Phe Arg Gly Ala	54	Gly Gly	58
Tyr Trp Arg Lys	65	Trp Gln Ala Gln	65	Asp Leu Ala Thr	69	Pro Leu	73
Phe Ala His Asn	80	Pro Ser Arg Val	80	Trp Glu Phe Tyr His	84	Tyr Arg	88
Arg Glu Val Met	95	Gly Ser Lys Glu	95	Pro Asn Ala Gly His	99	Arg Ala	103
	110		110		114		118



WO 00/44900

PCT/US00/02237

```

Ile Ala Glu Cys Glu Thr Arg Leu Gly Lys Gln Gly Arg Arg Val
125 130 135
Val Val Ile Thr Gln Asn Ile Asp Glu Leu His Arg Lys Ala Gly
140 145 150
Thr Lys Asn Leu Leu Glu Ile His Gly Ser Leu Phe Lys Thr Arg
155 160 165
Cys Thr Ser Cys Gly Val Val Ala Glu Asn Tyr Lys Ser Pro Ile
170 175 180
Cys Pro Ala Leu Ser Gly Lys Gly Ala Pro Glu Pro Gly Thr Gln
185 190 195
Asp Ala Ser Ile Pro Val Glu Lys Leu Pro Arg Cys Glu Glu Ala
200 205 210
Gly Cys Gly Gly Leu Leu Arg Pro His Val Val Trp Phe Gly Glu
215 220 225
Asn Leu Asp Pro Ala Ile Leu Glu Glu Val Asp Arg Glu Leu Ala
230 235 240
His Cys Asp Leu Cys Leu Val Val Gly Thr Ser Ser Val Val Tyr
245 250 255
Pro Ala Ala Met Phe Ala Pro Gln Val Ala Ala Arg Gly Val Pro
260 265 270
Val Ala Glu Phe Asn Thr Glu Thr Thr Pro Ala Thr Asn Arg Phe
275 280 285
Arg Phe His Phe Gln Gly Pro Cys Gly Thr Thr Leu Pro Glu Ala
290 295 300
Leu Ala Cys His Glu Asn Glu Thr Val Ser
305 310

```

&lt;210&gt; 26

&lt;211&gt; 331

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt; -

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 3988457CD1

&lt;400&gt; 26

```

Met Ala Ile Asn Arg Phe Arg Leu Glu Asn Asp Leu Glu Glu Leu
1 5 10 15
Ala Leu Tyr Gln Ile Gln Leu Leu Lys Asp Leu Arg His Thr Glu
20 25 30
Asn Glu Glu Asp Lys Val Ser Ser Ser Ser Phe Arg Gln Arg Met
35 40 45
Leu Gly Asn Leu Leu Arg Pro Pro Tyr Glu Arg Pro Glu Leu Pro
50 55 60
Thr Cys Leu Tyr Val Ile Gly Leu Thr Gly Ile Ser Gly Ser Gly
65 70 75
Lys Ser Ser Ile Ala Gln Arg Leu Lys Gly Leu Gly Ala Phe Val
80 85 90
Ile Asp Ser Asp His Leu Gly His Arg Ala Tyr Ala Pro Gly Gly
95 100 105
Pro Ala Tyr Gln Pro Val Val Glu Ala Phe Gly Thr Asp Ile Leu
110 115 120
His Lys Asp Gly Ile Ile Asn Arg Lys Val Leu Gly Ser Arg Val
125 130 135
Phe Gly Asn Lys Lys Gln Leu Lys Ile Leu Thr Asp Ile Met Trp
140 145 150
Pro Ile Ile Ala Lys Leu Ala Arg Glu Glu Met Asp Arg Ala Val
155 160 165
Ala Glu Gly Lys Arg Val Cys Val Ile Asp Ala Ala Val Leu Leu
170 175 180

```

29/91

SUBSTITUTE SHEET (RULE 26)

Glu Ala Gly Trp Gln Asn Leu Val His Glu Val Trp Thr Ala Val  
 185 190 195  
 Ile Pro Glu Thr Glu Ala Val Arg Arg Ile Val Glu Arg Asp Gly  
 200 205 210  
 Leu Ser Glu Ala Ala Gln Ser Arg Leu Gln Ser Gln Met Ser  
 215 220 225  
 Gly Gln Gln Leu Val Glu Gln Ser His Val Val Leu Ser Ser Pro  
 230 235 240  
 Cys Gly Ser Arg Ile Ser Pro Asn Ala Arg Trp Arg Lys Pro Gly  
 245 250 255  
 Pro Ser Cys Arg Ser Ala Phe Pro Arg Leu Ile Arg Pro Ser Thr  
 260 265 270  
 Glu Lys Phe Ser Val Gly Pro Asp Trp Leu Leu Glu Leu Thr Ser  
 275 280 285  
 Asp Pro Val Val Arg Arg Asn Gly Gly Leu Asp Ala His Pro Gly  
 290 295 300  
 Ser Gly Pro Glu Val Gln Ala Ile Leu Cys Arg Thr Trp Pro Gly  
 305 310 315  
 Leu Val Asp Thr Gly Ser Leu Pro Asn Thr Leu Val Phe Gly Gln  
 320 325 330  
 His

&lt;210&gt; 27

&lt;211&gt; 200

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 131890CD1

&lt;400&gt; 27

Met Met Thr Ala Glu Ser Arg Glu Ala Thr Gly Leu Ser Pro Gln  
 1 5 10 15  
 Ala Ala Gln Glu Lys Asp Gly Ile Val Ile Val Lys Val Glu Glu  
 20 25 30  
 Glu Asp Glu Glu Asp His Met Trp Gly Gln Asp Ser Thr Leu Gln  
 35 40 45  
 Asp Thr Pro Pro Pro Asp Pro Glu Ile Phe Arg Gln Arg Phe Arg  
 50 55 60  
 Arg Phe Cys Tyr Gln Asn Thr Phe Gly Pro Arg Glu Ala Leu Ser  
 65 70 75  
 Arg Leu Lys Glu Leu Cys His Gln Trp Leu Arg Pro Glu Ile Asn  
 80 85 90  
 Thr Lys Glu Gln Ile Leu Glu Leu Leu Val Leu Glu Gln Phe Leu  
 95 100 105  
 Ser Ile Leu Pro Lys Glu Leu Gln Val Trp Leu Gln Glu Tyr Arg  
 110 115 120  
 Pro Asp Ser Gly Glu Glu Ala Val Thr Leu Leu Asp Leu Glu  
 125 130 135  
 Leu Asp Leu Ser Gly Gln Gln Val Pro Gly Gln Val His Gly Pro  
 140 145 150  
 Glu Met Leu Ala Arg Gly Met Val Pro Leu Asp Pro Val Gln Glu  
 155 160 165  
 Ser Ser Ser Phe Asp Leu His His Glu Ala Thr Gln Ser His Phe  
 170 175 180  
 Lys His Ser Ser Arg Lys Pro Arg Leu Leu Gln Ser Arg Gly Lys  
 185 190 195  
 Lys Gln Gly Phe Ile

200

<210> 28  
 <211> 100  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 238642CD1

<400> 28  
 Met Gln Lys Pro Cys Lys Glu Asn Glu Gly Lys Pro Lys Cys Ser  
 1 5 10 15  
 Val Pro Lys Arg Glu Glu Lys Arg Pro Tyr Gly Glu Phe Glu Arg  
 20 25 30  
 Gln Gln Thr Glu Gly Asn Phe Arg Gln Arg Leu Leu Gln Ser Leu  
 35 40 45  
 Glu Glu Phe Lys Glu Asp Ile Asp Tyr Arg His Phe Lys Asp Glu  
 50 55 60  
 Glu Met Thr Arg Glu Gly Asp Glu Met Glu Arg Cys Leu Glu Glu  
 65 70 75  
 Ile Arg Gly Leu Arg Lys Lys Phe Arg Ala Leu His Ser Asn His  
 80 85 90  
 Arg His Ser Arg Asp Arg Pro Tyr Pro Ile  
 95 100

<210> 29  
 <211> 528  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 669862CD1

<400> 29  
 Met Ser Ser Pro Tyr Pro Leu Leu Leu Glu Asn Ser Ile Cys Leu  
 1 5 10 15  
 Phe Phe His Phe Leu Pro Asp Phe Asn Phe Thr Thr Glu Ser Asn  
 20 25 30  
 Lys Leu Ser Ser Glu Lys Arg Asn Tyr Glu Val Asn Ala Tyr His  
 35 40 45  
 Gln Glu Thr Trp Lys Arg Asn Lys Thr Phe Asn Leu Met Arg Phe  
 50 55 60  
 Ile Phe Arg Thr Asp Pro Gln Tyr Thr Ile Glu Phe Gly Arg Gln  
 65 70 75  
 Gln Arg Pro Lys Val Gly Cys Phe Ser Gln Met Ile Phe Lys Lys  
 80 85 90  
 His Lys Ser Leu Pro Leu His Lys Arg Asn Thr Arg Glu Lys  
 95 100 105  
 Ser Tyr Glu Cys Lys Glu Tyr Lys Lys Gly Phe Arg Lys Tyr Leu  
 110 115 120  
 His Leu Thr Glu His Leu Arg Asp His Thr Gly Val Ile Pro Tyr  
 125 130 135  
 Glu Cys Asn Glu Cys Gly Lys Ala Phe Val Val Phe Gln His Phe

Ile Arg His Arg	140	Ile His Thr Asp	145	Leu Lys Pro Tyr Glu Cys	150
Asn Gly Cys Glu	155	Ala Phe Arg Phe	160	Tyr Ser Gln Leu Ile Gln	165
His Gln Ile Ile	170	Thr Gly Met Lys	175	Pro Tyr Glu Cys Lys Gln	180
Cys Gly Lys Ala	185	Phe Arg Arg His Ser	190	His Leu Thr Glu His Gln	195
Lys Ile His Val	200	Gly Leu Lys Pro Phe	205	Glu Cys Lys Glu Cys Gly	210
Glu Thr Phe Arg	215	Leu Tyr Arg His Met	220	Cys Leu His Gln Lys Ile	225
His His Gly Val	230	Lys Pro Tyr Lys Cys	235	Lys Glu Cys Gly Lys Ala	240
Phe Gly His Arg	245	Ser Ser Leu Tyr Gln	250	His Lys Lys Ile His Ser	255
Gly Glu Lys Pro	260	Tyr Lys Cys Glu Gln	265	Cys Glu Lys Ala Phe Val	270
Arg Ser Tyr Leu	275	Val Glu His Gln Arg	280	Ser His Thr Gly Glu	285
Lys Pro His Glu	290	Cys Met Glu Cys Gly	295	Lys Ala Phe Ser Lys Gly	300
Ser Ser Leu Leu	305	Lys His Lys Arg Ile	310	His Ser Ser Glu Lys Leu	315
Tyr Asp Cys Lys	320	Asp Cys Gly Lys Ala	325	Phe Cys Arg Gly Ser Gln	330
Leu Thr Gln His	335	Gln Arg Ile His Thr	340	Gly Glu Lys Pro His Glu	345
Cys Lys Glu Cys	350	Gly Lys Thr Phe Lys	355	Leu His Ser Tyr Leu Ile	360
Gln His Gln Ile	365	Ile His Thr Asp Leu	370	Lys Pro Tyr Glu Cys Lys	375
Gln Cys Gly Lys	380	Ala Phe Ser Arg Val	385	Gly Asp Leu Lys Thr His	390
Gln Ser Ile His	395	Ala Gly Glu Lys Pro	400	Tyr Glu Cys Lys Glu Cys	405
Gly Lys Thr Phe	410	Arg Leu Asn Ser Gln	415	Leu Ile Tyr His Gln Thr	420
Ile His Thr Gly	425	Leu Lys Pro Tyr Val	430	Cys Lys Glu Cys Lys Lys	435
Ala Phe Arg Ser	440	Ile Ser Gly Leu Ser	445	Gln His Lys Arg Ile His	450
Thr Gly Glu Lys	455	Pro Tyr Glu Cys Lys	460	Glu Cys Asp Lys Ala Phe	465
Asn Arg Ser Asp	470	Arg Leu Thr Gln His	475	Glu Thr Ile His Thr Gly	480
Val Lys Pro Gln	485	Lys Cys Lys Glu Cys	490	Gly Lys Ala Phe Ser His	495
Cys Tyr Gln Leu	500	Ser Gln His Gln Arg	505	Phe His His Gly Glu Arg	510
Leu Leu Met	515		520		525

&lt;210&gt; 30

&lt;211&gt; 350

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 1003663CD1

```

<400> 30
Met His Pro Ala Ala Phe Pro Leu Pro Val Val Val Ala Ala Val
1      5      10      15
Leu Trp Gly Ala Ala Pro Thr Arg Gly Leu Ile Arg Ala Thr Ser
20      25      30
Asp His Asn Ala Ser Met Asp Phe Ala Asp Leu Pro Ala Leu Phe
35      40      45
Gly Ala Thr Leu Ser Gln Glu Gly Leu Gln Gly Phe Leu Val Glu
50      55      60
Ala His Pro Asp Asn Ala Cys Ser Pro Ile Ala Pro Pro Pro Pro
65      70      75
Ala Pro Val Asn Gly Ser Val Phe Ile Ala Leu Leu Arg Arg Phe
80      85      90
Asp Cys Asn Phe Asp Leu Lys Val Leu Asn Ala Gln Lys Ala Gly
95      100      105
Tyr Gly Ala Ala Val Val His Asn Val Asn Ser Asn Glu Leu Leu
110      115      120
Asn Met Val Trp Asn Ser Glu Glu Ile Gln Gln Gln Ile Trp Ile
125      130      135
Pro Ser Val Phe Ile Gly Glu Arg Ser Ser Glu Tyr Leu Arg Ala
140      145      150
Leu Phe Val Tyr Glu Lys Gly Ala Arg Val Leu Leu Val Pro Asp
155      160      165
Asn Thr Phe Pro Leu Gly Tyr Tyr Leu Ile Pro Phe Thr Gly Ile
170      175      180
Val Gly Leu Leu Val Leu Ala Met Gly Ala Val Met Ile Ala Arg
185      190      195
Cys Ile Gln His Arg Lys Arg Leu Gln Arg Asn Arg Leu Thr Lys
200      205      210
Glu Gln Leu Lys Gln Ile Pro Thr His Asp Tyr Gln Lys Gly Asp
215      220      225
Gln Tyr Asp Val Cys Ala Ile Cys Leu Asp Glu Tyr Glu Asp Gly
230      235      240
Asp Lys Leu Arg Val Leu Pro Cys Ala His Ala Tyr His Ser Arg
245      250      255
Cys Val Asp Pro Trp Leu Thr Gln Thr Arg Lys Thr Cys Pro Ile
260      265      270
Cys Lys Gln Pro Val His Arg Gly Pro Gly Asp Glu Asp Gln Glu
275      280      285
Glu Glu Thr Gln Gly Gln Glu Glu Gly Asp Glu Gly Glu Pro Arg
290      295      300
Asp His Pro Ala Ser Glu Arg Thr Pro Leu Leu Gly Ser Ser Pro
305      310      315
Thr Leu Pro Thr Ser Phe Gly Ser Leu Ala Pro Ala Pro Leu Val
320      325      330
Phe Pro Gly Pro Ser Thr Asp Pro Pro Leu Ser Pro Pro Ser Ser
335      340      345
Pro Val Ile Leu Val
350

```

```

<210> 31
<211> 315
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> misc-feature
<223> Incyte ID No.: 1432557CD1

```

<400> 31  
 Met Ala Ala Leu Gly Val Leu Glu Ser Asp Leu Pro Ser Ala Val  
 1 5 10 15  
 Thr Leu Leu Lys Asn Leu Gln Glu Gln Val Met Ala Val Thr Ala  
 20 25 30  
 Gln Val Lys Ser Leu Thr Gln Lys Val Gln Ala Gly Ala Tyr Pro  
 35 40 45  
 Thr Glu Lys Gly Leu Ser Phe Leu Glu Val Lys Asp Gln Leu Leu  
 50 55 60  
 Leu Met Tyr Leu Met Asp Leu Thr His Leu Ile Leu Asp Lys Ala  
 65 70 75  
 Ser Gly Gly Ser Leu Gln Gly His Asp Ala Val Leu Arg Leu Val  
 80 85 90  
 Glu Ile Arg Thr Val Leu Glu Lys Leu Arg Pro Leu Asp Gln Lys  
 95 100 105  
 Leu Lys Tyr Gln Ile Asp Lys Leu Ile Lys Thr Ala Val Thr Gly  
 110 115 120  
 Ser Leu Ser Glu Asn Asp Pro Leu Arg Phe Lys Pro His Pro Ser  
 125 130 135  
 Asn Met Met Ser Lys Leu Ser Ser Glu Asp Glu Glu Glu Asp Glu  
 140 145 150  
 Ala Glu Asp Asp Gln Ser Glu Ala Ser Gly Lys Lys Ser Val Lys  
 155 160 165  
 Gly Val Ser Lys Lys Tyr Val Pro Pro Arg Leu Val Pro Val His  
 170 175 180  
 Tyr Asp Glu Thr Glu Ala Glu Arg Glu Lys Lys Arg Leu Glu Arg  
 185 190 195  
 Ala Lys Arg Arg Ala Leu Ser Ser Ser Val Ile Arg Glu Leu Lys  
 200 205 210  
 Glu Gln Tyr Ser Asp Ala Pro Glu Glu Ile Arg Asp Ala Arg His  
 215 220 225  
 Pro His Val Thr Arg Gln Ser Gln Glu Asp Gln His Arg Ile Asn  
 230 235 240  
 Tyr Glu Glu Ser Met Met Val Arg Leu Ser Val Ser Lys Arg Glu  
 245 250 255  
 Lys Gly Arg Arg Lys Arg Ala Asn Val Met Ser Ser Gln Leu His  
 260 265 270  
 Ser Leu Thr His Phe Ser Asp Ile Ser Ala Leu Thr Gly Gly Thr  
 275 280 285  
 Val His Leu Asp Glu Asp Gln Asn Pro Ile Lys Lys Arg Lys Lys  
 290 295 300  
 Ile Pro Gln Lys Gly Arg Lys Lys Lys Gly Phe Arg Arg Arg Arg  
 305 310 315

<210> 32  
 <211> 120  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> 1441770CD1

<400> 32  
 Met Asp Asp Ser Lys Val Val Gly Gly Lys Val Lys Lys Pro Gly  
 1 5 10 15  
 Lys Arg Gly Arg Lys Pro Ala Lys Ile Asp Leu Lys Ala Lys Leu  
 20 25 30  
 Glu Arg Ser Arg Gln Ser Ala Arg Glu Cys Arg Ala Arg Lys Lys  
 35 40 45

Leu	Arg	Tyr	Gln	Tyr	Leu	Glu	Glu	Leu	Val	Ser	Ser	Arg	Glu	Arg	
				50					55						60
Ala	Ile	Cys	Ala	Leu	Arg	Glu	Glu	Leu	Glu	Met	Tyr	Lys	Gln	Trp	
				65					70						75
Cys	Met	Ala	Met	Asp	Gln	Gly	Lys	Ile	Pro	Ser	Glu	Ile	Lys	Ala	
				80					85						90
Leu	Leu	Thr	Gly	Glu	Glu	Gln	Asn	Lys	Ser	Gln	Gln	Asn	Ser	Ser	
				95					100						105
Arg	His	Thr	Lys	Ala	Gly	Lys	Thr	Asp	Ala	Asn	Ser	Asn	Ser	Trp	
				110					115						120

&lt;210&gt; 33

&lt;211&gt; 326

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 1456684CD1

&lt;400&gt; 33

Met	Gln	Glu	Glu	Pro	Leu	Pro	Gln	Gly	Asn	Asp	Pro	Glu	Pro	Ser	
1				5					10						15
Gly	Asp	Ser	Pro	Leu	Gly	Leu	Cys	Gln	Ser	Glu	Cys	Met	Glu	Met	
				20					25						30
Ser	Glu	Val	Phe	Asp	Asp	Ala	Ser	Asp	Gln	Asp	Ser	Thr	Asp	Lys	
				35					40						45
Pro	Trp	Arg	Pro	Tyr	Tyr	Asn	Tyr	Lys	Pro	Lys	Lys	Lys	Ser	Arg	
				50					55						60
Gln	Leu	Lys	Lys	Met	Arg	Lys	Val	Asn	Trp	Arg	Lys	Glu	His	Gly	
				65					70						75
Asn	Arg	Ser	Pro	Ser	His	Lys	Cys	Lys	Tyr	Pro	Ala	Glu	Leu	Asp	
				80					85						90
Cys	Ala	Val	Gly	Lys	Ala	Pro	Gln	Asp	Lys	Pro	Phe	Glu	Glu	Glu	
				95					100						105
Glu	Thr	Lys	Glu	Met	Pro	Lys	Leu	Gln	Cys	Glu	Leu	Cys	Asp	Gly	
				110					115						120
Asp	Lys	Ala	Val	Gly	Ala	Gly	Asn	Gln	Gly	Arg	Pro	His	Arg	His	
				125					130						135
Leu	Thr	Ser	Arg	Pro	Tyr	Ala	Cys	Glu	Leu	Cys	Ala	Lys	Gln	Phe	
				140					145						150
Gln	Ser	Pro	Ser	Thr	Leu	Lys	Met	His	Met	Arg	Cys	His	Thr	Gly	
				155					160						165
Glu	Lys	Pro	Tyr	Gln	Cys	Lys	Thr	Cys	Gly	Arg	Cys	Phe	Ser	Val	
				170					175						180
Gln	Gly	Asn	Leu	Gln	Lys	His	Glu	Arg	Ile	His	Leu	Gly	Leu	Lys	
				185					190						195
Glu	Phe	Val	Cys	Gln	Tyr	Cys	Asn	Lys	Ala	Phe	Thr	Leu	Asn	Glu	
				200					205						210
Thr	Leu	Lys	Ile	His	Glu	Arg	Ile	His	Thr	Gly	Glu	Lys	Arg	Tyr	
				215					220						225
His	Cys	Gln	Phe	Cys	Phe	Gln	Arg	Phe	Leu	Tyr	Leu	Ser	Thr	Lys	
				230					235						240
Arg	Asn	His	Glu	Gln	Arg	His	Ile	Arg	Glu	His	Asn	Gly	Lys	Gly	
				245					250						255
Tyr	Ala	Cys	Phe	Gln	Cys	Pro	Lys	Ile	Cys	Lys	Thr	Ala	Ala	Ala	
				260					265						270
Leu	Gly	Met	His	Gln	Lys	Lys	His	Leu	Phe	Lys	Ser	Pro	Ser	Gln	
				275					280						285

WO 00/44900

PCT/US00/02237

Gln	Glu	Lys	Ile	Gly	Asp	Val	Cys	His	Glu	Asn	Ser	Asn	Pro	Leu	
				290						295				300	
Glu	Asn	Gln	His	Phe	Ile	Gly	Ser	Glu	Asp	Asn	Asp	Gln	Lys	Asp	
				305					310					315	
Asn	Ile	Gln	Thr	Gly	Val	Glu	Asn	Val	Val	Leu					
				320					325						

<210> 34  
 <211> 106  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 1602916CD1

<400> 34															
Met	Phe	Pro	Trp	Met	Arg	Pro	Gln	Ala	Ala	Pro	Gly	Arg	Arg	Arg	
1				5					10					15	
Gly	Arg	Gln	Thr	Tyr	Ser	Arg	Phe	Gln	Thr	Leu	Glu	Leu	Glu	Lys	
				20					25					30	
Glu	Phe	Leu	Phe	Asn	Pro	Tyr	Leu	Thr	Arg	Lys	Arg	Arg	Ile	Glu	
				35					40					45	
Val	Ser	His	Ala	Leu	Ala	Leu	Thr	Glu	Arg	Gln	Val	Lys	Ile	Trp	
				50					55					60	
Phe	Gln	Asn	Arg	Arg	Met	Lys	Trp	Lys	Lys	Glu	Asn	Asn	Lys	Asp	
				65					70					75	
Lys	Phe	Pro	Val	Ser	Arg	Gln	Glu	Val	Lys	Asp	Gly	Glu	Thr	Lys	
				80					85					90	
Lys	Glu	Ala	Gln	Glu	Leu	Glu	Glu	Asp	Arg	Ala	Glu	Arg	Leu	Thr	
				95					100					105	
Asn															

<210> 35  
 <211> 209  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 1692816CD1

<400> 35															
Met	Asn	Pro	Ser	Met	Lys	Gln	Lys	Gln	Glu	Glu	Ile	Lys	Glu	Asn	
1				5					10					15	
Ile	Lys	Asn	Ser	Ser	Val	Pro	Arg	Arg	Thr	Leu	Lys	Met	Ile	Gln	
				20					25					30	
Pro	Ser	Ala	Ser	Gly	Ser	Leu	Val	Gly	Arg	Glu	Asn	Glu	Leu	Ser	
				35					40					45	
Ala	Gly	Leu	Ser	Lys	Arg	Lys	His	Arg	Asn	Asp	His	Leu	Thr	Ser	
				50					55					60	
Thr	Thr	Ser	Ser	Pro	Gly	Val	Ile	Val	Pro	Glu	Ser	Ser	Glu	Asn	
				65					70					75	
Lys	Asn	Leu	Gly	Gly	Val	Thr	Gln	Glu	Ser	Phe	Asp	Leu	Met	Ile	
				80					85					90	
Lys	Glu	Asn	Pro	Ser	Ser	Gln	Tyr	Trp	Lys	Glu	Val	Ala	Glu	Lys	



			95					100				105		
Arg	Arg	Lys	Ala	Leu	Tyr	Glu	Ala	Leu	Lys	Glu	Asn	Glu	Lys	Leu
			110											120
His	Lys	Glu	Ile	Glu	Gln	Lys	Asp	Asn	Glu	Ile	Ala	Arg	Leu	Lys
			125											135
Lys	Glu	Asn	Lys	Glu	Leu	Ala	Glu	Val	Ala	Glu	His	Val	Gln	Tyr
			140											150
Met	Ala	Glu	Leu	Ile	Glu	Arg	Leu	Asn	Gly	Glu	Pro	Leu	Asp	Asn
			155											165
Phe	Glu	Ser	Leu	Asp	Asn	Gln	Glu	Phe	Asp	Ser	Glu	Glu	Glu	Thr
			170											180
Val	Glu	Asp	Ser	Leu	Val	Glu	Asp	Ser	Glu	Ile	Gly	Thr	Cys	Ala
			185											195
Glu	Gly	Thr	Val	Ser	Ser	Thr	Asp	Ala	Lys	Pro	Cys	Ile		
			200											

<210> 36  
 <211> 212  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 1968191CD1

<400> 36  
 Met Leu Gly Asn Glu Trp Ser Lys Leu Pro Pro Glu Glu Lys Gln  
 1 5 10 15  
 Arg Tyr Leu Asp Glu Ala Asp Arg Asp Lys Glu Arg Tyr Met Lys  
 20 25 30  
 Glu Leu Glu Gln Tyr Gln Lys Thr Glu Ala Tyr Lys Val Phe Ser  
 35 40 45  
 Arg Lys Thr Gln Asp Arg Gln Lys Gly Lys Ser His Arg Gln Asp  
 50 55 60  
 Ala Ala Arg Gln Ala Thr His Asp His Glu Lys Glu Thr Glu Val  
 65 70 75  
 Lys Glu Arg Ser Val Phe Asp Ile Pro Ile Phe Thr Glu Glu Phe  
 80 85 90  
 Leu Asn His Ser Lys Ala Arg Glu Ala Glu Leu Arg Gln Leu Arg  
 95 100 105  
 Lys Ser Asn Met Glu Phe Glu Glu Arg Asn Ala Ala Leu Gln Lys  
 110 115 120  
 His Val Glu Ser Met Arg Thr Ala Val Glu Lys Leu Glu Val Asp  
 125 130 135  
 Val Ile Gln Glu Arg Ser Arg Asn Thr Val Leu Gln Gln His Leu  
 140 145 150  
 Glu Thr Leu Arg Gln Val Leu Thr Ser Ser Phe Ala Ser Met Pro  
 155 160 165  
 Leu Pro Gly Ser Gly Glu Thr Pro Thr Val Asp Thr Ile Asp Ser  
 170 175 180  
 Tyr Met Asn Arg Leu His Ser Ile Ile Leu Ala Asn Pro Gln Asp  
 185 190 195  
 Asn Glu Asn Phe Ile Ala Thr Val Arg Glu Val Val Asn Arg Leu  
 200 205 210  
 Asp Arg

WO 00/44900

PCT/US00/02237

<210> 37  
<211> 359  
<212> PRT  
<213> Homo sapiens

<220>  
<221> misc-feature  
<223> Incyte ID No.: 2052061CD1

<400> 37  
Met Val Asp Met Asp Lys Leu Ile Asn Asn Leu Glu Val Gln Leu  
1 5 10 15  
Asn Ser Glu Gly Gly Ser Met Gln Val Phe Lys Gln Val Thr Ala  
20 25 30  
Ser Val Arg Asn Arg Asp Pro Pro Glu Ile Glu Tyr Thr Ser Asn  
35 40 45  
Met Thr Ser Pro Thr Leu Leu Asp Ala Asn Pro Met Glu Asn Pro  
50 55 60  
  
Ala Leu Phe Asn Asp Ile Lys Ile Glu Pro Pro Glu Glu Leu Leu  
65 70 75  
Ala Ser Asp Phe Ser Leu Pro Gln Val Glu Pro Val Asp Leu Ser  
80 85 90  
Phe His Lys Pro Lys Ala Pro Leu Gln Pro Ala Ser Met Leu Gln  
95 100 105  
Ala Pro Ile Arg Pro Pro Lys Pro Gln Ser Ser Pro Gln Thr Leu  
110 115 120  
Val Val Ser Thr Ser Thr Ser Asp Met Ser Thr Ser Ala Asn Ile  
125 130 135  
Pro Thr Val Leu Thr Pro Gly Ser Val Leu Thr Ser Ser Gln Ser  
140 145 150  
Thr Gly Ser Gln Gln Ile Leu His Val Ile His Thr Ile Pro Ser  
155 160 165  
Val Ser Leu Pro Asn Lys Met Gly Gly Leu Lys Thr Ile Pro Val  
170 175 180  
Val Val Gln Ser Leu Pro Met Val Tyr Thr Thr Leu Pro Ala Asp  
185 190 195  
Gly Gly Pro Ala Ala Ile Thr Val Pro Leu Ile Gly Gly Asp Gly  
200 205 210  
Lys Asn Ala Gly Ser Val Lys Val Asp Pro Thr Ser Met Ser Pro  
215 220 225  
Leu Glu Ile Pro Ser Asp Ser Glu Glu Ser Thr Ile Glu Ser Gly  
230 235 240  
Ser Ser Ala Leu Gln Ser Leu Gln Gly Leu Gln Gln Glu Pro Ala  
245 250 255  
Ala Met Ala Gln Met Gln Gly Glu Glu Ser Leu Asp Leu Lys Arg  
260 265 270  
Arg Arg Ile His Gln Cys Asp Phe Ala Gly Cys Ser Lys Val Tyr  
275 280 285  
Thr Lys Ser Ser His Leu Lys Ala His Arg Arg Ile His Thr Gly  
290 295 300  
Glu Lys Pro Tyr Lys Cys Thr Trp Asp Gly Cys Ser Trp Lys Phe  
305 310 315  
Ala Arg Ser Asp Glu Leu Thr Arg His Phe Arg Lys His Thr Gly  
320 325 330  
Ile Lys Pro Phe Arg Cys Thr Asp Cys Asn Arg Ser Phe Ser Arg  
335 340 345  
Ser Asp His Leu Ser Leu His Arg Arg Arg His Asp Thr Met  
350 355

<210> 38  
 <211> 445  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 2056207CD1

<400> 38

Met	Val	Glu	Cys	Ile	Arg	Glu	Val	Asn	Glu	Val	Ile	Gln	Asn	Pro	
1				5				10						15	
Ala	Thr	Ile	Thr	Arg	Ile	Leu	Leu	Ser	His	Phe	Asn	Trp	Asp	Lys	
				20				25						30	
Glu	Lys	Leu	Met	Glu	Arg	Tyr	Phe	Asp	Gly	Asn	Leu	Glu	Lys	Leu	
				35				40						45	
Phe	Ala	Glu	Cys	His	Val	Ile	Asn	Pro	Ser	Lys	Lys	Ser	Arg	Thr	
				50				55						60	
Arg	Gln	Met	Asn	Thr	Arg	Ser	Ser	Ala	Gln	Asp	Met	Pro	Cys	Gln	
				65				70						75	
Ile	Cys	Tyr	Leu	Asn	Tyr	Pro	Asn	Ser	Tyr	Phe	Thr	Gly	Leu	Glu	
				80				85						90	
Cys	Gly	His	Lys	Phe	Cys	Met	Gln	Cys	Trp	Ser	Glu	Tyr	Leu	Thr	
				95				100						105	
Thr	Lys	Ile	Met	Glu	Glu	Gly	Met	Gly	Gln	Thr	Ile	Ser	Cys	Pro	
				110				115						120	
Ala	His	Gly	Cys	Asp	Ile	Leu	Val	Asp	Asp	Asn	Thr	Val	Met	Arg	
				125				130						135	
Leu	Ile	Thr	Asp	Ser	Lys	Val	Lys	Leu	Lys	Tyr	Gln	His	Leu	Ile	
				140				145						150	
Thr	Asn	Ser	Phe	Val	Glu	Cys	Asn	Arg	Leu	Leu	Lys	Trp	Cys	Pro	
				155				160						165	
Ala	Pro	Asp	Cys	His	His	Val	Val	Lys	Val	Gln	Tyr	Pro	Asp	Ala	
				170				175						180	
Lys	Pro	Val	Arg	Cys	Lys	Cys	Gly	Arg	Gln	Phe	Cys	Phe	Asn	Cys	
				185				190						195	
Gly	Glu	Asn	Trp	His	Asp	Pro	Val	Lys	Cys	Lys	Trp	Leu	Lys	Lys	
				200				205						210	
Trp	Ile	Lys	Lys	Cys	Asp	Asp	Asp	Ser	Glu	Thr	Ser	Asn	Trp	Ile	
				215				220						225	
Ala	Ala	Asn	Thr	Lys	Glu	Cys	Pro	Lys	Cys	His	Val	Thr	Ile	Glu	
				230				235						240	
Lys	Asp	Gly	Gly	Cys	Asn	His	Met	Val	Cys	Arg	Asn	Gln	Asn	Cys	
				245				250						255	
Lys	Ala	Glu	Phe	Cys	Trp	Val	Cys	Leu	Gly	Pro	Trp	Glu	Pro	His	
				260				265						270	
Gly	Ser	Ala	Trp	Tyr	Asn	Cys	Asn	Arg	Tyr	Asn	Glu	Asp	Asp	Ala	
				275				280						285	
Lys	Ala	Ala	Arg	Asp	Ala	Gln	Glu	Arg	Ser	Arg	Ala	Ala	Leu	Gln	
				290				295						300	
Arg	Tyr	Leu	Phe	Tyr	Cys	Asn	Arg	Tyr	Met	Asn	His	Met	Gln	Ser	
				305				310						315	
Leu	Arg	Phe	Glu	His	Lys	Leu	Tyr	Ala	Gln	Val	Lys	Gln	Lys	Met	
				320				325						330	
Glu	Glu	Met	Gln	Gln	His	Asn	Met	Ser	Trp	Ile	Glu	Val	Gln	Phe	
				335				340						345	
Leu	Lys	Lys	Ala	Val	Asp	Val	Leu	Cys	Gln	Cys	Arg	Ala	Thr	Leu	
				350				355						360	
Met	Tyr	Thr	Tyr	Val	Phe	Ala	Phe	Tyr	Leu	Lys	Lys	Asn	Asn	Gln	
				365				370						375	
Ser	Ile	Ile	Phe	Glu	Asn	Asn	Gln	Ala	Asp	Leu	Glu	Asn	Ala	Thr	
				380				385						390	

Glu	Val	Leu	Ser	Gly	Tyr	Leu	Glu	Arg	Asp	Ile	Ser	Gln	Asp	Ser	
				395					400					405	
Leu	Gln	Asp	Ile	Lys	Gln	Lys	Val	Gln	Asp	Lys	Tyr	Arg	Tyr	Cys	
				410					415					420	
Glu	Ser	Arg	Arg	Arg	Val	Leu	Leu	Gln	His	Val	His	Glu	Gly	Tyr	
				425					430					435	
Glu	Lys	Asp	Leu	Trp	Glu	Tyr	Ile	Glu	Asp						
				440					445						

<210> 39  
 <211> 433  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 2101803CD1

<400> 39

Met	Arg	Ala	Glu	Gly	Leu	Gly	Gly	Leu	Glu	Arg	Phe	Cys	Ser	Pro	
1				5					10					15	
Gly	Lys	Gly	Arg	Gly	Leu	Arg	Ala	Leu	Gln	Pro	Phe	Gln	Val	Gly	
				20					25					30	
Asp	Leu	Leu	Phe	Ser	Cys	Pro	Ala	Tyr	Ala	Tyr	Val	Leu	Thr	Val	
				35					40					45	
Asn	Glu	Arg	Gly	Asn	His	Cys	Glu	Tyr	Cys	Phe	Thr	Arg	Lys	Glu	
				50					55					60	
Gly	Leu	Ser	Lys	Cys	Gly	Arg	Cys	Lys	Gln	Ala	Phe	Tyr	Cys	Asn	
				65					70					75	
Val	Glu	Cys	Gln	Lys	Glu	Asp	Trp	Pro	Met	His	Lys	Leu	Glu	Cys	
				80					85					90	
Ser	Pro	Met	Val	Val	Phe	Gly	Glu	Asn	Trp	Asn	Pro	Ser	Glu	Thr	
				95					100					105	
Val	Arg	Leu	Thr	Ala	Arg	Ile	Leu	Ala	Lys	Gln	Lys	Ile	His	Pro	
				110					115					120	
Glu	Arg	Thr	Pro	Ser	Glu	Lys	Leu	Leu	Ala	Val	Lys	Glu	Phe	Glu	
				125					130					135	
Ser	His	Leu	Asp	Lys	Leu	Asp	Asn	Glu	Lys	Lys	Asp	Leu	Ile	Gln	
				140					145					150	
Ser	Asp	Ile	Ala	Ala	Leu	His	His	Phe	Tyr	Ser	Lys	His	Leu	Glu	
				155					160					165	
Phe	Pro	Asp	Asn	Asp	Ser	Leu	Val	Val	Leu	Phe	Ala	Gln	Val	Asn	
				170					175					180	
Cys	Asn	Gly	Phe	Thr	Ile	Glu	Asp	Glu	Glu	Leu	Ser	His	Leu	Gly	
				185					190					195	
Ser	Ala	Ile	Phe	Pro	Asp	Val	Ala	Leu	Met	Asn	His	Ser	Cys	Cys	
				200					205					210	
Pro	Asn	Val	Ile	Val	Thr	Tyr	Lys	Gly	Thr	Leu	Ala	Glu	Val	Arg	
				215					220					225	
Ala	Val	Gln	Glu	Ile	Lys	Pro	Gly	Glu	Glu	Val	Phe	Thr	Ser	Tyr	
				230					235					240	
Ile	Asp	Leu	Leu	Tyr	Pro	Thr	Glu	Asp	Arg	Asn	Asp	Arg	Leu	Arg	
				245					250					255	
Asp	Ser	Tyr	Phe	Phe	Thr	Cys	Glu	Cys	Gln	Glu	Cys	Thr	Thr	Lys	
				260					265					270	
Asp	Lys	Asp	Lys	Ala	Lys	Val	Glu	Ile	Arg	Lys	Leu	Ser	Asp	Pro	
				275					280					285	
Pro	Lys	Ala	Glu	Ala	Ile	Arg	Asp	Met	Val	Arg	Tyr	Ala	Arg	Asn	
				290					295					300	

WO 00/44900

PCT/US00/02237

Val	Ile	Glu	Glu	Phe	Arg	Arg	Ala	Lys	His	Tyr	Lys	Ser	Pro	Ser	
				305					310					315	
Glu	Leu	Leu	Glu	Ile	Cys	Glu	Leu	Ser	Gln	Glu	Lys	Met	Ser	Ser	
				320					325					330	
Val	Phe	Glu	Asp	Ser	Asn	Val	Tyr	Met	Leu	His	Met	Met	Tyr	Gln	
				335					340					345	
Ala	Met	Gly	Val	Cys	Leu	Tyr	Met	Gln	Asp	Trp	Glu	Gly	Ala	Leu	
				350					355					360	
Gln	Tyr	Gly	Gln	Lys	Ile	Ile	Lys	Pro	Tyr	Ser	Lys	His	Tyr	Pro	
				365					370					375	
Leu	Tyr	Ser	Leu	Asn	Val	Ala	Ser	Met	Trp	Leu	Lys	Leu	Gly	Arg	
				380					385					390	
Leu	Tyr	Met	Gly	Leu	Glu	His	Lys	Ala	Ala	Gly	Glu	Lys	Ala	Leu	
				395					400					405	
Lys	Lys	Ala	Ile	Ala	Ile	Met	Glu	Val	Ala	His	Gly	Lys	Asp	His	
				410					415					420	
Pro	Tyr	Ile	Ser	Glu	Ile	Lys	Gln	Glu	Ile	Glu	Ser	His			
				425					430						

<210> 40

<211> 355

<212> PRT

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 2112362CD1

<400> 40

Met	Ser	Val	Asn	Tyr	Ala	Ala	Gly	Leu	Ser	Pro	Tyr	Ala	Asp	Lys	
1				5					10					15	
Gly	Lys	Cys	Gly	Leu	Pro	Glu	Ile	Phe	Asp	Pro	Pro	Glu	Glu	Leu	
				20					25					30	
Glu	Arg	Lys	Val	Trp	Glu	Leu	Ala	Arg	Leu	Val	Trp	Gln	Ser	Ser	
				35					40					45	
Asn	Val	Val	Phe	His	Thr	Gly	Ala	Gly	Ile	Ser	Thr	Ala	Ser	Gly	
				50					55					60	
Ile	Pro	Asp	Phe	Arg	Gly	Pro	His	Gly	Val	Trp	Thr	Met	Glu	Glu	
				65					70					75	
Arg	Gly	Leu	Ala	Pro	Lys	Phe	Asp	Thr	Thr	Phe	Glu	Ser	Ala	Arg	
				80					85					90	
Pro	Thr	Gln	Thr	His	Met	Ala	Leu	Val	Gln	Leu	Glu	Arg	Val	Gly	
				95					100					105	
Leu	Leu	Arg	Phe	Leu	Val	Ser	Gln	Asn	Val	Asp	Gly	Leu	His	Val	
				110					115					120	
Arg	Ser	Gly	Phe	Pro	Arg	Asp	Lys	Leu	Ala	Glu	Leu	His	Gly	Asn	
				125					130					135	
Met	Phe	Val	Glu	Glu	Cys	Ala	Lys	Cys	Lys	Thr	Gln	Tyr	Val	Arg	
				140					145					150	
Asp	Thr	Val	Val	Gly	Thr	Met	Gly	Leu	Lys	Ala	Thr	Gly	Arg	Leu	
				155					160					165	
Cys	Thr	Val	Ala	Lys	Ala	Arg	Gly	Leu	Arg	Ala	Cys	Arg	Gly	Glu	
				170					175					180	
Leu	Arg	Asp	Thr	Ile	Leu	Asp	Trp	Glu	Asp	Ser	Leu	Pro	Asp	Arg	
				185					190					195	
Asp	Leu	Ala	Leu	Ala	Asp	Glu	Ala	Ser	Arg	Asn	Ala	Asp	Leu	Ser	
				200					205					210	
Ile	Thr	Leu	Gly	Thr	Ser	Leu	Gln	Ile	Arg	Pro	Ser	Gly	Asn	Leu	
				215					220					225	

```

Pro Leu Ala Thr Lys Arg Arg Gly Gly Arg Leu Val Ile Val Asn
230 235 240
Leu Gln Pro Thr Lys His Asp Arg His Ala Asp Leu Arg Ile His
245 250 255
Gly Tyr Val Asp Glu Val Met Thr Arg Leu Met Lys His Leu Gly
260 265 270
Leu Glu Ile Pro Ala Trp Asp Gly Pro Arg Val Leu Glu Arg Ala
275 280 285
Leu Pro Pro Leu Pro Arg Pro Pro Thr Pro Lys Leu Glu Pro Lys
290 295 300
Glu Glu Ser Pro Thr Arg Ile Asn Gly Ser Ile Pro Ala Gly Pro
305 310 315
Lys Gln Glu Pro Cys Ala Gln His Asn Gly Ser Glu Pro Ala Ser
320 325 330
Pro Lys Arg Glu Arg Pro Thr Ser Pro Ala Pro His Arg Pro Pro
335 340 345
Lys Arg Val Lys Ala Lys Ala Val Pro Ser
350 355

```

```

<210> 41
<211> 443
<212> PRT
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 2117346CD1

```

```

<400> 41
Met Asp Arg Leu Gly Ser Phe Ser Asn Asp Pro Ser Asp Lys Pro
1 5 10 15
Pro Cys Arg Gly Cys Ser Ser Tyr Leu Met Glu Pro Tyr Ile Lys
20 25 30
Cys Ala Glu Cys Gly Pro Pro Pro Phe Phe Leu Cys Leu Gln Cys
35 40 45
Phe Thr Arg Gly Phe Glu Tyr Lys Lys His Gln Ser Asp His Thr
50 55 60
Tyr Glu Ile Met Thr Ser Asp Phe Pro Val Leu Asp Pro Ser Trp
65 70 75
Thr Ala Gln Glu Glu Met Ala Leu Leu Glu Ala Val Met Asp Cys
80 85 90
Gly Phe Gly Asn Trp Gln Asp Val Ala Asn Gln Met Cys Thr Lys
95 100 105
Thr Lys Glu Glu Cys Glu Lys His Tyr Met Lys His Phe Ile Asn
110 115 120
Asn Pro Leu Phe Ala Ser Thr Leu Leu Asn Leu Lys Gln Ala Glu
125 130 135
Glu Ala Lys Thr Ala Asp Thr Ala Ile Pro Phe His Ser Thr Asp
140 145 150
Asp Pro Pro Arg Pro Thr Phe Asp Ser Leu Leu Ser Arg Asp Met
155 160 165
Ala Gly Tyr Met Pro Ala Arg Ala Asp Phe Ile Glu Glu Phe Asp
170 175 180
Asn Tyr Ala Glu Trp Asp Leu Arg Asp Ile Asp Phe Val Glu Asp
185 190 195
Asp Ser Asp Ile Leu His Ala Leu Lys Met Ala Val Val Asp Ile
200 205 210
Tyr His Ser Arg Leu Lys Glu Arg Gln Arg Arg Lys Lys Ile Ile
215 220 225

```

```

Arg Asp His Gly Leu Ile Asn Leu Arg Lys Phe Gln Leu Met Glu
230 235 240
Arg Arg Tyr Pro Lys Glu Val Gln Asp Leu Tyr Glu Thr Met Arg
245 250 255
Arg Phe Ala Arg Ile Val Gly Pro Val Glu His Asp Lys Phe Ile
260 265 270
Glu Ser His Ala Leu Glu Phe Glu Leu Arg Arg Glu Ile Lys Arg
275 280 285
Leu Gln Glu Tyr Arg Thr Ala Gly Ile Thr Asn Phe Cys Ser Ala
290 295 300
Arg Thr Tyr Asp His Leu Lys Lys Thr Arg Glu Glu Glu Arg Leu
305 310 315
Lys Arg Thr Met Leu Ser Glu Val Leu Gln Tyr Ile Gln Asp Ser
320 325 330
Ser Ala Cys Gln Gln Trp Leu Arg Arg Gln Ala Asp Ile Asp Ser
335 340 345
Gly Leu Ser Pro Ser Ile Pro Met Ala Ser Asn Ser Gly Arg Arg
350 355 360
Ser Ala Pro Pro Leu Asn Leu Thr Gly Leu Pro Gly Thr Glu Lys
365 370 375
Leu Asn Glu Lys Glu Lys Glu Leu Cys Gln Met Val Arg Leu Val
380 385 390
Pro Gly Ala Tyr Leu Glu Tyr Lys Ser Ala Leu Leu Asn Glu Cys
395 400 405
Asn Lys Gln Gly Gly Leu Arg Leu Ala Gln Ala Arg Ala Leu Ile
410 415 420
Lys Ile Asp Val Asn Lys Thr Arg Lys Ile Tyr Asp Phe Leu Ile
425 430 435
Arg Glu Gly Tyr Ile Thr Lys Gly
440

```

&lt;210&gt; 42

&lt;211&gt; 164

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 2119917CD1

&lt;400&gt; 42

```

Met Thr Ala Ser Ala Gln Pro Arg Gly Arg Arg Pro Gly Val Gly
1 5 10 15
Val Gly Val Val Val Thr Ser Cys Lys His Pro Arg Cys Val Leu
20 25 30
Leu Gly Lys Arg Lys Gly Ser Val Gly Ala Gly Ser Phe Gln Leu
35 40 45
Pro Gly Gly His Leu Glu Phe Gly Glu Thr Trp Glu Glu Cys Ala
50 55 60
Gln Arg Glu Thr Trp Glu Glu Ala Ala Leu His Leu Lys Asn Val
65 70 75
His Phe Ala Ser Val Val Asn Ser Phe Ile Glu Lys Glu Asn Tyr
80 85 90
His Tyr Val Thr Ile Leu Met Lys Gly Glu Val Asp Val Thr His
95 100 105
Asp Ser Glu Pro Lys Asn Val Glu Pro Glu Lys Asn Glu Ser Trp
110 115 120
Glu Trp Val Pro Trp Glu Glu Leu Pro Pro Leu Asp Gln Leu Phe
125 130 135

```

WO 00/44900

PCT/US00/02237

Trp Gly Leu Arg Cys Leu Lys Glu Gln Gly Tyr Asp Pro Phe Lys  
 140 145  
 Glu Asp Leu Asn His Leu Val Gly Tyr Lys Gly Asn His Leu  
 155 160

<210> 43  
 <211> 215  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 2123456CD1

<400> 43  
 Met Leu Gly Ala Glu Trp Ser Lys Leu Gln Pro Thr Glu Lys Gln  
 1 5 10 15  
 Arg Tyr Leu Asp Glu Ala Glu Arg Glu Lys Gln Gln Tyr Met Lys  
 20 25 30  
 Glu Leu Arg Ala Tyr Gln Gln Ser Glu Ala Tyr Lys Met Cys Thr  
 35 40 45  
 Glu Lys Ile Gln Glu Lys Lys Ile Lys Lys Glu Asp Ser Ser Ser  
 50 55 60  
 Gly Leu Met Asn Thr Leu Leu Asn Gly His Lys Gly Gly Asp Cys  
 65 70 75  
 Asp Gly Phe Ser Thr Phe Asp Val Pro Ile Phe Thr Glu Glu Phe  
 80 85 90  
 Leu Asp Gln Asn Lys Ala Arg Glu Ala Glu Leu Arg Arg Leu Arg  
 95 100 105  
 Lys Met Asn Val Ala Phe Glu Glu Gln Asn Ala Val Leu Gln Arg  
 110 115 120  
 His Thr Gln Ser Met Ser Ser Ala Arg Glu Arg Leu Glu Gln Glu  
 125 130 135  
 Leu Ala Leu Glu Glu Arg Arg Thr Leu Ala Leu Gln Gln Gln Leu  
 140 145 150  
 Gln Ala Val Arg Gln Ala Leu Thr Ala Ser Phe Ala Ser Leu Pro  
 155 160 165  
 Val Pro Gly Thr Gly Glu Thr Pro Thr Leu Gly Thr Leu Asp Phe  
 170 175 180  
 Tyr Met Ala Arg Leu His Gly Ala Ile Glu Arg Asp Pro Ala Gln  
 185 190 195  
 His Glu Lys Leu Ile Val Arg Ile Lys Glu Ile Leu Ala Gln Val  
 200 205 210  
 Ala Ser Glu His Leu  
 215

<210> 44  
 <211> 539  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 2148792CD1

<400> 44



WO 00/44900

PCT/US00/02237

Met	Ala	Ala	Leu	Phe	Leu	Ser	Ala	Pro	Pro	Gln	Ala	Glu	Val	Thr	
1				5					10					15	
Phe	Glu	Asp	Val	Ala	Val	Tyr	Leu	Ser	Arg	Glu	Glu	Trp	Gly	Arg	
				20					25					30	
Leu	Gly	Pro	Ala	Gln	Arg	Gly	Leu	Tyr	Arg	Asp	Val	Met	Leu	Glu	
				35					40					45	
Thr	Tyr	Gly	Asn	Leu	Val	Ser	Leu	Gly	Val	Gly	Pro	Ala	Gly	Pro	
				50					55					60	
Lys	Pro	Gly	Val	Ile	Ser	Gln	Leu	Glu	Arg	Gly	Asp	Glu	Pro	Trp	
				65					70					75	
Val	Leu	Asp	Val	Gln	Gly	Thr	Ser	Gly	Lys	Glu	His	Leu	Arg	Val	
				80					85					90	
Asn	Ser	Pro	Ala	Leu	Gly	Thr	Arg	Thr	Glu	Tyr	Lys	Glu	Leu	Thr	
				95					100					105	
Ser	Gln	Glu	Thr	Phe	Gly	Glu	Glu	Asp	Pro	Gln	Gly	Ser	Glu	Pro	
				110					115					120	
Val	Glu	Ala	Cys	Asp	His	Ile	Ser	Lys	Ser	Glu	Gly	Ser	Leu	Glu	
				125					130					135	
Lys	Leu	Val	Glu	Gln	Arg	Gly	Pro	Arg	Ala	Val	Thr	Leu	Thr	Asn	
				140					145					150	
Gly	Glu	Ser	Ser	Arg	Glu	Ser	Gly	Gly	Asn	Leu	Arg	Leu	Leu	Ser	
				155					160					165	
Arg	Pro	Val	Pro	Asp	Gln	Arg	Pro	His	Lys	Cys	Asp	Ile	Cys	Glu	
				170					175					180	
Gln	Ser	Phe	Glu	Gln	Arg	Ser	Tyr	Leu	Asn	Asn	His	Lys	Arg	Val	
				185					190					195	
His	Arg	Ser	Lys	Lys	Thr	Asn	Thr	Val	Arg	Asn	Ser	Gly	Glu	Ile	
				200					205					210	
Phe	Ser	Ala	Asn	Leu	Val	Val	Lys	Glu	Asp	Gln	Lys	Ile	Pro	Thr	
				215					220					225	
Gly	Lys	Lys	Leu	His	Tyr	Cys	Ser	Tyr	Cys	Gly	Lys	Thr	Phe	Arg	
				230					235					240	
Tyr	Ser	Ala	Asn	Leu	Val	Lys	His	Gln	Arg	Leu	His	Thr	Glu	Glu	
				245					250					255	
Lys	Pro	Tyr	Lys	Cys	Asp	Glu	Cys	Gly	Lys	Ala	Phe	Ser	Gln	Ser	
				260					265					270	
Cys	Glu	Phe	Ile	Asn	His	Arg	Arg	Met	His	Ser	Gly	Glu	Ile	Pro	
				275					280					285	
Tyr	Arg	Cys	Asp	Glu	Cys	Gly	Lys	Thr	Phe	Thr	Arg	Arg	Pro	Asn	
				290					295					300	
Leu	Met	Lys	His	Gln	Arg	Ile	His	Thr	Gly	Glu	Lys	Pro	Tyr	Lys	
				305					310					315	
Cys	Gly	Glu	Cys	Gly	Lys	His	Phe	Ser	Ala	Tyr	Ser	Ser	Leu	Ile	
				320					325					330	
Tyr	His	Gln	Arg	Ile	His	Thr	Gly	Glu	Lys	Pro	Tyr	Lys	Cys	Asn	
				335					340					345	
Asp	Cys	Gly	Lys	Ala	Phe	Ser	Asp	Gly	Ser	Ile	Leu	Ile	Arg	His	
				350					355					360	
Arg	Arg	Thr	His	Thr	Gly	Glu	Lys	Pro	Phe	Glu	Cys	Lys	Glu	Cys	
				365					370					375	
Gly	Lys	Gly	Phe	Thr	Gln	Ser	Ser	Asn	Leu	Ile	Gln	His	Gln	Arg	
				380					385					390	
Ile	His	Thr	Gly	Glu	Lys	Pro	Tyr	Lys	Cys	Asn	Glu	Cys	Glu	Lys	
				395					400					405	
Ala	Phe	Ile	Gln	Lys	Thr	Lys	Leu	Val	Glu	His	Gln	Arg	Ser	His	
				410					415					420	
Thr	Gly	Glu	Lys	Pro	Tyr	Glu	Cys	Asn	Asp	Cys	Gly	Lys	Val	Phe	
				425					430					435	
Ser	Gln	Ser	Thr	His	Leu	Ile	Gln	His	Gln	Arg	Ile	His	Thr	Gly	
				440					445					450	
Glu	Lys	Pro	Tyr	Lys	Cys	Ser	Glu	Cys	Gly	Lys	Ala	Phe	His	Asn	
				455					460					465	

Ser	Ser	Arg	Leu	Ile	His	His	Gln	Arg	Leu	His	His	Gly	Glu	Lys
				470					475					480
Pro	Tyr	Arg	Cys	Ser	Asp	Cys	Lys	Lys	Ala	Phe	Ser	Gln	Ser	Thr
				485					490					495
Tyr	Leu	Ile	Gln	His	Arg	Arg	Ile	His	Thr	Gly	Glu	Lys	Pro	Tyr
				500					505					510
Lys	Cys	Ser	Glu	Cys	Gly	Lys	Ala	Phe	Arg	His	Ser	Ser	Asn	Met
				515					520					525
Cys	Gln	His	Gln	Arg	Ile	His	Leu	Arg	Glu	Asp	Phe	Ser	Met	
				530					535					

&lt;210&gt; 45

&lt;211&gt; 182

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 2751943CD1

&lt;400&gt; 45

Met	Ala	Arg	Leu	Leu	Trp	Leu	Leu	Arg	Gly	Leu	Thr	Leu	Gly	Thr
1				5					10					15
Ala	Pro	Arg	Arg	Ala	Val	Arg	Gly	Gln	Ala	Gly	Gly	Gly	Gly	Pro
				20					25					30
Gly	Thr	Gly	Pro	Gly	Leu	Gly	Glu	Ala	Gly	Ser	Leu	Ala	Thr	Cys
				35					40					45
Glu	Leu	Pro	Leu	Ala	Lys	Ser	Glu	Trp	Gln	Lys	Lys	Leu	Thr	Pro
				50					55					60
Glu	Gln	Phe	Tyr	Val	Thr	Arg	Glu	Lys	Gly	Thr	Glu	Pro	Pro	Phe
				65					70					75
Ser	Gly	Ile	Tyr	Leu	Asn	Asn	Lys	Glu	Ala	Gly	Met	Tyr	His	Cys
				80					85					90
Val	Cys	Cys	Asp	Ser	Pro	Leu	Phe	Ser	Ser	Glu	Lys	Lys	Tyr	Cys
				95					100					105
Ser	Gly	Thr	Gly	Trp	Pro	Ser	Phe	Ser	Glu	Ala	His	Gly	Thr	Ser
				110					115					120
Gly	Ser	Asp	Glu	Ser	His	Thr	Gly	Ile	Leu	Arg	Arg	Leu	Asp	Thr
				125					130					135
Ser	Leu	Gly	Ser	Ala	Arg	Thr	Glu	Val	Val	Cys	Lys	Gln	Cys	Glu
				140					145					150
Ala	His	Leu	Gly	His	Val	Phe	Pro	Asp	Gly	Pro	Gly	Pro	Asn	Gly
				155					160					165
Gln	Arg	Phe	Cys	Ile	Asn	Ser	Val	Ala	Leu	Lys	Phe	Lys	Pro	Arg
				170					175					180
Lys	His													

&lt;210&gt; 46

&lt;211&gt; 534

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 3128913CD1

&lt;400&gt; 46

Met	Ala	Val	Glu	Ser	Gly	Val	Ile	Ser	Thr	Leu	Ile	Pro	Gln	Asp	15
1				5					10						15
Pro	Pro	Glu	Gln	Glu	Leu	Ile	Leu	Val	Lys	Val	Glu	Asp	Asn	Phe	30
				20					25						30
Ser	Trp	Asp	Glu	Lys	Phe	Lys	Gln	Asn	Gly	Ser	Thr	Gln	Ser	Cys	45
				35					40						45
Gln	Glu	Leu	Phe	Arg	Gln	Gln	Phe	Arg	Lys	Phe	Cys	Tyr	Gln	Glu	60
				50					55						60
Thr	Pro	Gly	Pro	Arg	Glu	Ala	Leu	Ser	Arg	Leu	Gln	Glu	Leu	Cys	75
				65					70						75
Tyr	Gln	Trp	Leu	Met	Pro	Glu	Leu	His	Thr	Lys	Glu	Gln	Ile	Leu	90
				80					85						90
Glu	Leu	Leu	Val	Leu	Glu	Gln	Phe	Leu	Ser	Ile	Leu	Pro	Glu	Glu	105
				95					100						105
Leu	Gln	Ile	Trp	Val	Gln	Gln	His	Asn	Pro	Glu	Ser	Gly	Glu	Glu	120
				110					115						120
Ala	Val	Thr	Leu	Leu	Glu	Asp	Leu	Glu	Arg	Glu	Phe	Asp	Asp	Pro	135
				125					130						135
Gly	Gln	Gln	Val	Pro	Ala	Ser	Pro	Gln	Gly	Pro	Ala	Val	Pro	Trp	150
				140					145						150
Lys	Asp	Leu	Thr	Cys	Leu	Arg	Ala	Ser	Gln	Glu	Ser	Thr	Asp	Ile	165
				155					160						165
His	Leu	Gln	Pro	Leu	Lys	Thr	Gln	Leu	Lys	Ser	Trp	Lys	Pro	Cys	180
				170					175						180
Leu	Ser	Pro	Lys	Ser	Asp	Cys	Glu	Asn	Ser	Glu	Thr	Ala	Thr	Lys	195
				185					190						195
Glu	Gly	Ile	Ser	Glu	Glu	Lys	Ser	Gln	Gly	Leu	Pro	Gln	Glu	Pro	210
				200					205						210
Ser	Phe	Arg	Gly	Ile	Ser	Glu	His	Glu	Ser	Asn	Leu	Val	Trp	Lys	225
				215					220						225
Gln	Gly	Ser	Ala	Thr	Gly	Glu	Lys	Leu	Arg	Ser	Pro	Ser	Gln	Gly	240
				230					235						240
Gly	Ser	Phe	Ser	Gln	Val	Ile	Phe	Thr	Asn	Lys	Ser	Leu	Gly	Lys	255
				245					250						255
Arg	Asp	Leu	Tyr	Asp	Glu	Ala	Glu	Arg	Cys	Leu	Ile	Leu	Thr	Thr	270
				260					265						270
Asp	Ser	Ile	Met	Cys	Gln	Lys	Val	Pro	Pro	Glu	Glu	Arg	Pro	Tyr	285
				275					280						285
Arg	Cys	Asp	Val	Cys	Gly	His	Ser	Phe	Lys	Gln	His	Ser	Ser	Leu	300
				290					295						300
Thr	Gln	His	Gln	Arg	Ile	His	Thr	Gly	Glu	Lys	Pro	Tyr	Lys	Cys	315
				305					310						315
Asn	Gln	Cys	Gly	Lys	Ala	Phe	Ser	Leu	Arg	Ser	Tyr	Leu	Ile	Ile	330
				320					325						330
His	Gln	Arg	Ile	His	Ser	Gly	Glu	Lys	Ala	Tyr	Glu	Cys	Ser	Glu	345
				335					340						345
Cys	Gly	Lys	Ala	Phe	Asn	Gln	Ser	Ser	Ala	Leu	Ile	Arg	His	Arg	360
				350					355						360
Lys	Ile	His	Thr	Gly	Glu	Lys	Ala	Cys	Lys	Cys	Asn	Glu	Cys	Gly	375
				365					370						375
Lys	Ala	Phe	Ser	Gln	Ser	Ser	Tyr	Leu	Ile	Ile	His	Gln	Arg	Ile	390
				380					385						390
His	Thr	Gly	Glu	Lys	Pro	Tyr	Glu	Cys	Asn	Glu	Cys	Gly	Lys	Thr	405
				395					400						405
Phe	Ser	Gln	Ser	Ser	Lys	Leu	Ile	Arg	His	Gln	Arg	Ile	His	Thr	420
				410					415						420
Gly	Glu	Arg	Pro	Tyr	Glu	Cys	Asn	Glu	Cys	Gly	Lys	Ala	Phe	Arg	435
				425					430						435
Gln	Ser	Ser	Glu	Leu	Ile	Thr	His	Gln	Arg	Ile	His	Ser	Gly	Glu	450
				440					445						450
Lys	Pro	Tyr	Glu	Cys	Ser	Glu	Cys	Gly	Lys	Ala	Phe	Ser	Leu	Ser	

Ser	Asn	Leu	Ile	455	Arg	His	Gln	Arg	Ile	460	His	Ser	Gly	Glu	Glu	465
				470						475						480
Tyr	Gln	Cys	Asn	Glu	Cys	Gly	Lys	Thr	Phe	Lys	Arg	Ser	Ser	Ser	Ala	
				485						490						495
Leu	Val	Gln	His	Gln	Arg	Ile	His	Ser	Gly	Asp	Glu	Ala	Tyr	Ile		
				500						505						510
Cys	Asn	Glu	Cys	Gly	Lys	Ala	Phe	Arg	His	Arg	Ser	Val	Leu	Met		
				515						520						525

Arg His Gln Arg Val His Thr Ile Lys  
530

&lt;210&gt; 47

&lt;211&gt; 206

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 3282941CD1

&lt;400&gt; 47

Met	Ser	Thr	Gly	Ser	Val	Ser	Asp	Pro	Glu	Glu	Met	Glu	Leu	Arg		
1				5					10						15	
Gly	Leu	Gln	Arg	Glu	Tyr	Pro	Val	Pro	Ala	Ser	Lys	Arg	Pro	Pro		
				20					25						30	
Leu	Arg	Gly	Val	Glu	Arg	Ser	Tyr	Ala	Ser	Pro	Ser	Asp	Asn	Ser		
				35					40						45	
Ser	Ala	Glu	Glu	Glu	Asp	Pro	Asp	Gly	Glu	Glu	Glu	Arg	Cys	Ala		
				50					55						60	
Leu	Gly	Thr	Ala	Gly	Ser	Ala	Glu	Gly	Cys	Lys	Arg	Lys	Arg	Pro		
				65					70						75	
Arg	Val	Ala	Gly	Gly	Gly	Gly	Ala	Gly	Gly	Ser	Ala	Gly	Gly	Gly		
				80					85						90	
Gly	Lys	Lys	Pro	Leu	Pro	Ala	Lys	Gly	Ser	Ala	Ala	Glu	Cys	Lys		
				95					100						105	
Gln	Ser	Gln	Arg	Asn	Ala	Ala	Asn	Ala	Arg	Glu	Arg	Ala	Arg	Met		
				110					115						120	
Arg	Val	Leu	Ser	Lys	Ala	Phe	Ser	Arg	Leu	Lys	Thr	Ser	Leu	Pro		
				125					130						135	
Trp	Val	Pro	Pro	Asp	Thr	Lys	Leu	Ser	Lys	Leu	Asp	Thr	Leu	Arg		
				140					145						150	
Leu	Ala	Ser	Ser	Tyr	Ile	Ala	His	Leu	Arg	Gln	Leu	Leu	Gln	Glu		
				155					160						165	
Asp	Arg	Tyr	Glu	Asn	Gly	Tyr	Val	His	Pro	Val	Asn	Leu	Thr	Trp		
				170					175						180	
Pro	Phe	Val	Val	Ser	Gly	Arg	Pro	Asp	Ser	Asp	Thr	Lys	Glu	Val		
				185					190						195	
Ser	Ala	Ala	Asn	Arg	Leu	Cys	Gly	Thr	Thr	Ala						
				200					205							

&lt;210&gt; 48

&lt;211&gt; 172

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 3286656CD1

<400> 48  
 Met Glu Ser Val Thr Phe Glu Asp Val Ala Val Glu Phe Ile Gln  
 1 5 10 15  
 Glu Trp Ala Leu Leu Asp Ser Ala Arg Ser Leu Cys Lys Tyr  
 20 25 30  
 Arg Met Leu Asp Gln Cys Arg Thr Leu Ala Ser Arg Gly Thr Pro  
 35 40 45  
 Pro Cys Lys Pro Ser Cys Val Ser Gln Leu Gly Gln Arg Ala Glu  
 50 55 60  
 Pro Lys Ala Thr Glu Arg Gly Ile Leu Arg Ala Thr Gly Val Ala  
 65 70 75  
 Trp Glu Ser Gln Leu Lys Pro Glu Glu Leu Pro Ser Met Gln Asp  
 80 85 90  
 Leu Leu Glu Glu Ala Ser Ser Arg Asp Met Gln Met Gly Pro Gly  
 95 100 105  
 Leu Phe Leu Arg Met Gln Leu Val Pro Ser Ile Glu Glu Arg Glu  
 110 115 120  
 Thr Pro Leu Thr Arg Glu Asp Arg Pro Ala Leu Gln Glu Pro Pro  
 125 130 135  
 Trp Ser Leu Gly Cys Thr Gly Leu Lys Ala Ala Met Gln Ile Gln  
 140 145 150  
 Arg Val Val Ile Pro Val Pro Thr Leu Gly His Arg Asn Pro Trp  
 155 160 165  
 Val Ala Arg Asp Ser Ala Met  
 170

<210> 49  
 <211> 275  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 3490802CD1

<400> 49  
 Met Gly Pro Leu Gln Phe Arg Asp Val Ala Ile Glu Phe Ser Leu  
 1 5 10 15  
 Glu Glu Trp His Cys Leu Asp Thr Ala Gln Arg Asn Leu Tyr Arg  
 20 25 30  
 Asp Val Met Leu Glu Asn Tyr Arg Asn Leu Val Phe Leu Gly Ile  
 35 40 45  
 Val Val Ser Lys Pro Asp Leu Val Thr Cys Leu Glu Gln Gly Lys  
 50 55 60  
 Lys Pro Leu Thr Met Glu Arg His Glu Met Ile Ala Lys Pro Pro  
 65 70 75  
 Val Met Ser Ser His Phe Ala Gln Asp Leu Trp Pro Glu Asn Ile  
 80 85 90  
 Gln Asn Ser Phe Gln Ile Gly Met Leu Arg Arg Tyr Glu Glu Cys  
 95 100 105  
 Arg His Asp Asn Leu Gln Leu Lys Lys Gly Cys Lys Ser Val Gly  
 110 115 120  
 Glu His Lys Val His Lys Gly Gly Tyr Asn Gly Leu Asn Gln Cys  
 125 130 135  
 Leu Thr Thr Thr Gln Lys Glu Ile Phe Gln Cys Asp Lys Tyr Gly



215 220 225  
 Lys Met Glu Gln Met Ser Arg Val Ser Lys Asn  
 230 235

<210> 51  
 <211> 214  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 3573060CD1

<400> 51  
 Met Asn Leu Ser Ser Ala Ser Ser Thr Glu Glu Lys Ala Val Thr  
 1 5 10 15  
 Thr Val Leu Trp Gly Cys Glu Leu Ser Gln Glu Arg Arg Thr Trp  
 20 25 30  
 Thr Phe Arg Pro Gln Leu Glu Gly Lys Gln Ser Cys Arg Leu Leu  
 35 40 45  
 Leu His Thr Ile Cys Leu Gly Glu Lys Ala Lys Glu Glu Met His  
 50 55 60  
 Arg Val Glu Ile Leu Pro Pro Ala Asn Gln Glu Asp Lys Lys Met  
 65 70 75

Gln Pro Val Thr Ile Ala Ser Leu Gln Ala Ser Val Leu Pro Met  
 80 85 90  
 Val Ser Met Val Gly Val Gln Leu Ser Pro Pro Val Thr Phe Gln  
 95 100 105  
 Leu Arg Ala Gly Ser Gly Pro Val Phe Leu Ser Gly Gln Glu Arg  
 110 115 120  
 Tyr Glu Ala Ser Asp Leu Thr Trp Glu Glu Glu Glu Glu Glu  
 125 130 135  
 Gly Glu Glu Glu Glu Glu Glu Glu Asp Asp Glu Asp Glu Asp  
 140 145 150  
 Ala Asp Ile Ser Leu Glu Glu Gln Ser Pro Val Lys Gln Val Lys  
 155 160 165  
 Arg Leu Val Pro Gln Lys Gln Ala Ser Val Ala Lys Lys Lys Lys  
 170 175 180  
 Leu Glu Lys Glu Glu Glu Glu Ile Arg Ala Ser Val Arg Asp Lys  
 185 190 195  
 Ser Pro Val Lys Lys Ala Lys Ala Thr Ala Arg Ala Lys Lys Pro  
 200 205 210  
 Gly Phe Lys Lys

<210> 52  
 <211> 396  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 3573661CD1

<400> 52  
 Met Asn Phe Thr Val Gly Phe Lys Pro Leu Leu Gly Asp Ala His

1	5	10	15
Ser Met Asp Asn	Leu Glu Lys Gln Leu	Ile Cys Pro Ile Cys Leu	
	20	25	30
Glu Met Phe Ser	Lys Pro Val Val Ile	Leu Pro Cys Gln His Asn	
	35	40	45
Leu Cys Arg Lys	Cys Ala Asn Asp Val	Phe Gln Ala Ser Asn Pro	
	50	55	60
Leu Trp Gln Ser	Arg Gly Ser Thr Thr	Val Ser Ser Gly Gly Arg	
	65	70	75
Phe Arg Cys Pro	Ser Cys Arg His Glu	Val Val Leu Asp Arg His	
	80	85	90
Gly Val Tyr Gly	Leu Gln Arg Asn Val	Leu Val Glu Asn Ile Ile	
	95	100	105
Asp Ile Tyr Lys	Gln Glu Ser Ser Lys	Pro Leu His Ser Lys Ala	
	110	115	120
Glu Gln His Leu	Met Cys Glu Glu His	Glu Glu Lys Ile Asn	
	125	130	135
Ile Tyr Cys Leu	Ser Cys Glu Val Pro	Thr Cys Ser Leu Cys Lys	
	140	145	150
Val Phe Gly Ala	His Lys Asp Cys Glu	Val Ala Pro Leu Pro Thr	
	155	160	165
Ile Tyr Lys Arg	Gln Lys Ser Glu Leu	Ser Asp Gly Ile Ala Met	
	170	175	180
Leu Val Ala Gly	Asn Asp Arg Val Gln	Ala Val Ile Thr Gln Met	
	185	190	195
Glu Glu Val Cys	Gln Thr Ile Glu Asp	Asn Ser Arg Arg Gln Lys	
	200	205	210
Gln Leu Leu Asn	Gln Arg Phe Glu Ser	Leu Cys Ala Val Leu Glu	
	215	220	225
Glu Arg Lys Gly	Glu Leu Leu Gln Ala	Leu Ala Arg Glu Gln Glu	
	230	235	240
Glu Lys Leu Gln	Arg Val Arg Gly Leu	Ile Arg Gln Tyr Gly Asp	
	245	250	255
His Leu Glu Ala	Ser Ser Lys Leu Val	Glu Ser Ala Ile Gln Ser	
	260	265	270
Met Glu Glu Pro	Gln Met Ala Leu Tyr	Leu Gln Gln Ala Lys Glu	
	275	280	285
Leu Ile Asn Lys	Val Gly Ala Met Ser	Lys Val Glu Leu Ala Gly	
	290	295	300
Arg Pro Glu Pro	Gly Tyr Glu Ser Met	Glu Gln Phe Thr Val Arg	
	305	310	315
Val Glu His Val	Ala Glu Met Leu Arg	Thr Ile Asp Phe Gln Pro	
	320	325	330
Gly Ala Ser Gly	Gly Gly Arg Gly Gly	Gly Pro Arg Arg Lys Lys	
	335	340	345
Arg Ala Thr Arg	Gly Pro Glu Glu Lys	Thr Ala Arg Met Gly Pro	
	350	355	360
Tyr Arg Pro Leu	Arg Pro Asn Pro Asp	Pro Leu Leu Arg Lys Ser	
	365	370	375
Pro Arg Arg Leu	Arg Ile Ser Gly Gly	Arg Asn Ser Cys Arg Lys	
	380	385	390
Lys Thr Pro Ala	Ser Phe		
	395		

<210> 53  
 <211> 486  
 <212> PRT  
 <213> Homo sapiens



<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 3633422CD1

<400> 53

Met Arg Arg Leu Val	His Asp Leu Leu Pro	Pro Glu Val Cys Ser	15
1	5	10	25
Leu Leu Asn Pro Ala	Ala Ile Tyr Ala Asn	Asn Glu Ile Ser Leu	30
20	25	30	45
Arg Asp Val Glu Val	Tyr Gly Phe Asp Tyr	Asp Tyr Thr Leu Ala	50
35	40	45	60
Gln Tyr Ala Asp Ala	Leu His Pro Glu Ile	Phe Ser Thr Ala Arg	65
50	55	60	75
Asp Ile Leu Ile Glu	His Tyr Lys Tyr Pro	Glu Gly Ile Arg Lys	80
65	70	75	90
Tyr Asp Tyr Asn Pro	Ser Phe Ala Ile Arg	Gly Leu His Tyr Asp	95
80	85	90	105
Ile Gln Lys Ser Leu	Leu Met Lys Ile Asp	Ala Phe His Tyr Val	110
95	100	105	120
Gln Leu Gly Thr Ala	Tyr Arg Gly Leu Gln	Pro Val Pro Asp Glu	125
110	115	120	135
Glu Val Ile Glu Leu	Tyr Gly Gly Thr Gln	His Ile Pro Leu Tyr	140
125	130	135	150
Gln Met Ser Gly Phe	Tyr Gly Lys Gly Pro	Ser Ile Lys Gln Phe	155
140	145	150	165
Met Asp Ile Phe Ser	Leu Pro Glu Met Ala	Leu Leu Ser Cys Val	170
155	160	165	180
Val Asp Tyr Phe Leu	Gly His Ser Leu Glu	Phe Asp Gln Ala His	185
170	175	180	195
Leu Tyr Lys Asp Val	Thr Asp Ala Ile Arg	Asp Val His Val Lys	200
185	190	195	210
Gly Leu Met Tyr Gln	Trp Ile Glu Gln Asp	Met Glu Lys Tyr Ile	215
200	205	210	225
Leu Arg Gly Asp Glu	Thr Phe Ala Val Leu	Ser Arg Leu Val Ala	230
215	220	225	240
His Gly Lys Gln Leu	Phe Leu Ile Thr Asn	Ser Pro Phe Ser Phe	245
230	235	240	255
Val Asp Lys Gly Met	Arg His Met Val Gly	Pro Asp Trp Arg Gln	260
245	250	255	270
Leu Phe Asp Val Val	Ile Val Gln Ala Asp	Lys Pro Ser Phe Phe	275
260	265	270	285
Thr Asp Arg Arg Lys	Pro Phe Arg Lys Leu	Asp Glu Lys Gly Ser	290
275	280	285	300
Leu Gln Trp Asp Arg	Ile Thr Arg Leu Glu	Lys Gly Lys Ile Tyr	305
290	295	300	315
Arg Gln Gly Asn Leu	Phe Asp Phe Leu Arg	Leu Thr Glu Trp Arg	320
305	310	315	330
Gly Pro Arg Val Leu	Tyr Phe Gly Asp His	Leu Tyr Ser Asp Leu	335
320	325	330	345
Ala Asp Leu Met Leu	Arg His Gly Trp Arg	Thr Gly Ala Ile Ile	350
335	340	345	360
Pro Glu Leu Glu Arg	Glu Ile Arg Ile Ile	Asn Thr Glu Gln Tyr	365
350	355	360	375
Met His Ser Leu Thr	Trp Gln Gln Ala Leu	Thr Gly Leu Leu Glu	380
365	370	375	390
Arg Met Gln Thr Tyr	Gln Asp Ala Glu Ser	Arg Gln Val Leu Ala	395
380	385	390	405
Ala Trp Met Lys Glu	Arg Gln Glu Leu Arg	Cys Ile Thr Lys Ala	410
395	400	405	420
Leu Phe Asn Ala Gln	Phe Gly Ser Ile Phe	Arg Thr Phe His Asn	415
410	415	420	430
Pro Thr Tyr Phe Ser	Arg Arg Leu Val Arg	Phe Ser Asp Leu Tyr	

WO 00/44900

PCT/US00/02237

Met Ala Ser Leu	425	430	435
Met Ser Cys Leu Leu Asn Tyr Arg Val Asp Phe Thr			
440	445	450	
Phe Tyr Pro Arg Arg Thr Pro Leu Gln His Glu Ala Pro Leu Trp			
455	460	465	
Met Asp Gln Leu Cys Thr Gly Cys Met Lys Thr Pro Phe Leu Gly			
470	475	480	
Asp Met Ala His Ile Arg			
485			

<210> 54  
 <211> 555  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 3993377CD1

<400> 54

Met Gly Ala Glu Asp Lys Leu Pro Leu Glu Asp Ser Pro Val Ile	
1	5
Ala Ala Leu Asp Cys Pro Ser Leu Asn Asn Ala Thr Ala Phe Ser	10
20	25
Leu Leu Ala Asp Asp Ser Gln Thr Ser Thr Ser Ile Phe Ala Ser	30
35	40
Pro Thr Ser Pro Pro Val Leu Gly Glu Ser Val Leu Gln Asp Asn	45
50	55
Ser Phe Asp Leu Asn Asn Gly Ser Asp Ala Glu Gln Glu Glu Met	60
65	70
Glu Thr Gln Ser Ser Asp Phe Pro Pro Ser Leu Thr Gln Pro Ala	75
80	85
Pro Asp Gln Ser Ser Thr Ile Gln Leu His Pro Ala Thr Ser Pro	90
95	100
Ala Val Ser Pro Thr Thr Ser Pro Ala Val Ser Leu Val Val Ser	105
110	115
Pro Ala Ala Ser Pro Glu Ile Ser Pro Glu Val Cys Pro Ala Ala	120
125	130
Ser Thr Val Val Ser Pro Ala Val Phe Ser Val Val Ser Pro Ala	135
140	145
Ser Ser Ala Val Leu Pro Ala Val Ser Leu Glu Val Pro Leu Thr	150
155	160
Ala Ser Val Thr Ser Pro Lys Ala Ser Pro Val Thr Ser Pro Ala	165
170	175
Ala Ala Phe Pro Thr Ala Ser Pro Ala Asn Lys Asp Val Ser Ser	180
185	190
Phe Leu Glu Thr Thr Ala Asp Val Glu Glu Ile Thr Gly Glu Gly	195
200	205
Leu Thr Ala Ser Gly Ser Gly Asp Val Met Arg Arg Arg Ile Ala	210
215	220
Thr Pro Glu Glu Val Arg Leu Pro Leu Gln His Gly Trp Arg Arg	225
230	235
Glu Val Arg Ile Lys Asn Ser Ser His Arg Trp Gln Gly Glu Thr	240
245	250
Trp Tyr Tyr Gly Pro Cys Gly Lys Arg Met Lys Gln Phe Pro Glu	255
260	265
Val Ile Lys Tyr Leu Ser Arg Asn Val Val His Ser Val Arg Arg	270
275	280
Glu His Phe Ser Phe Ser Pro Arg Met Pro Val Gly Asp Phe Phe	285

Glu Glu Arg Asp	290	Pro Glu Gly Leu	295	Trp Val Gln Leu	300
Thr	305	Gln	310	Ile Thr Gly Lys	315
Ala Glu Glu Ile	320	Pro Ser Arg Ile	325	Ala Ile Thr Gly Lys	330
Arg	335	Asn Thr Glu Lys	340	Ala Thr Lys Glu Val	345
Gly Arg Pro Arg	350	Thr Arg Gly Arg	355	Pro Lys Val Lys Ile	360
Lys Val Lys Arg	365	Gly Thr Asp Asn Arg	370	Pro Leu Lys Lys Leu	375
Glu Leu Leu Asn	380	Lys Asn Glu Glu Asp	385	Lys Ala Lys Ile Ala	390
Thr	395	Met Arg Gln Lys Val	400	Gln Arg Gly Glu Cys	405
Ser Lys Lys Lys	410	Gln Ala Arg Asn	415	Lys Arg Lys Gln Glu	420
Thr Thr Ile Gln	425	Gly Lys Glu Ala Lys	430	Lys Lys Ser Lys Ala	435
Lys Ser Leu Lys	440	Gln Thr Lys Gln Glu	445	Lys Leu Lys Glu Lys	450
Gly	455	Lys Glu Lys Val Lys	460	Met Lys Glu Lys Glu	465
Lys Arg Glu Lys	470	Lys Pro Ala Cys Lys	475	Ala Asp Lys Thr Leu	480
Val Thr Lys Ala	485	Glu Glu Arg Gln	490	Gln Gln Met Ile Leu	495
Thr Gln Arg Arg	500	Leu Thr Glu Asp	505	Met Cys Leu Thr Asp	510
Glu Asp Met Lys	515	Lys Phe Ser Arg Val	520	Pro Gly Leu Thr Leu	525
Gln Pro Leu Pro	530	Asp Asp Cys Leu Thr	535	Ile Val Glu Phe Leu	540
Ser Gly Ala Phe	545	Val Leu Gly Leu Asp	550	Pro Ala Gln Gly Cys	555
Ser Phe Gly Lys					

&lt;210&gt; 55

&lt;211&gt; 61

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 4717936CD1

&lt;400&gt; 55

Met Gln Pro Arg Thr	Gln Pro Leu Ala Gln Thr Leu Pro Phe	Phe
1	5	10
Leu Gly Gly Ala Pro	Arg Asp Thr Gly Leu Arg Val Pro Val	Ile
20	25	30

Lys Met Gly Thr Gly	Trp Glu Gly Phe Gln Arg Thr Leu Lys	Glu
35	40	45
Val Ala Tyr Ile Leu	Leu Cys Cys Trp Cys Ile Lys Glu Leu	Leu
50	55	60

Asp

WO 00/44900

PCT/US00/02237

<210> 56  
<211> 2781  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc-feature  
<223> Incyte ID No.: 025733CBI

<400> 56  
gaagtggggt gcacgcttcg ggttggtgtc atggcagctg cggggagccg caagaggcgc 60  
ctggcgggagc tgacgggtgga cgagttccta gcttcgggct ttgactccga gtcccgaatcc 120  
gagtcgccaaa attctccaca agcggagaca cgggaagcac gcgaggctgc cgggagctccg 180  
gataagccggg gcgggagccc ctcgccagc cggcgtaaa ggcgtgacct tgagcacaaa 240  
gaccagctctc ctcggtctgaa ggacagagac ccaggtttct acaagttcct gcaggagaaat 300  
gaccacagac tgctaaactt cagcgactcg gacagctctg aggaggaaga ggggccgttct 360  
cactccctgc cagatgtgct ggaggaagcc agtgaggagg aggatggagc ggaggaagga 420  
gaagatgggg acagagtcct cagagggctg aaggggaaga agaattctgt tctctgacc 480  
gtcgccatgg ttgagagatg gaagcaggca gcaaaacaa cccctactcc aaagctgttct 540  
catgaagtgg tacaggcgct ccgagcagct gtggccacca cccgagggga ccaggaaaagt 600  
gctgaggcca acaaatctca ggtcacggac agtgctgcat tcaattgctct gggttacctt 660  
tgcacacagag acctcattgg ctgtctccag aagctgtctgt ttggaaaagt ggcaaaaggt 720  
agcagcaggga tgcctgcagcc gtcacagcag ccgctctggg ggaagctctg tgctgggcatc 780  
aaggctctacc tgggctcggc catacagctg gtgtctctgtc ttctggagac gacgagctgtg 840  
cgcccgctgc tggcgcaaat cagcgtgctg gtgcccctgtc tcttgacctt ccccaagcag 900  
tgccgcatgc tgcctcagag aatgggtggc atgcacagct gtgaggaact agctgtctgac 960  
gtgctggctt tctgtgtcct agcagagtc ccatcacgtat accttgaagg tgcgcataca 1020  
ccgctctctca agcaaatgta catcagctat cctcctctac atcccgccagc acaactggga 1080  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 1140  
gggtgtggcct accagcacgc ggtcctgagc actcggggac agctctctcc 1200  
gccatgacca cccgcaagaa ggcctctgagc actcggggac agctctctcc 1260  
ttctaccgcc tegtgtgccc ggtcctgagc actcggggac agctctctcc 1320  
ttctaccgcc tegtgtgccc ggtcctgagc actcggggac agctctctcc 1380  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 1440  
gggtgtggcct accagcacgc ggtcctgagc actcggggac agctctctcc 1500  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 1560  
gggtgtggcct accagcacgc ggtcctgagc actcggggac agctctctcc 1620  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 1680  
gggtgtggcct accagcacgc ggtcctgagc actcggggac agctctctcc 1740  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 1800  
gggtgtggcct accagcacgc ggtcctgagc actcggggac agctctctcc 1860  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 1920  
gggtgtggcct accagcacgc ggtcctgagc actcggggac agctctctcc 2040  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 2100  
gggtgtggcct accagcacgc ggtcctgagc actcggggac agctctctcc 2160  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 2220  
gggtgtggcct accagcacgc ggtcctgagc actcggggac agctctctcc 2280  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 2340  
gggtgtggcct accagcacgc ggtcctgagc actcggggac agctctctcc 2400  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 2460  
gggtgtggcct accagcacgc ggtcctgagc actcggggac agctctctcc 2520  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 2580  
gggtgtggcct accagcacgc ggtcctgagc actcggggac agctctctcc 2640  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 2700  
gggtgtggcct accagcacgc ggtcctgagc actcggggac agctctctcc 2760  
gcccctccctc tcatcagttt accagcacgc ggaacacatac cagctctgtgt 2781

<210> 57  
<211> 2544  
<212> DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 079702CB1

&lt;400&gt; 57

```

cgggaaacac aaatggcgag gggctgtatt gaagtgggct gtgtttggag cgggtgtaag 60
aacgctcatt ctaccccacaa cctctgtctc caaggacctc ggtttgtgag tgcattatgtg 120
cgggttaccc ggtggggcgg gtgccacgta agtgctcgga ctgcaggggg aagcggccac 180
gggggacggat tggttgtttt ttctgtatg aagcgggttg caccactgaa gtgacccgaat 240
gagagactct acaggggcag gtaattcact ggtccacaag cggctctcct tacgtcgaaa 300
ccaaaagacc ccaacatcct tgaccaagct gtctttacag gatggacata aagccaaaaa 360
gccagcatgt aaatttgaag agggtcagga tgtctagct agatgtctcag atggcttgtt 420
ttatcttggc actatcaaaa agataaacat attgaaacag agctgcttca tcattattga 480
agacagttct aaactctggg ttctctggaa ggacattcaa acaggagcca ctgggaagtgg 540
ggaaatggctc tgtacaatat gtcaagaaga gtaattcaga gctcccaatg aaatggttat 600
atgtgacaa gtgtggcgaag gatattcatca gttgtgtcac acacctcata ttgattccag 660
tgtgatgtat tcagatgaaa aatggctctg tcggcagttg gtttttgcaa caacaacaaa 720
gaggggtgggt gcaacttaaga aaggaccaaa tgccaagca ttgcaagtca tgaagcagac 780
attaccctat agtgtggcag accttgaatg ggtgacaggt cataaaacca atgtccagca 840
gtgttactgc tatttggag gccctggaga ctggtatttt caaatgctac agtgcctcaa 900
atgtaagcag tggtttcagt aggccttgtgt gcaatgcctt caaagccaa tgctatttgg 960
agacagattt tatagttta tatgctctgt ctgcagttct ggaccagaa accccaacg 1020
tctaccatta cagtgggtag atatagcaca cctatgcctt tacaacctaa gtgtatttca 1080
taagaagaaa tactttgatt ctgaacttga gcttatatgc tacattaatg aaacttggga 1140
tgatttgcaa cctggagagc tggcagacac accaaaaatg gaaagatatg agcatgttct 1200
gaagggcata aatgattaca agaccatggt tatgtctggg taatgtctgg aaagaaataa aaggaagtaa 1260
gcatttggtt ggggttgcgaa ttctgtgtcc tctgtgtcca ccaaatgtgg ctcttcaaac 1320
agaaaagaaa cctgaaggaa catctcatga attttaaatt ttctgtgcca aaagggcaga aggcattcaa 1380
acctatctct gattcaaggg aagtaagcaa tggcatagaa aaaaaaaaaa aaaaaaaatc 1440
tgtaggctgt ccacctggcc catatacaag aaaaatgatt aaaaaaaact ctgagccact 1500
tttgataagg gaatcaattt cagagaatcc tactttggat ttaccttgtt ctatagggag 1560
aactgaggga actgcacatt catccaatcc ctgagattgt gatttccagg gtgcttccag 1620
tgcaaaagaa actacctcgt ctagcatctt caggcattat ggatttatct actccagaaa 1680
aagaacgggt acaggaagat ctggcctgct tgcaatacca catttgcgga atgttgcgga gaagaagagg 1740
tcgtcttcca agaagagcac tccagactca gaaactcaga attgtaaaaa atgatgaagg 1800
caaaagaagt tatcagt ttg atgaactcaa cacagagatt ctgaataact tagcagatca 1860
ggagtacaaa ctcaatcatt taaagaactc cattaccagt tatttttggt ctgcaggtag 1920
aatagcatgt ggcgaaaaat accgagtttt ggcacgtcgg gtgacacttg atggaagagt 1980
gcagtatctt gttgaaatggg aaggagcaac tgcattctga ctgtaggact gaacattatg 2040
ttcaactcac tctgattttt ttgaggtaca gttcaaaagg cttaaggagt cgggttttta 2100
ctatcttctc taaaaaaaaa aaaaagtcaa aaaaattcaa aaaaaggggt gatactagcc 2160
ttaacatgta cctgtcaatg ttatggatat tgcataaaaa aggtactctt taataatcag 2220
aacagagact taatttttta aatettaaga ttgttagaat gtttctcaga taggatatta 2280
aaaaatgatg aaacccatgc atggtgttag acaatttttc taattattcc atgtagttag 2340
ttttttgtga tctgtgatta tcagagcaaa catcatgtag atagcacaag tatttggaga 2400
aacgttgtgt gttctgttac caaaatgtgt gaaaaattta ttccaatccc tttttagatt 2460
cataaagtg agtgcataat atgcctactg aaagactgta aatatatgaa attttcttct 2520
aagcaagtg taaataata tacc 2544

```

&lt;210&gt; 58

&lt;211&gt; 1627

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 116208CB1

&lt;400&gt; 58

```

ctgatgaggg cgctgcattt attgaagagc ggcgcagcc ctgcggtcca gattaaaatc 60
cgagaattgt atagagcccg atatccacga actcttgaag gactttctga tttatccaca 120
atcaaatcat cggttttcag tttggatgggt ggcctcatcac ctgtagaacc tgactttggcc 180
gtggctggaa tccacttcgtt gccctccact tccactcacc atccctctct 240
gttggtttctg tgcctgtcca agatactaag ccacattttg agatgcagca gccatctccc 300
ccaattctctc ctgtccatcc tgatgtgcag ttataaaaatc tgccttttta tgatgtcctt 360
gatgtttctca tcaagccccc gagttttagtt caaagcagta ttcagcgatt tcaagagaa 420
ttttttatttt ttgctttgac acctcaacaa gtttagagaga atcagttcgt tgagactttt cagggtattt 480
tggccaggtg tagggagaga ttatacagtc caagttcagt tgagactttt ctgggcagag 540
acaagttgcc ctcaagaaga taactatcca aatagttctat gtataaaaagt aaatgggaa 600
ctatttctct tgcctggcta tgcaccaccg cctaaaaaatg ggtatgaaca gaagcgcct 660
ggagcgcctt tgaatat tac atcttttagtt aggtttatct tgcctgtgcc aaaccctgag 720
tccatttctt gggcatcaga aattgggaag aattactcta tgtctgtata tcttctaagg 780
cagctttacat cagccatggt attacagaga ttataaaatga aaaccctgat 840
cattccagag cactaagtaa agaaaaactt actgcagatc ctgatagtga aattgtcata 900
actagccttc gggtaacctt gatgtgccct ttaggaaaaa tgaggctgac aatcccatgc 960
cgtgcagtga ctgttacaca tctgcagttg ttgtacataa cctctctatc acaaatgaat 1020
gagaaaaagc ccacctggat ttgtcctgtg tgtgacaaaa agctgtcata tgaagtcta 1080
atatagatgt ggtcttttat ggaaattctc aatgactgtt ctgatgtaga tgagatcaaa 1140
ttccaaagaag atggttcttg tgtcccaatg agaccgaaga aagaagctat gaaagtatcc 1200
agccaaccgt gtacaaaaat agaaagtcca agcgtctcca gtaaagcctt ttcagtgact 1260
gtagccagtg aggcaagcaa gaagaagaat gatgttat tg atcttacaat tccagctct 1320
tctgacgaag aggaagaccc tcttgccaaa aggaaatgca tctttatgtc agaaacacaa 1380
agcagcccaa ccaaggggtt tctcatgtat cagccatctg atcactcagt accattccac 1440
acctcggttg atctctctgc tattccgcct tcatcagat ttgcaggag aactatgat 1500
catacgccaa tatcaagcat gtcactagat gactactgtc aacaaagaa acaaatgata 1560
aataatgaac tgaagcttgg aacatctctt gatactgtgc aacagtgaa acaaaaataa 1620
accgata

```

&lt;210&gt; 59

&lt;211&gt; 1043

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 179261CB1

&lt;400&gt; 59

```

gtgggatatt tcagtcaaat gaaaatcacc tctgaaaatg tgcccagtta caaaactcat 60
gaatctctta ctttaccctg gagaatccat gacagtgaga agccctatga atacaaaggaa 120
tatgagaagg tcttcagttg tgacttagag tttgatgaat atcagagaaat acatcatgg 180
ggaaaaaact atgaatgtaa tcaatgttgg taaacctttg atcagatata cccagtatg 240
ttacaactga atatttcata tgggtgtgaaa cctgttaaat tatatggaata tgggaaataa 300
tgtagttttt ataaagacct taatgtatac cagaaaaatc ataatgagaa gtctctataa 360
tgtaaagaa acagaaggac ctttgaaaga gttggaaaaa ttaactccac tcaaaagatt 420
catgatgggt tgaatgcctc ttctgtggga gatgtgaaaa aatcctttag agtgcattga 480
caacttactc gacatcagaa aatcccatat gatgagaaaa cttaacaatt tatggaa 540
ggcaaggact tccagatttc ttccacagct accgaacatc agagaattca tctcattgag 600
aaacccatca aatgtatgca ctgtgagaag gttcttagaa ttagtccaca gctcattgaa 660
catcagaa ttccactctg ttgagaaact tatgtcatga aggaattctg gaaggtcttt 720
ggagtatgta gagaacttgc tgcctcatcag agaattcata ctggaaaaat ctgtggatgg 780
atttcaatagg taatcagtat aattcagtat cctcctctct taaagctctg ttttagactt 840
catggctcatt ctgtatgtag acgtagaatt gcttagtcct agctgtatata tactatcagc 900
ttgtgcaaat gtgtgcaaat ttttagttaa aaagtttttt ctcatgattt 960
ctcatctctg catgtttttg ttacaatagc ttttgatggt tgtattatgt ctcttttgac 1020
taatacagttt tttatctctg gat

```

&lt;210&gt; 60

&lt;211&gt; 2448



```

<400> 61
ttccgggttct gtaccccccat cctttctctc gcccctccta cccgcagctc ctggcgctcg 60
gccggcgctaa ctgcagcggg gagatctcgg ccgccaaagct ccgcctcccc ccccgggctg 120
tgcggcgctaa ctgccttgag gccgaccacc ctctagccgc ttgtctcgag 180
gccccaccttc ttcccaaccc ccgcaaaact cagtaggctc gctctcgctg actcccccgc 240
cccgctgcaa ctgcaagggg cccgcccata gccagttccg gggcggttgc tcacatcgac 300
cgggaactccc cgcctccctc ccgcgccctt gggggccgtag gaggcccgag cgaggaggta 360
gggggtggcg ggggtgcact aggggtgtcc tagagaaacg ggactctgag ggcgggacat 420
ttgggcgacg cccggcgcgg gccagccatt aaacagtcct acttctgtgc cagacactga 480
actgggctct tgacggcgaa catctcttaa tctcagaac atcccaggga gctccacagg 540
atccccatat cctggggccat gagttagttg aaagactgcc ccttgacagt ccacgacttc 600
aagtctgtgg atcacctgaa ggtctgtccc cgctacacgg cagtgtcggc acgctctgag 660
gatgatggca tcggcatcga ggagctggac accctgcagc tggagctgga gacctctgct 720
tctctcgcca gccggcgccct gcgtgtgctt gaggccgaaa ccagatcct caccgactgg 780
caggataaaga aaggtgacag acgattcctg aagctgggtc gagaccatga acttggagct 840
cccccaaac atgggaagcc caagaagcag aaactggaa ggaaggcagg acatggggccg 900
ggccctggcc caggacggcc caaatccaaa aacttcagc ccaagatcca ggaatatgaa 960
ttcactgatg accctatcga cgtgccacgg atccccaaaa atgatgcctc caacagggtc 1020
tggtgcttca tgagagcccta ctgtctgtac atcaccagcg agggaggtccc cacacttgaag 1080
gagttaactga agcccccaga agatgaggct gaggattaca agatccacc ctgggggag 1140
cactactccc agcgtctgggc ccaggaggac ctgctggagg agcagaagga tggggcccg 1200
cgacggctgt ttgctgacaa gaagaaggc ctcatggggc cactgaaccg actggacact 1260
aaagatgttg atgctctgct gaagaagctc gaggcccgag atgaacagcc ggaagatgga 1320
tgcccccttg gtgccttgaa gcagcgcctc ctgcaggccc tggttggagg aaatattatt 1380
tccccctatg aggatattctc tatctctgac atgtctggga aagaatcagg ttgctgacgg 1440
gcaagacact cccctcgcaa tcagaacaag cccctcagtg tgcgcatac taagtccctg 1500
gagagccgca tcaaggaggga gctaattgcc cagggccttt tggagcttga ggaccgccc 1560
cgagaggact ccgagatgga ggtccttgct cagctctgca aacggcaggc tgagctgaag 1620
gcacttagtg ccacaaacgc caccagaag cagcactctg tgaggtctgc aaagaggagg 1680
gtgagccggc aggagctgag gcagcgggtg ccagcggagt acaacagggt catggagcc 1740
tttcgcaaga tcatggtctg ccggcgagaag aagcggactc ccaccaagaa agaaaggagc 1800
caggctgaga agactctgaa ggagcgtgag agcatcctga agtctctgga tgggtatgcc 1860
tcaccctctg ctacaggctga ttatctggcc taggggaggg gaagggaggc cactctctct 1920
ctttgggcac aggaacaact ggccctgggc tgtcccttgg cctctagagg 1980
gccgtggccc ttccccctgag gtcttttggc ctatgctctg cttagctctg 2040
cctgtctggc tagcctgagg cctagctctc gcttggctcc gcttggctcc 2100
cttctgtttc ggggtcttct ctccccctct ttaccatccc ccactcccta atccccatcc 2160
cctgtctccc ctcaaggac ttctcccttg tggttttgta aagtgcaaac ttaagaataa 2220
agtgaactgt gtggtttttc aaaaaaaaaa aaaaaa 2255

```

```

<210> 62
<211> 2982
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 491271CB1

```

```

<400> 62
gcgtggggat gtttaccgcc gttttaccgg gatagagact ccactgtgct gacagatccc 60
ttttattcac cgccttcgaa ttgtcaaaag ggaggaaagga gggacttctt ggcttctccc 120
agcatagccc cagtttatgcc atctcagaaa tatgaccttc ccagagaaga cgaggagaaa 180
atgacaaagt ttccaggaggc tgtgacattc aagggtgtgg cctgtggtct ctcaggaggag 240
gaactctgac ttccagatct taaccagagg aagctgtacc gagatgtcat gtcggagaaac 300
ttcaagaacc tggttgcagt ggggcatctt ccttcccaac cagatatggt atcccaattg 360
gaagcgaaac aaaagctctg gatgatggaa acagaaaacc aaagaagcag caagcatcaa 420
aataaagtgg aaacactcca aaaaatttga ttaaaaaacc tttaaaatca agagctgtcc 480
tgctggacaaa ttgggaacaa ggttgcaagt ggtttgcaac aaaggaagag ggtgttcc 540
tcccagttat tacaaggctga ctctattcag gtttttgaaa atgagaacaa tataatgaac 600
cctaaggagg atagccctat ttatattgaa aatacaagag ttccatcttg gagaacccag 660

```



WO 00/44900

PCT/US00/02237

```

cattcttgcg ggaatacata tctgagtgag tcacagattc agagtagagg taagcaaat 720
gagtgtgaaa ataacctgca aatcagtgaa gacttcgtga agaatacacc attcatgag 780
catatataaaa ctgacacagaa accaaaaccc tgcagaaggtg atgaataatgg caaaatcatt 840
agtgatggctt ccaaccagaa attaccctta ggagagaaaac cccatccatg ttggtgagtg 900
ggaaggggctc tcagttatag cccaagggctt ccccttcctc cgaatgttca ccaggagaaa 960
aaatgccttca gtcaaaagctt acatctgcga gaattcacc aggaagagaaa 1020
ctcaaatagat gtcatgaatc tgggtgatgg tcacaataaga gctcttttca ttcttatcaa 1080
tctaatcatac caggagagaaa gtcttataga tgcgcacagt tgcggcaggg attcagtgagc 1140
agcagcgggtc ttatcattca ttacagaact catactggag agaaacccta taagtgcgag 1200
gaatgtggta aatgtctttag tcaaaagtcca aattttcagt gccatcagag agtccacact 1260
gaagaaaaac catcaaaagt cgaagagtggt ggttaagggct tcggttgagg tgtaaatctc 1320
cgtgttcacc agaggggtcca caggggtgag aagccctata aatgtgaggg atgtggtaag 1380
ggcttcactc aggcctgcac ttttcacatc catcagagag tccacactgg agagaaaacc 1440
tacaagtgtg atgtgtgtgg taaggggcttc agccacaatt caccattaat atgacctgag 1500
agagtcacca caggagagaaa gccatacaag tgtgaggcgt gtgggaaagg cttaccctgt 1560
aatacagatc tgcatttcca tttcagagtt cacacgggag agaaacccta taaatgtaag 1620
gagtggtgta agggcttcag tcaggcttca aatcttcaag tccatcagaa tctccacact 1680
ggggagagaaa gatccaagtg tgaacgtgtg gggaagggct tcagtcatc ctcaagagt 1740
caaacaccatc agcaggttcca cactggagag aaaccatata gatgtgatgt gtgtggtaag 1800
gacttcagtt atagtctcaa tcttaactca caccagtaaa ttcacactgg agaaaaacca 1860
tataaatgtg aggaatgtgg gaagggcttc agttggagat caaatcttca tgcacatcaa 1920
agagttcact caggagaaaa accctataaa tgtgagcagt gtgataagg agtccagctc 1980
gccatagatt ttccgggtaca tcagagagtc catactggag agaagccata caaatgtggt 2040
gtctgtggta agggcttcag tcagtctctt ggtcttcaat cccatcagag agtccacacg 2100
ggggaaaaagc catcaaaagt tgaatgtgtg ggaagggctt ttgagacag ttccagtttt 2160
atatacactc agagaggcca cactggagaa aaaccttaca aatgtggaaga gtgtgggaaa 2220
ggctttgggca ggagcttgaa tcttcgcatc catcagaggg tccacacggg agagaaaacc 2280
catatagtg aggagtggtg taaggccttc agtctccctc caaatcttcc agtccagctc 2340
ggtgttcaca ccagggaaaa actctttaa tgtgaagagt gtggttaaagg ctctcagtcag 2400
agtgacagtc ttgaagccca tcagagagtc cacactggag aaaaaccata caaatgtgac 2460
gtatgtgata aggaactccg tcaccgttca cgtcttcatc atcatcagaa agtccatcat 2520
gttaaaaaagc tttagaaaag agaaaatgtg taacaaatct ttgtctgaatg cactcttcca 2580
agtttttcgc tagtccatgc tgggtgtaaa cctctgtaaa ctactgagag tggaaagggg 2640
ttgtgtcaca cttggaatct ttctaacaaa tccatacaaga tgataacaga gaacctatga 2700
caggaataga actcgtattt aggggagaaa tagggctggt ggctctcttg gtaagatcta 2760
gttaatatata atgatcattt ttcattgtga atatactctt gaagataatg gtatgagaga 2820
tatttgccat atgctaactg gttttttggc cagggagagt ttggggttat tatccctttt 2880
ctttaatttt cattttatc ttacagtgat catatttttc ataaaagctg taaagctatg 2940
aaaaatgaat aaaattacta aaaattttct gtaaaaaaaa aa 2982

```

```

<210> 63
<211> 1185
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 585172CB1

```

```

<400> 63
ggagcgtcca gagtctcggc cctgagcggg aatcgcagtg gccgagcgtg agcggcaggt 60
agaaggggct tctcgggggc ttacagggga acacaggggc ttccggcccaa ccacaagtat 120
gcagttgcac atgcccactt ttgcccatt ctttgcaaat acccaacggcg 180
ctcgcgctgc ggcctttcta atcttcgctt gtcccttcac ttccacagct ggaggtcgaa 240
ttcacacagt cagtagagaa aaagggggcg tgtctcgggt ctcccctgtt ggcataagg 300
cgggcctccc ataggagctg gtgtggagct cggggcgttg ggttagggcg gatcggggga 360
tcgcgggggt ggtgttgaca ttgtgctctc accagggcga tcgcccgcac cctcactcct 420
ggcgctetgag tctctggcgt agccatgctg agtggggcgc tggctcctggg tctggtctcc 480
atggctgcgc gctcttgctt gtccagggc agcgcgggat ccggggccat cgttccgggtg 540
gagggcgcca ttgcacgaa gtggaggag gccctgagcc cggaggtgct agagcttcgc 600
aacgagagcg gtggccacgc ggtcccgcct ggcagtgaga ctcacttcg cgtggctgtg 660

```

```

gtgagctctc gtttcgaggg actgagcccc ctacaacgac accggctggt ccacgcagcg 720
ctggccgaggg agctggggagg tccggtccat gcgctggcca tccaggcaag gacccccgcc 780
cagtcggagag agaactctca gctggacact agccccccat gccctgggtgg gaacaagaaa 840
actctaggaa cccccggaac cccaagagag ggaggaccag gatccgaatg ggctgggtga 900
gcacgaatta ccgagggcctt ccccttgata cagtcagagg tttgtaaggg atgaagaccc 960
ctggggcccca ttctgttggg gtccatacat actctccgaa gatagcaact tgcctcaggt 1020
caaaagtgaa ccgagaaaag agaagaaatca ctactactg ctcttgcctt ggactattca 1080
ggaaggcgag ccgggatgtt ccatgtttaa tctgtacaga attgcaccag acctgatgag 1140
ttggaacaaa tctatatacat taaaagaaat tacactaaaa aaaaa 1185

```

<210> 64  
 <211> 1191  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 615200CB1

```

<400> 64
tggacggagc cgtgggcttg atttctgatt tatgactgct ttttcttgta cccaatagt 60
cgcttaagaa aggtgattat tttgagagcg ctggggagac acacatgctc attctcgagg 120
gtggcggtgg tgcagaggcg agagccatgc tctgtcttgc tctcttgaga ttggttgcta 180
tctgtttctt ttgctgctgt gtttttttct gtccagtatga aaggtggaag aaggttccata 240
ttcttttctg tgggtgtctc aagtgttgtt ggaagtggag gcagcagtga caaggggaag 300
ctttccctgc aggatgtagc tgagctgatt cgggcccagag cctgccagag ggtggttgct 360
atggtggggg ccggcatcag cacaccaggt ggcattccag acctcagatc tccggggagt 420
ggcctgtaca gcaacctcca gcagtaecat ctcccgtagc ccgaggccat ttttgaaact 480
ccatttctct ttccacaacc caagcccttt ttcaacttgg ccaaggagat tttcgaaact 540
aactacaagc ccaacatcac tcaactattt tctcggtgct tccatgaca gggcgtgctt 600
ctgaggctct acacgcagaa catcgatggg ctgcagagag tgtcggcgat cctgtccctca 660
aagctgggtt aagctcatcg aacctttgcc ttgtgccact gcacagtctg ccaagacccc 720
ttcccccggg aggaacttcg ggctgacgtg atggcagaca ggtttccctc ctgcccggct 780
tgcaacggcg ttgtgagacc cgactattgt ttctttgggg agccctgcgc ccagaggctt 840
ttgctgcatt ttgttgattt ccccatggca gatctgtctg tcatccttgg gacctccctg 900
gtgcgtgagc ccttgaccag ccttgaccag gccgtgcgga ctacgttccc cgactgctca 960
tcaaccggga cttgttgggg ccttggcttt ggcattctcg cagcagggac gtggcccagc 1020
attccaggaa cgttccagcg gtggaaagcc tagtggagct ttgggctctg acagaagaga 1080
tgccggacct tgtgcagcgg gaaactggga agcttgatgg accagacaaa taggatgatg 1140
gctgccccca cacaataaat ggttaacatag gagacatcca catccaatt c 1191

```

<210> 65  
 <211> 2596  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 997067CB1

```

<400> 65
ccggtgtctg ggtctgtgac aggggtccaa agggcctggt ccgtgtccgg tcccccaaat 60
ctgtctgtcc tgccccagc cattggcact aacaaaagtc agaattcccg ggaacttgaa 120
cagaggctgc taaattccca gtaattgctc ctttggcctt ctagggactg acttcaagaa 180
aggaagtgaa gaattcccca gtgcttctct attctctttt aaaaccgcct tcccctgag 240
cttgacacca ggaacacaga gacacacctg ccttccatcg gaaactcagg ctgatctgt 300
attccaggaa cctcagcgcc tgcttgacag tgctcttctc tcaaaagttc ctgctctttc 360
cgacaaggac agcctggggg atgagatgtt ggcggctgcg ctcttaaggg ccaagtcaca 420
ggagcttgga acccttgagg atgtagctgt gtaatttata cggaaaggag ggaagcgttt 480
ggaactctgt cagaggggacc tctatagaga tgtgatgctg gagaattacg ggaatgtgtt 540

```

WO 00/44900

PCT/US00/02237

ctcactggat	cgtgagacca	ggactgaaaa	tgatcaagaa	attctctgaag	acacaagatc	600
acatgggggt	ctactgggaa	gatttcaaaa	ggatatttct	cagggtctca	agtttaaaaga	660
agcctatgaa	cgagaagtca	gtctgaaaaa	gccgctgggg	aactccccctg	gagaaagact	720
gaacaggaaa	atgccagatt	ttggtcaagt	gacagttag	gagaaagctaa	ccccaggggg	780
agagagaagc	gagaaatata	atgatttttg	gaacagcttc	actgtgaatt	ccaaccttat	840
ctcacatcag	agactccccg	tgggagacag	acccataag	tgtgtgaat	gtagcaagag	900
ctttaatcat	acttcagacc	ttattcaaca	tcagagaatc	cacactgggg	aaaagcccta	960
tgaatgtaat	gagtggtgga	aggccttcag	ccagagctca	caccttattc	agcatcagag	1020
aatccacact	ggggaaaaac	cttatgaatg	tagtgattgt	gggaaaacct	tcagctgtag	1080
ctctgccttc	attctgcctc	ggaggatcca	cacgggggag	aaacccctatg	aatgtaatga	1140
gtgtgggaag	accttcagct	ggagctccac	cctcaccac	catcagagaa	tcacactgtg	1200
tgagaaacct	tacgctgca	atgaatgtgg	gaaggccttc	agcaggagct	caacctttat	1260
tcaccatcag	agaatccaca	ctggagaaaa	accctatgaa	tgtaatgag	gtgggaaaagc	1320
cttcagccag	agctcacacc	tctatcagca	ccagagaatc	cacactggat	agaagcccta	1380
cgaatgtcat	gaatgtggag	gaaagtttac	ctacagtcca	ggccttattc	agcatcaaa	1440
aatccacacc	gggggagaacc	cctatgaatg	tagtgattgt	gggaaaacct	tcaggtacag	1500
ctcgggtctt	gttcgccatc	agagaattca	cactggagag	aagcctttga	atgggactcg	1560
catgagcaaa	agctccctca	gagttacag	cgagttaaat	atcagattgt	ccactgtgaa	1620
gagcccaaca	cccattttcc	tcactttccc	tgagctcaaa	gagctcttgc	cttaccttat	1680
aaatctcaac	agcttaggat	gtgtcccttt	caactcagac	ttttcaattt	agagaatggg	1740
gcagatgggg	caaatcggtg	aattttccca	gaaatcacac	cagccttaga	aaagcgtcaag	1800
ccaagtggat	ggcgtgc tgg	gaatagaaa	cagctctggg	accagttaac	ccatttagga	1860
aaggagtttg	cactaaactg	tttttctcac	agcagaggaa	ccttctcaag	gtgggtagtg	1920
aagcagcagt	ccagcagaa	tcgattggat	cctttagttg	gcttcgcgtg	atccagcaca	1980
gacagcagtg	gaaaggagcg	tctgggtctc	gttatttget	aggggagcta	aaaggagact	2040
atttcaagc	tactgttctc	agtcacagct	taagt ttcgt	taagaaagct	gcgttttgt	2100
ttcatgatgt	cgttaattat	ggaaaatttg	cattgaggga	ttttttatt	gagggtagaa	2160
gagatctcag	aatcatcatc	gtgtccttta	gtgtccttta	ggactcttgg	agcagccaga	2220
ccatgtttcc	aagagaatac	tggtgatatt	gccagcagac	cccttcccat	ccccccagat	2280
gtgtctgggg	ctgaatgggc	aaatctgtcc	aaacagctac	taacccggctg	gtggggagag	2340
ggtcagaagc	acttagcgtt	ggcctctgat	tgtctctctc	tcttctcttc	ttccactccc	2400
aatgtagtaa	atgattttct	ctaaatgctc	gggtaaggag	gcttccaagg	agctcaactg	2460
gctctgtctc	ctctgccctc	cacctctgac	accagccccc	agggaccaga	ccactcctgc	2520
ctccacctct	gactcttcag	cagcttgaga	ttaatgcaga	gaaagagcaa	agcccaaaag	2580
ggagaaaaaa	aaaaaa					2596

<210> 66  
 <211> 1574  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 1443262CB1

<400> 66						
gacgcccgtc	gacgcccgtc	gctggacgcg	ggtagccgtc	tgaggtgcgc	gagctgcggg	60
aggaatggag	cgctgaaagt	ggaaaagtgc	gcaaccggca	acaggggaaa	cgggctgcgc	120
gcgctgaccc	cgctgcgcgc	cggaagagta	ctcttcgcgt	cggaatccct	ggcgtacacg	180
gtgtgccaagg	ggagtcgctg	cgctgcgtgc	gacgcgtgcc	ttctcgggaa	gaaaagctgc	240
atgcagtgct	ctcagctgcg	cgctgcgcaca	tactgttagt	ctaagtgtca	gaaaaaagct	300
tgccagacag	acaagcggga	atgcaaatgc	cttaaaagct	gcaaacccag	atatctcca	360
gactccgttc	gactctctgg	cagagattgc	ttcaaaacta	tggatgggac	accttcagaa	420
tcagagaagc	tttactcatt	ttagtgatctg	gagtcacaat	tttaacaact	gactgaagat	480
aagaaaaggg	gcctcaggca	actcgttaat	acatttcaac	atttcatgag	agaagaaata	540
caggatgcct	ctcagctgcc	acctgccttt	gaccttttgg	aagcctttgc	aaagtgtact	600
tgcaactctt	tcaccatctg	taatgcggag	ctcagggaag	ttggtgtgtg	ccatatccc	660
agtatcctct	tgctcaatca	cagctgtgac	cccaactgtt	cgatgtgtgt	taagggccc	720
cacctctta	tcgcagcagt	ccgagacatc	gaggtgggag	agtagctcca	catctgcttc	780
ctgggatatg	tgatgaccag	tgaggagcgc	cgggaagcag	tgagggacaa	gtaactcttt	840
gaatgtgact	gtttccagtg	ccaaaccag	gacaagagtg	ctgatattgc	aactgtgtag	900

WO 00/44900

PCT/US00/02237

```

gagcaagtat ggaaggaagt tcaagaatcc ctgaaaaaaa ttgaagaact gaaggcacac 960
tggaaagtggg agcaggtttc ggccatgtgc caggcgatca taagcagcaa tctcgaaagg 1020
cttcoccgata tcaacatctca ccagctgaag gtgctcgact gcgcccattgga tgcctgcactc 1080
aacctcggcc ttgtggagga agcctgtgtc tatggtactc ggaccattgga gccatacagg 1140
atttttttcc caggaagacca tcccgctcaga ggggttcaag tgatgaaagt tggcgaactgc 1200
cagctacatc aagggcatgt tccccaagca atgaagaatc tgagactggc ttttgatatt 1260
atgagagtga cacatggcag agaacacagc ctgattgaag atttgattct acttttagaa 1320
gaatgcgagc ccaacatcag agcatcctaa gggaaacgag tcagagggaa atacggcgtg 1380
tgtctttgtt gaatgcctta ttgaggtcac acactctatg ctttggtagc tgtgtgaacc 1440
tctcctatgt gaaaatttgt tccgtgtttg ttgagtgtaa taagggtttg 1500
aaaccacaag aatcatttag ttgtagagaag cactgattata ataaattcaa aacatttgg 1560
tgaaaaaaa aaaa

```

&lt;210&gt; 67

&lt;211&gt; 2197

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 1521648CB1

&lt;400&gt; 67

```

cccgacgaac gcgaggaggg cgaggcgagg ggtgcaggag ggaccctcgc catgggtcca 60
cgggcctaga gtggcggaag ataccggcct ggtgccaaac ttgctactgc tgcctcctgt 120
ggcctccatg gctgaggact ggctggactg cccggccctg ggcctgtgct ggaagcgccg 180
cgaagtcttt cgcaagtcag gggccacctg tggagctcga gacacctatt accagagccc 240
cacaggagac aggaatcgaa gcaaaagtta gctgactcga taactggccc ctgcgtgtga 300
tctcaccttc ttcgacttca aacaaggcat cttgtgctat ccagccccca aggcccatcc 360
cgtggcggtt gccagcaaga agcgaagaa gccctcaagg ccagccaaga ctcgaaaaag 420
tcagggttga cccagagatg gtgaggtcag gaaggaggcg ccgaggggatg agaccaaggc 480
tgacactgac acagccccag cttcattccc tgctctgggg acgtgtgaga actgtggaat 540
cagctttcca ggggatggca cccaaggca ggggtcctaa acgttgttga aagactgtcg 600
agcacagaga attgccttca accgggaaca gagaatgttt aagcgtgtggt gctgtgggga 660
gtgtgcagcg tgcagggtaa tgcaggctgc tggggcctgc tccacctgcc tctgcagct 720
gccccatgat gtggcatcgg ggctgttctg caagtgtgaa cggagacgct gccctcggat 780
tgtggaaaagg agccgagggt gtggagtatg gtggggctgt cagacccaag aggatgtggt 840
ccattgcccc atctgccttc gccctccccg ccttggtctc agggcccgat ggaaa tgtgt 900
cagcgacgtg tgcctacggg gtaaacatgc ccgcgcgaag ggaggtgtg actccaagat 960
gggtgccagg cggcgccccg gagccccagc actgcctcca ccacccccat cagtctcccc 1020
agagccccaca gagccgaccc ccagagccct ggccccctcg ccacctgcgc agtctactta 1080
tactgtgtga gacgaggagc agctacagcc ctacacgaac cggccgcaga accgcaagt 1140
cgggactctg gcaagcctgc tacggcggat ggaactgtgc cgtctgcact tctgtcgga 1200
caagccccaa ttccgggggca gcaaccagaa gcgccagaag tgcctgtggc gcaaatgtct 1260
gcagttttgc atgaagcgcg tgctgccagc tgctgtgcta gactctgaga atgggcagg 1320
atcgccccca ccttaccgtg gtgcgaagag gccagctct cggccagcgc accactctgg 1380
ccctaccttg aagccccact tggctacacg cacagcccaa ccagaccata ccaggctccc 1440
aacgaagcag gaagcaggtg gtggctttgt gctgcccccg cctgtgcact accttgttgt 1500
tttaccggaa ggcgcaagca gtccgtgtga gttgcggggc cctgtgtcag ctctccaga 1560
agccctgttg cagggtgaag aagagaaggc ggataccaga gacgagtgtg caccaggcac 1620
gtcctctctg acttctcccg tatgtgtgct ttggtgccct ccaagaggag gcaagaggag 1680
cctgcctctc tccaatttgg cccagaggga agaggcagga gggcctgtga caccgttgt 1740
cacggagatt ttacagcttg gtggaaccgg cctccgagat actcggagct ggttgccaag 1800
gtccaaaagc cttaaaaaac ctggagctag aaagcagtag gatggatgt aagtgtggga 1860
tagggattca agtctgcagg gcaggcactc ggggaagggaa gatggatgt aagtggtgga 1920
gaccagagga acagtggagc ccacagcac gactgtgaa acacagagat ggctgtgga 1980
ccatgtcagt ctctaccac ctcagctctc gatgatgtgt gttgcttga gaagaagct 2040
tgccctcttc tcacagagt aatatgcgat ctggcccagg aattagagaa gctgaaagga 2100
tgactctggg gaaggtggaa cagctcagag cctggct

```

WO 00/44900

PCT/US00/02237

<210> 68  
<211> 2081  
<212> DNA  
<213> Homo sapiens  
  
<220>  
<221> misc-feature  
<223> Incyte ID No.: 1685494CB1

<400> 68  
gcatacgagcag tccgtgccta agcggcacga ccatcgagcc aaggctctcc agcgttcccta 60  
gagcggagaa gaaagcgctc cgaagagcta gagctgacac tcggcgatga gctaagacgc 120  
tgtttcagag cgttttgggtc ccttgaggcc ccttgaccag gagtgtctct gaagatcacag 180  
tccaaagaag gttctccaaa acaaggagag cagtctgaag ctggggatgg caacagcatt 240  
ggtgagtgcc catttcctgg ctcccctgag tctgaagaag gaggggcttc gggtagtgag 300  
ggagagatcac tactctactt attcaagctg caaggaaca gtaaaaggct 360  
tgagacaggag ccatttgcca acaatttcag gcagttgcgt tatgaagaga ccacaggacc 420  
tcgagaagca ctaagtccgc tcggggagct ctgtcaacag tggctacagc ccagaccaca 480  
taccaaaggag cagatccctg agctgctggt gctggagcag ttcttgatca tctgtcctaa 540  
ggagctccag gcccggtgctc aggagcatca ccagagagc agggaggacg tggttgttgt 600  
tctggagatg ttgcagctgg atctttggaga aacaggacaa caggtggacc cagaccagcc 660  
aaagaacaaca aaaatacttg tggaggagat ggccctctg aaaggagtac aggaacagca 720  
ggttcggcat gagtgtggaag ttacaagcc tgagaagaag aaggggtgag agacaaggat 780  
tgagaatggg aagcttattg tagtaacaga ctcttgtgga agagttagat cactctggaa 840  
aatatctgaa cccatggagg ctctaataga gggctctaac ttggaaaggc atcaggccaa 900  
gccaaaaggag aagatttgat ataaatgtct agaaagtgag cagagattca tccagcactt 960  
ggacgtgatt gaacatcgga gtacacacac gggaaagaaa ctctgcgagt ctgattgtgtg 1020  
tcagagttcc agtcttaccg gacataagaa agtccctctc agagagaaga gctatcagtg 1080  
tcagtgtgag gggaaagcct ttacaggagg ttacacacct gtcagatcac agaaaaacca 1140  
tctgtgtgat agtgcattatc agtgcaatga gtctggcaaa gtctttatgc agaatgcagg 1200  
ccttttggaa catctcagaa ttcatacttg agagaacct tctatgtga tctatgttg 1260  
aaaaaatttt aggcgcagct ctacacttaa ctgcacatc agaatccaca gtcaggagg 1320  
gcctgtgag tgcaggaggt gtggaaaaaa ctcttagtcag gccttaactc tcaccacaca 1380  
tcagagaatc catagtcaat ccaaaagcca gacaccacag aatctatac ttgagaaaac ctccaagt 1440  
tttgacctca gaccttattc gacaccacag atctgtagtga atgtggagaa gcttgcagc 1500  
taacatatgc cagaaaagcct tccgactaaa ctacacacct gctcagcag taagaatcca 1560  
caatgaagaa aaacctctatc agtgcacatg atcaccacaa agacaaactg gcttcagcgc 1620  
ctcttttcaa catcagagat gacacattga gcaacattga tagcaaacag cacttttaga aaagtcaacg 1680  
tagcccaatc tggcattcag aagtctcttc gaggctctcc tgctctatt 1800  
ctctctcatt tgccttcttc caagatttaa cagcattctt ctggaccaca ataattcac tgcagatgat 1860  
atgtcagtat ttatatactc taagcaatag agagtcagg catctcagag cacttaaacat 1920  
gtcagcagtg aatcaagctg ccacagattt cagggcttaa gactgagtaa gagttttgaa 1980  
tatttttggg tatgtatatg cagctttctc ctatgaaata a ggaagattga 2040  
2081

<210> 69  
<211> 2785  
<212> DNA  
<213> Homo sapiens  
  
<220>  
<221> misc-feature  
<223> Incyte ID No.: 1730829CB1

<400> 69  
gacctgtggg cgeccagcgg agtaggggct gcgcttgggg ttgtgtgaag ctggctgcct 60  
ctcccaactc cctctttgggt gcaaaagcgc gctcagcggg agacatgggc cggcgccgga 120  
caggggcaac tgcgtacttg actgagagcc tgcggccagc ttacatggag ccaccccgcc 180  
ccggcgccga cccgacgcga cccgaatcga tccgatccca tctccatcgt tctcgtctc 240  
ctcccgctct gaccgtgtgc ccggccgtgg ttgcgcacac caggcatcca aagctgaggt 300

WO 00/44900

PCT/US00/02237

```

cgctctcaagc gctctgggctc gccttcgctt tagagatggt tggcctcttc cctcccaaac 360
agctcatctt caaaactctg actcttggac tggcacctgg ccacctttcc ctctaccaag 420
actccacttc cgtcttaacc actctctctc cagatctctg gtacccccct ggttggagac 480
tgctcaattt ccttccaaat taatcccaaa cccctcaaaa tattgacaac ctctgacaac 540
ccccaccgga ggagccacagc tttgttttgg actaacttcc atagccctat catggaggga 600
gtgtacctgg tagtgaatgg gttgggctcg gtgctggagc tgctgacctt ggtgttggac 660
ctcaactctc tgctgtgttg ctccctctcg gcttccctgg cctggctcctt ggccttctgt 720
tacaactctc cgcacacggt actgactagt ctctcgact tgggcccgagg agtcttgcct 780
tcaatgtctgg ccttgactga agccgtggte cggttcacat gtgggggctt gagggctctg 840
tgtatcttgg tgtatagctc ctgctctggc ctagagagcc taaagctctt ccggccacctg 900
gcctctcatg gggcacttgc gaggcaggag atactgcacc gggggcgtctc caatgttggtc 960
tccagtgccc atgctttgct gcgccaggcc tgtgacatct gtgccattgc catgagcttg 1020
gtggcttatg tgatcaacag cctggctcaac atctgcccca tcggcactca gaacctcttt 1080
tccctgtgtc tggccctgtg gggatgcagt accgggctct tgtggaggat gacagacgta 1140
gtggctgcct tccatgccca catttccagc agtgctgtgg ccactggccat cctcctttgg 1200
acaccctgcc aactagcctt gggagctgctg gcctcagctg ccgcctcctc gggcagcttt 1260
gtgcttgtca atctcactgg cttggtgttg ctactcctct tgctggcagt gacggtgact 1320
gtgttgcata cggacttca cctgaggctg gctaccagg cactcagcca gctccatgcc 1380
cggccatctc accaccgtct tcgagaggat gtcacgcggc tctctcgctc agcactgggc 1440
tcagaggcct ggcgcgcagt ctggagccgc agtctgcagc tggcgagttg gccaaaccgg 1500
ggagggggcac ctggagctcc ccagggtgac cctatgaggg tattctcagt taggaccggg 1560
agacacggca cttctctctga agcgggggcgc agatcagagg cagaaggagga gggaggccagg 1620
accatcagag tgacacactgt caggggccga gagaggctca atgaggaggga gcctccagg 1680
gggaagagag cttggaaatt gctgaaggag caagaggagc ggaagaaagt tgtcatctgc 1740
caggacacga gcaagacagt gttgctcctg ccttgcggcg atctgtgcct gtgcacggcc 1800
tgcctgtaaa tccctgtgcg ccaccctctc taaccacgca attgcctcct ctgcgcggcg 1860
ggcatctctg agacccctca tgctacactc tgaagcctcc ttccctgcct gccaccctc 1920
ccatgctcca cgcagcgact cagcttagga cagcattaac accctcctct cgggtcctgg 1980
tctgaatccc ctctaccccc tgtggccatc ctgccatata tccaggacat tgaagtggaa 2040
gaactatgat tgggtggggg caggataaca tggcttctct ttaccaggct ggtccctctg 2100
atgctgaggg tgggtgagatg gtcactatgc aagggccctg agactatttg ctgtggggctc 2160
tctccacagg tgcaccaggc ccaccagatg gcctctgggg ttaccctgtg ctgcctctgg 2220
ttttctctgt ggagatctat aggtcctttt cctgcctcct tcacatttcc tcccagctt 2280
tgcgggccac aacacatcag tgtcatttgg gtgttttggc aactcagggg ccttcggatg 2340
atcttaaaac ttgtgttcca gccagagccc ctgtgcccct gtaggcggtg ggggttagat 2400
ctctcgctcc cctcagagcc cactctgcct tgtgatctgt tgatgaggct cctcccaac 2460
ctgatccaaa agccagctctc agggatttacc ccttgggagt ggggatgcat ctgcacctga 2520
ctttggggct tggcgccttg tggcaccoca gctcactggg agtctcagga ttctcaaacg 2580
gatttctgct ctttcccttg tcactccca cacacacaga aaaaatggcat tctctctgt 2640
ctctccctgg actggagagg cagagactgt cacatttacc tagggttcaa atcagaagg 2700
ccaggggccc aggggttgca gctctgtgag ggttctctgg cccagtttcc aatgaataaa 2760
gttctcttga cagctaaaaa aaaaaa

```

&lt;210&gt; 70

&lt;211&gt; 12131

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 1864641CB1

&lt;400&gt; 70

```

tccgtctcta tgacgtcagc cgttaaggcgc tgtgtctgta aaaggacgtc cggctcgtct 60
cctagtgtac ggaatcggtc gtcagcctcc ctggctgtta gtacctcttt tcccgaggat 120
ctgtgtccagc agttggattt actgtgtgtc cgggtgggcc tcaagccatt cctgttccct 180
cggccccctc agtgagtccg gctctccggc gaaagtgtagc gaggtttggc cggagcgccg 240
acgaggggaa aatgcctaaa aaaaagactg gtgcaggaaa gaagcgttag aaccgcggag 300
aacgtgaaaa acaactaaga gcatcaagaa gcactataga tttagctaaa ctctcatcta 360
atgcctcaat ggaatgtgac aagtgtcaga ggccgacgaa gaatagagca tatctgtact 420

```

WO 00/44900

PCT/US00/02237

tttgaatttc	tgtacagaag	ttaccaattt	gtgcacagtg	tgggaaaaca	aagtgcata	480
tgaagtcttc	agactgtgtc	ataaagcatg	ctgggtgtata	cagtactggc	cttgcattgg	540
tggtgtcgaat	atgtgcattc	tgtgaagcct	gggtttgtcca	tggtaggaaa	tgtctcagta	600
cacatgcttg	tgcctgccct	cttaccgatg	ctgagtggtg	tgaatgtgaa	cgaggcgtgt	660
gggaccatgg	aggcagaata	ttcagtgtgt	ctttttgtcca	taactttctc	tgtgaagatg	720
atcaattttga	gcatacaagg	agctgccagg	tttttagagg	agaaacattt	aaatgtgttt	780
catgcaatcg	gcttgggtcag	cactcatgtc	tccgttgttaa	ggcttgtttc	tgtgatgata	840
atacaaggag	caaagtgttt	aagcaagaaa	aaggaaaaca	gcctcctgtg	cttaaatgtg	900
ggcatgaac	tcagggaact	aaggacctta	gcattgtcaac	acgctccctg	aaatttggca	960
ggcagactgg	aggtgaagag	ggagatggag	ctcttgggta	tgatgtctat	tgggaagacc	1020
tttcatctga	taagtatggt	gataccagct	accacgatga	ggaggaggat	gagtatgaag	1080
cagaggatga	tgaagaggaa	gaagatgaag	gcagaaaagg	ttcagatact	gagtcacag	1140
atttgtttac	taatttgaat	ttaggaagga	ccatgtctag	tggctatgct	cactatgagg	1200
aacaagagaa	ctaggggagc	tgccttggtg	g			1231

<210> 71  
 <211> 700  
 <212> DNA  
 <213> Homo sapiens  
 <220> -  
 <221> misc-feature  
 <223> Incyte ID No.: 2444604CB1

<400> 71	gccccagggtc	acacaatggc	cgagctccat	ggcggtctgc	ttcttcacgc	cttcatgtgc	60
	accgcgcttc	ccagggggcc	cccggcccac	cctgcggatg	ccgagtcagc	ctcccgcagc	120
	ccctccctgg	tcccagcccc	tcttccctgg	cgccatggag	ccctcccccac	gagccccagg	180
	gcataccgagc	atggggcgcc	caatgcagag	gggtgacgct	cctctgtggca	tggccagcgt	240
	ggggccccag	agctatggag	gtggcatgag	acccccaccc	aactccctcg	ccggccccagg	300
	ctcgctctgc	atgaacatgg	gcccaggagt	tcgtggcccc	tggggcagcc	ccagtgaaaa	360
	ctcgatcccc	tactctctct	catccccccg	cagctacacc	ggacccccag	gaggagggtg	420
	gccccctgga	acacccatca	tgcctagccc	tggagaltcc	accaactcca	gcgaaaaact	480
	gtacacatct	atgaacccca	tcgggcaggg	cgccggcagg	gctaatttcc	cgctcgggcc	540
	tggccctgag	ggcccattgc	gcctatgagc	gcgatggagc	ctcaccaagt	gaacgattcc	600
	ctgggctcgg	gcgacatgga	cggtgtgcgc	aagagtctcc	ccggcgccgt	ggccggcctg	660
	agcaacgccc	cggggcaccc	cgcgggacga	cgcgcgagatg			700

<210> 72  
 <211> 2332  
 <212> DNA  
 <213> Homo sapiens  
 <220> -  
 <221> misc-feature  
 <223> Incyte ID No.: 2445008CB1

<400> 72	agcgaggagg	ggaacgagc	cgctggccat	tttgtgtctg	cttctctgtg	gacgtgggtg	60
	tagccgtttg	gttgggaaa	tgagggtatt	ttggctcgt	ttctcctgct	tcttttctcc	120
	tccctttttc	tttgcggta	gaacacagtt	atgggtgtcca	agaagaagaa	cgagctgaag	180
	cgctgggtgt	ggtattgtaa	tagagatttt	gatgatgaga	agatctctat	tcagacacca	240
	aaagcaaaag	atttttaagt	ccatatattt	cacaagaaat	tgtatcacag	acctgggtta	300
	gctattcatt	gcattgcaggt	acataaagaa	acaaatagatg	ccgtaccaca	tgcataacct	360
	ggaagaagac	acatagagtt	ggaatatatat	ggtatgggaag	gtattccaga	aaaagacatt	420
	gatgaagaag	gacgactctt	tgaacagaaa	acacaagaaa	gtccaaaaaaa	gaagcaacaa	480
	gatgattctg	atgaa tatga	tgatgcagac	tctgcagcct	caactctatt	tcagccacag	540
	ctcgtttcac	ctgcagcaagg	ttatatctct	ccaatggcac	agccaggact	ggccaccagta	600
	ccaggagcac	caggaattgcc	tccaggcata	ctctccattaa	tgccagggtg	tctctctctg	660
	atgcaggaaa	tggccaccagt	tatgccaggc	atgccacctg	gattgtcatca	tcagagaaaa	720

```

tacaccagct catcttgcgg tgaaaacata atgatgccaa tgggtggaat gatgccacct 780
ggaccaggaa taccacctct gatgcctgga atgccaccag gtatgccccc acctgttcca 840
cgtctctgaa ttcctccaat gactcaagca caggctgttt cagcgccagg tattcttaat 900
agaccactcg caccacaacgc aactgtacct gccccacagc ctccagttac taagcctctt 960
ttccccagtg ctggcagcagc ggggacacct gtcacaagct caagtacagc ttcatccaat 1020
tcgaaaagtc tgtctgcatc ttctaagctc ctgttttcta gcacagcaca agctcaggca 1080
gctgttccaa gacctgttgg tacagatttc aaaccttaaa atagtaccct tgcaacaact 1140
acagaaacccc caaagcctac attccctgct tatcacagct ctacagcttc acaactagct 1200
acaacaaata gtactgcagc taaccacgcy gcttcaataa caagttaagc tgcctacact 1260
acaacaaata gtgcaaccag taagtgtatc catccagatg aggatatact cctgggaagag 1320
agaagggcac agttaccata gtatcaacgt aatcttctcc gcccaggaca ggcccccatc 1380
ggtaatccac cagttggacc aattggaggt atgatgccac cacagccagg catcccacag 1440
caacaaggaa tgagaccccc aatgccacct catggctcag atggttggtca tcatcaaggc 1500
atgccagagt accttctcgg tgctatgccc ccgtatgggc agggcaccgc aatgggtgcc 1560
ccttaccagg gtgggacctc tcgacctccg atgggaatga gacctcctgt aatgtcgcaa 1620
gggtggcgtt actgattcta ctctatccag tctaataagt ttggagatta aaccttttct 1680
caacttctgc tgtttatata gccaaagctc cgtcaataag gcttcatagg gaacttaaca 1740
aacattatct tcccacatac caggaaacta tggacattta ttttaccatg aacattttct 1800
tgggaataat aaagcaggaa cttttctcga agttgcaatt tataactgtt ggtctctttt 1860
tcattgttca tctagggttt tagaagttaa gttatgttaa tttggttcgt taaaattgtg 1920
agggcctgga attacatgca cataccacc ctagtaaggg aagtctctga agcttcaact 1980
gctatttcta aagtttgcct tcacagcatt tcagatgctg ttggactgtt tgcctccaac 2040
ctagcttggt gagggtctga actgtttcca agtacttgta cattggaagt ctgaatgtgt 2100
aacaataatt attgtattta gagtctctca tttgtcaggg ttgttcaggg taacccacc 2160
aaggtcattt gacttttctg tactgttaaa cttcatttga ataaaatgag agaaaaattt 2220
atgctctttt attcataacc cagctgtgga ccactgcctg aaaggtttgt acagatgcat 2280
gccacagtag atgtccacat aataaaattc atagttacca aaaaaaaaaa aa 2332

```

&lt;210&gt; 73

&lt;211&gt; 1936

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 2572462CB1

&lt;400&gt; 73

```

ttggttctat attactgttt tatagctgag gagaagaact cacatcattt actaagatca 60
tatagctagc tagtaaatgt tttagctgaa atacaaaacaa aggtttttctg acttttaagag 120
ctttagtttt ttccactata ccatattgca tctgttgtaa ttgttaacta atcttgactt 180
taaaattctc atttgcctta tgtactgagc ccttatacca gtgctaattt atgtgactcc 240
ttctctctgc agaagaagaa aaaaatcctt tttaattcat ttatagtagc cagtttttaa 300
agaagattta ttttgtaaaa ttttgcctat ggtacatgct atcttagcct gtaataaata 360
taaaagcata atttttatec ctccctggtc ttctctctct tetgacttta tacgtctctc 420
tagagagctt atcttctata ataaaatctc ttgtttttaa agtggaagaa atcagttcaa 480
agaaaaggag aagaaaagtg aaaaaacatc tctctctcgg gctacctctt ctgcacctca 540
gctagccagg gccccagaac aaacaccgat ggcctctctc ccacgtccca agatggatgc 600
aatcttaact agggccatta aggcattgct ccagaagagt ggtgcatcag tgggtgctat 660
tcgaaaatac atcatccata agtatccttc tctggagctg gagagaaggg gttatctcct 720
taaaacaagc ctgaaaagag aattaaaatg aggatgcac aaacaggtta aaggaataag 780
tgctctcgga agttttgttg ttggttcagaa atcaagaaaa acacctcaga aatccagaaa 840
cagaagaagt aggagctctg cagtggatcc atgaaccaca gtaaaaattg gtttgaagaa 900
cccactggcc tttaactcgc tttgtgaacc taaagaagct tctcactcagc tcatcaggaa 960
atatgtgctt cagttatctc cttaagcttag agtggttagc atggtgacatc agggcctcagc tgttgaagaa 1020
ggctctgtag agagcagtag agagggggcca gttagaacag ataacctggca aaggtgtctc 1080
ggggaacatc cagctgaaga aatcagggga gaaacccctg ctgtgtggaa cggctgatgaa 1140
atatgcaact ttgtctgcca ttgtctgcat gaatgagccg aagacctgtg ctaccactgc 1200
tctgaagaag tatgtctctg agaatacacc aggaaccaat ttaactatc atcactattt 1260
gctgaaaaaa accctcgaga aatgcgaaaa gaatgggttg atggaacaga tctctgggaa 1320
agggctcagt gcgacctctc agctctgttt tctctattat cccagccagc gagttctgtt 1380

```



WO 00/44900

PCT/US00/02237

```

tccgaagaaa gagccagatg attctagaga tgaggatgaa gatgaagatg agtcatacaga 1440
aagaagactct gaggatgaag agccgccacc taagagaaagg ttgcagaaga aaaccccgagc 1500
caagtccccc ggaagggccg catctgtgaa gcagagagagg tccaaacctg caccataaagt 1560
ctcagctccc cagcggggga aagctaggcc cttgcctaag aaagcacctc ctaaggccaa 1620
aacgctctcc aagaagacca gacctccatc cacagtcac aagaaacctc gtggtggctc 1680
ctcaagaagac cctgcaacca gtgcaagaaa ggaagtataa ttgccgggca agggcaaatc 1740
caccatgaagc aatgtcttca gagtgaataa gtaaatatta taggaaaaaa ggggtatcatg 1800
atgaataata aatatctatt ttctaaggct agtgtgcatt tgtttagttt tgatgctttt 1860
caaatcacat tattttcttc ccctatgaac attgtgggga gggactctaa ataacacctg 1920
ttaggcaaaa aaaaaa
1936

```

&lt;210&gt; 74

&lt;211&gt; 1667

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 2572892CB1

&lt;400&gt; 74

```

cggaatgcc cgaccccgag gctctgacca ctatgccgag ctctttcttc gtcaggaagc 60
ctcccgagcc caatcggaag cctaactaca gcgagctgca ggactctaata ccagagttaa 120
ccttcaggaca gcctctagcac caggcccccac tgcctggcagc catccaccact ccggagatcc 180
tcaaccccac gcctcgcgtg ccaatgctca tctgggaactc tgcctgggagc ccccaagccc 240
agccaattgc ctgggctccc ctctggctcc aggagagctcc cagggtggcca gagctgacct 300
ccctgtcaga caggagcagt gggaaaagct cccagccccc cagcccccac tcaccggctc 360
ctctgctctt ctctctactc tcagcctctt ccttggaggc cgaggccctat gctgctcttc 420
caggtctggg ccaagtgcac aagcagctgg ccagctctc tgaggccaag gatctccagg 480
ctcgaagggc cttcaactgc aaatactgca acaaggaata cctcagcctg ggtgcctcca 540
agatgcacat ccgaagccac acgctgcccc gcgtctgcgg aacctcgggg aaggcctctc 600
ctaggccctg gctgctacaa gcccatgtcc ggaccacac tgccgagaaag cctctctctc 660
gtccccactg cagccgtgccc ttgcctgacc gctccaacct gcgggcccac ctccagacc 720
actcagatgt caagaagtac cagtgccagg cgtgtgctgc gactctctcc cgaatgtccc 780
tgctccacca gcaccaagag tccggctgct caggatgtcc ccgctgacc ctcaggctccc 840
ctcttctctt ctaactctgc cctgctctga cagcctctcc cagctccagc aggaaggacc 900
ccacatctct ctcaactcca tggaaatccc tctgtagtg ccactctctg gccacatcag 960
ccccacagga ctttgatgaa gaccatttct tgggtctctg tctctgctct gggctctgga 1020
agaggccctc ccgtggccat ttctgtggag ggaggggcagc tggccccagc cctgggggga 1080
ttctctgagct ggctctgtct cgtgggtttt tgtatccaga gctgtttgga tacagctgct 1140
ttgagctaca gacacaaagg tgacagactc actgggaagc tccccacca ctcaggggagc 1200
cccaactccc tcacacacac cccccacaa ggaacccctc ggccacctc cagcaggtgt 1260
gactaacatc gcaataatcc acccccaggt gcagccccag ggcctgggca gcgggtggca 1320
gactagagtc tgagatgccc cgagccaggt cagctatttc agcctcctgt ttggtggggt 1380
ggcactgtgt tccccgggcaa ttttaacatg tctgaaaagg gactgtgagt atgtgctgct 1440
actgttcggg ggccccaggt ggggtgctct gctcgaccga tgtgtctccc agaactatct 1500
tggggggccc acaggtgggc ctggggaggaa gatgtttaca tttttaaag tacactagta 1560
tttatatttc aaacattttg tatcaaggaa acgttttgtat tagttatatg tacagtttat 1620
tgaattatca taaagcagtt aatttatata ttaaaaaaaaa aaaaaa
1667

```

&lt;210&gt; 75

&lt;211&gt; 759

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 2785674CB1

&lt;400&gt; 75

```

tgggctcgcc      tccctggggac      taggtttcag      cggccgctgc      gatgacacaa      ataaaggcag      60
atccccgacgg      gcccgaggct      caggcggagg      cgtgttccgg      ggagcgaccc      taccaggagc      120
tgctgtgtcaa      ccagaacccc      atcgcgcagc      cctctggcttc      tcgcgcctc      acgcggaaagc      180
tctacaagt      catcaagaaa      gcggtgaaag      agaagcagat      tcggcgcggg      gtgaaaggag      240
ttcagaana      tgtaacacaa      ggagaaaaag      ggaatcatggt      ttgtgcagg      gacacactgc      300
ccattgaggt      atactgccat      ctccagtc      tgtgtgagga      ccgaatattt      cccatgtct      360
atatccccct      taagacggag      ctgggtgcag      ccgcaggctc      caagcgcccc      acctgtgtga      420
taatgtgtcaa      gccccatgag      gagtaccagg      aggcctacga      tgagtgcctg      gagggatgtg      480
agtcctctgcc      cctaccctca      tgagggggct      cggtagcacc      tggggacactg      ccgctggaaag      540
ctattgggct      gcgacaggga      cgaactggctg      tctctctgc      caccacact      gacggcatct      600
tcccagttcc      ccaaggcacg      ccttcttccc      aggcagctct      aacagccct      teatgaaggt      660
aatgctagtc      ttctgtccat      cagtgccatt      tctgtgagaa      ctaaaggctg      ttccaagaat      720
gtgggggtggg      gaaagtcaat      gctaagacta      aaaaaaaaaa

```

```

<210> 76
<211> 1421
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 2797479CB1

```

```

<400> 76
ccccagtcac      tgttctctgcc      cgttgccctgt      cagcctcact      gcctttaatc      cgcttcacagg      60
gctcatctcac      ctgaatatata      tggttgcccatt      gctgagatat      ggtgaggata      ctgcgcaagg      120
catccccggat      ccgaacccctc      tcaaagtagct      gttgatagtg      ctggagctcc      acgggtgacat      180
gagctaacacag      gcgctgatca      tcatgggtga      tgacatatcg      acgcacatag      cccgcacaaag      240
acttagacac      aaacatccca      gctctgttga      gaaagtgtcc      caggttgtta      agcagctcag      300
aatatattctt      cagcagcagg      tccgtccagg      gctctgtccc      tgagggcgcc      tcaggccgaa      360
tgtagacgag      atagaagcgc      cagatgtcag      cagggatccc      cgtgtcctgg      gctttgattt      420
tctctggttca      gcacgaattc      atgaaggaaac      tcacactcta      gagaaacct      atgaatgtaa      480
gcaattgtggg      aaatttgttat      cteatgcctc      aagctttcga      agacacatga      tggcacacac      540
tggagatggc      cctcataaat      gcaactgtatg      tgggaaagcc      ttgactctc      ctagtgtatt      600
tcaaaggacat      gaaaggagtc      acactggaga      gaaacctat      gaatgcaagc      aatgtgggaa      660
agccttccgt      acttcaggct      cccctcgaaa      acatgaaaca      acacacactg      gagagcaacc      720
ctataaactgt      aaattgtggaa      aagcttttag      tgattttatt      tcttttcaa      gtcattgaaac      780
taaacacagat      gaagcgatcg      cttatgaatg      taaggagtg      gggaaaagcat      ttagttcttt      840
aaaatacttt      tgtcgccatg      aacggactca      cagtgaaaga      aaactttatg      agtgtcaaat      900
tgtgtggcaaa      tcttcagtg      ttccagttac      ttaaaaactc      atgaaaggac      tcacacggca      960
gagaaagccat      atgaattgag      gcaatgcagg      aaagcattct      ttggccctc      ttctctcta      1020
agacatgaaa      ggaactcac      tggagaaga      cctatgaat      gtaaacactg      acataaaaagg      1080
ttcagctggt      ccagttctctg      agccttcagt      tctctcagt      acgctggaga      gaagccctat      1140
gaatgttaagg      aatgtgggaa      agccttcagt      ggaatgtggg      aaagactca      ttggttttat      1260
acacactgga      aggatattct      taattgaaag      ataataaatcc      taagtatgta      acgtgtgtaa      1320
cacattcaga      tacttgaaag      cggctcacac      ctgtaatccc      agcactttga      gaggccagg      1380
agtttccagg      ctggggcgag      acgaggccag      gagatcgaga      ccagcctggg      g      1421

```

```

<210> 77
<211> 2386
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 2960640CB1

```

```

<400> 77
gactttttaag      ggatcacaga      gctcacacca      aagaccaggg      gaacagtcag      aagcctggct      60

```

WO 00/44900

PCT/US00/02237

tgctcctcag	gtcctccagga	acctgcctca	aaacacaggt	ctccacgacc	aggagacagg	120
tgtctgtgtc	tggagagctg	ggctccaggg	accagccatg	cgtgacaaca	gagctgtatc	180
cctctgtcag	caagaatgga	tgtgcccagc	ccctgcacaa	agggccctct	acagggggtg	240
caccacagag	aaggacagtc	acgtctcgct	ggcaacaggt	gtgcccctggg	gctatgaaga	300
gaccaagacg	ctccttgcta	tctcttagtag	tcttcaattt	tatggaaaac	tccagaccgc	360
tccagcagaac	agccagatct	acagggccat	ggcggaaagga	ctctggggagc	aggggtttct	420
gcggaagccca	gaacagtgtc	gcaccaagtt	caaaaagccta	cagttgtagtt	accgcgaagt	480
gaggagaccag	cggtgtgcctg	agccttgtag	cttttatgag	gaaatgaatg	ctcttcagg	540
ctcctgggccc	tctgcaccctc	ctatggcaag	cgtatgctgt	cctggccaag	aagggaagtga	600
tattgagctc	ggagagctga	atcaccagaa	tggggaaaccc	acggaggtag	aagatggcac	660
tgtggaatggt	gcagacagggg	atgaaaagga	cttcagggaat	cctggccagg	aagtcaggaa	720
actagacctg	ccagtgcctgt	tcccaaacag	acttggtttt	gagtccaaga	acgagattaa	780
aaagaaaaat	ctaaaaatggg	atgatccaga	ggaagttagaa	ataaaccaag	ctttacagag	840
aaagtccaga	ggagtttatt	ggcactctga	gtacaaaaaa	ggcttggaga	gtgagccaac	900
atcaagaagg	caatgtagaa	attctccagg	ggagagttag	gagaaaaacc	catcccagga	960
gaagatgagt	accagagatt	tttgtgccag	ggacaaaagcc	tgtacacata	tcctctgtgg	1020
gaaaaactgc	tctcagagtg	tgcactctcc	ccacaagcca	gcgtctcaac	tggaaaaagt	1080
atctcaatgt	ctcgaatgtg	tgaaaaacct	tagccgaagt	tcttatcttg	ttcggcatac	1140
aagaatccac	acaggcgagga	agcctcacaa	gtcgagttag	tgcgggaaagg	gcttttagtga	1200
cgctctcaac	ctcagctccc	acctcagaac	tcacacaggg	cagaggccct	atcagtgttg	1260
gcaatgtgag	aaaagcttca	accagagttc	cagcctcaat	gtccaccaga	ggaccatac	1320
cggggaaaaa	ctttaccagt	gcattgtctg	tggaaaagaga	ttcaacaaca	gttccagctt	1380
cagtgctcac	cgggcgcatcc	acactgggga	gagcccatat	aagtgtgcag	tgtgtgggaa	1440
aatcttcaac	aatagctccc	acttcagtcg	ccaccgaaaa	accaccaactg	ctgaaaagcc	1500
ttaacagctg	tctcactgtg	agagaggcct	cactaaagaa	tctgcctcga	cccgatcata	1560
gacagtagct	atgaaaagag	tactctcatc	acaggaaagga	agagatgcgt	tatgagtgtg	1620
tcggtaaatc	gtcagatata	gttctctcag	tcagcatgta	tgagctttct	tctgtgtgtg	1680
agagatctag	ccagtccctg	actttgcaac	agacctactg	actactggat	cttaagaccc	1740
atgtctcagg	ccaggtagtca	gcataaggac	gctgaacctc	ctcggctgtg	ccgtgtgact	1800
cagagctcta	tcttactgtg	acttaagttt	tgatggagag	aaagcttgat	gatccataaa	1860
tctaccagg	aatccagggt	ctctctgttc	ctagcactga	gaatggggca	ccagtggtcc	1920
aagaacactt	tctgggcttaa	catagtctct	acacagggtc	gagaaaagaa	gttctctctt	1980
tctgtgaaag	acatgtataa	agttaaaggg	ctgaattctc	tctaaaccaa	taattgacct	2040
ctaggccacc	acogtaatac	tgctaacttc	agagcagaag	acactgtctc	ttgtaaccca	2100
gccaccacca	aaaagcacaac	agaaaagaagg	aaggatgggt	aaggccattgg	ataatactga	2160
atctgttttc	ctaatgtact	taaccttaga	gcaagccaaa	ctccagggtt	aattaatgtt	2220
aagatttttc	ctctcattat	tgcctctcat	atagtctctg	attgtctctt	aaagtaagtg	2280
gtttatagac	attactattt	ctgataataa	tttacaacct	actataaaca	aatttataac	2340
cattactaat	tctctgatgaa	ataaaagttg	tttctccttc	caccac		2386

<210> 78  
 <211> 1432  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <221> misc-feature  
 <223> Incyte ID No.: 3454051CBI

<400> 78						
agtatgagct	ggacctgtcc	gcgttgccag	caacctgttt	tctctcgctga	gaagggtgagc	60
tcctctgggca	agaactggca	ccgcttctgc	ctgaaatagt	agcgtctgcca	cagcatctctg	120
tcctctggcga	ggcatgacga	gcacaatggg	agggccatct	gccacaagcc	atgctatggg	180
gctctcttgct	gaccacaggg	ccctcccat	atgaagacat	ctactggggga	gacctcgtgt	240
gctctcttgct	gtggggagacc	cgcttatatt	gctgagaagg	tgatgtcaat	aggcagaagt	300
tggcaccgac	cggtgtctgag	gtccagcgt	tgccacaaga	ccctgactgc	tggtgagcat	360
gctgagcatg	atggagctcc	ctactgccac	gtccccctgt	acggctacct	gtttggcccc	420
aaaggtgtga	acatgtggga	tgtgggtctg	tactctatg	accagtgtaa	gataaatttc	480
aaatgagacg	ctccacaaaa	aggtcacacc	taactcaggc	tccecatctg	ccccctcagt	540
tccaattggaa	gtcacaaaaa	tctccagctc	catgggggtg	gggggaaggtg	ggaattctggg	600
ggcctgtggcc	taggtctccat	ggtagggcag	agagctcaga	ctttctctgc	caattttctc	660







WO 00/44900

PCT/US00/02237

gatctgtgtt	cacagacatc	tgggggaaga	aaaggagcag	gaaactacc	cgccacagagt	240
taagcaggaa	acaacaacaa	catcatgcaa	aaaccctgca	aagaaaacga	aggaaagcca	300
aagtgcagcg	tgccaaagag	ggaggaaaaa	cgcccgtatg	gagaatttga	acgccagcaa	360
acagaaggga	attttagaca	gaggctgtct	cagctctctg	aagaatttaa	agagacata	420
gactataggc	attttaaga	tgaagaaatg	acaagggagg	gagatgagat	ggaaagggtg	480
ttggaagaga	taaggggtct	gagaaagaaa	tttagggctc	tgcattctaa	ccataggcat	540
tctcggggac	gtccttatcc	catttaatta	atctctctga	caattcaatt	atcttctgtt	600
attaatgttg	ccactgtctt	ctgtttgtct	gcactttctt	gataaatttt	tgcattcgtt	660
ttactccagt	catttcgatg	tgctgagatt	tacatattgc	tcttgtcaac	atctcatctt	720
ttgaccacat	cttattcatt	taataagag	tctcattcat	tgtcatggaa	gaatggccat	780
tgtatatgcc	aaagtgaaaa	taacgagttg	caaaacagtg	tatacatata	tgtgtgtata	840
tatgtacact	tattttgtac	atctctatgt	gacataatgc	aaaggaaaat	gtctgtattt	900
attatcaccc	aaaggttaac	agtgaattct	tgtgtgatct	cttttttttt	ctttttgtcc	960
atctgcattc	tctcacttgc	caaaaaatga	atataatgtt	atgtgtgtat	at tactgtgt	1020
tcacaaaaaa	ccctaaagta	gacagtaaaa	gaacttgtca	atcgcccttg	gaaggcaatg	1080
aaacacttaa	taaacctcca	ataacagaga	cgtaaaaatg	aaatgttaac	ctccaattac	1140
ctctggtact	cttagccaga	gtaataaaat	ggtaatattt	acagataaaa	aaaaaaa	1197

<210> 84  
 <211> 2170  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <221> misc-feature  
 <223> Incyte ID No.: 669862CB1

<400> 84						
agcgtgcgcg	ataggacagg	gccttaaaat	catggttatg	tgtttgtgtt	tca:tctaa	60
ataaagccga	accactgaca	aaattatata	gagagcaatg	tgataatggt	aggaatagag	120
atattccagga	catactctcg	tgcattctga	tccactcagt	ctgagttgat	atctaaaga	180
tactgtatgt	aaaatgttgt	tatggttaat	ctgattctga	actaaatagt	agaactagtg	240
actttgtgtt	tgcctactca	taaatctttac	atctctctgt	gcactttcat	aattttctgt	300
gcctcttctg	ctctttatat	attgctttggc	agtaactttg	aattttctaa	actcgtttgat	360
tcaatacaga	tacacttttt	agttgatcac	tatttttaat	atagatgtcc	tctccctacc	420
ctgtattatt	gaaaataatgc	atttgtttgt	ttttctcatt	tctccagag	cttaatttca	480
caactgaaag	caacaagtta	tcttcagaaa	aaagaaaata	tgaagttaat	gcgtaccact	540
aggagacatg	gaaaagaatga	aaaacccttca	acccttagag	tgacttaatt	aggatgacc	600
ccagactcac	aattgaaatt	gggagacaac	agagacctaa	agtggaatgt	tttagtcaaa	660
tgtatttcaa	aaaacataaaa	tccctctctc	tacataagag	aaataacaca	agagagaat	720
catatgagtg	taaggaaatat	aaagaagggt	ttagaaaaat	tttgaccctt	actgaacatc	780
tgagagaccta	tactgtgtgtg	ataccctatg	aatgtaatga	atgttgaaaa	gcattttgat	840
ttttccagca	tttttataga	catcgaaaaa	tccacactga	tttgaacccc	tatgaatgca	900
atggatgtga	gaaggcccttt	aggttttatt	caacgcttat	gtttattatt	aatcttcaat	960
cttggtatgaa	acccttatgaa	tgttaagcaat	gcgggaagcg	ttttagagct	catcttcaac	1020
ttcagaacata	tcagaaaatt	catgttggct	tgaacccttc	tgaatgtaag	ggatgtgggg	1080
aaacgctttag	attatatcga	catatgtgtc	tgcatacaga	aattcatcat	ggtgtgaaac	1140
cgactcaaatg	taagaagtgt	ggaaaggctt	ttgtgtcatg	tccaagttct	taccaacata	1200
agaaaattca	tctctgtgtg	aaaccatata	aatgtgaaca	atgtgaaaag	gcctttgttc	1260
gcagctatct	actttgttga	catcaaaaga	gtcactatgg	tgagaaacct	catgaatgca	1320
tggaaatgtga	aaaggctttt	agtaagggtc	caagccttct	taaacataaa	agaattcata	1380
tgtatgcgaa	actctatgat	tgttaaggat	tgtggaagcg	cttttttaga	ggctctcaac	1440
ttacacagca	tcagagaatt	catactgggt	agaagccaca	tgaatgtaaa	gaatgtggga	1500
agacttttaa	gctttcttca	tattcttatt	aacatcagat	aattcatcat	gattttgaagc	1560
catatgaaat	taagcaatgt	gggaaagcct	tcagtcgtgt	tggagacctt	aagacacatc	1620
aatcaattca	tgtctggggag	aaaccctatg	aatgtaagga	atgtgaaaaa	gaatgtgcta	1680
tttaattctca	actaatttat	catcagacaa	ttcatactgg	tttgaaaccc	tatgtatgta	1740
aagaatgttaa	gaaggccttt	cgcttctatc	caggcttctc	tcaacataat	agaattcata	1800
tggttgaaaa	accctatgaa	tgtaaagaat	tgataaagcg	ctttaatcgc	agtgatcgac	1860
ttactcaaca	tgagacaatt	catactgggt	tgaaccacca	gaaatgcmaa	gaatgttgta	1920
aggccttttag	tcatgtctat	caacttagtc	aacatcaaat	atcttaccat	ggtgagagac	1980





gaaggggtgg	gctttgcgaa	gatggcgggc	ctgggggtgc	tggagctcga	cctgccaa	180
gccgtgacac	tcttgaaaaa	tctccaggag	caagtgtatg	ctgtaaactg	acaagtga	240
tcactgacac	aaaaagttca	agctgtgtgc	tatcctacag	aaaaggggtc	cagcttcttg	300
gaagtgaagg	accagctgct	gctcatgtac	cttatggatt	tgaccacact	catctctggc	360
aaagcctcag	gaggatctct	tcaggagcat	gatgcagttt	tgagacttgt	agagattcga	420
acgggttttg	aaaagcttcg	tcccttggac	caaaagctga	agtatacaat	tgacaagctg	480
atcaagactg	cagtgacagg	cagccttagt	gagaatgacc	cacttcgttt	taagcctcat	540
cccagcaata	tgatgagcaa	gttgagctct	gaggatgagg	aggaagatga	agcagaagat	600
gaccagcttg	aggcttcagg	gaagaaatct	gtgaaggagg	tgtctaaaga	atatgttcc	660
ccacgcttgg	tccagtcaca	ttatgatgaa	acagaagctg	agcgggagaa	gaagcgtcta	720
gaacgagcca	agagacgggc	attgagcagc	tctgtcatc	gtgaacttaa	ggagcagtag	780
tcagatgctc	cagaggaagt	ccgtgatgct	cggcatcccc	atgtttaccc	ccagagtcag	840
gaggaccaac	acaggattaa	ctatgaggag	agcatgatgg	tgcgttttag	cgctagtaag	900
cgagagaaag	gacggcgaaa	acgagcaaat	gtcatgagct	cacaacttca	ttcccttaca	960
cacttcagtg	acatcagttg	tttgacaggg	ggaactgttc	atcttgtatg	ggatcagaat	1020
cctattaaag	agcgcaagaa	gatacctcag	aaaggttcga	agaaaaaagg	ttttcggagg	1080
cgcggtgtat	tatgggtgtg	catatttgtg	tattttttgt	catcctgaga	tacttcta	1140
ttcatttgtat	atagttgtgt	ttccctggaa	ttcataaatt	gtttgtcttg	gacatgttga	1200
aagagcctta	ctaataaaat	tgattttact	tatgaaaaaa	aaaaaaaaa		1249

<210> 87  
 <211> 1064  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 1441770CB1

<400> 87						
cggtctccca	gagcgtctgt	aaacacccag	agactgtcat	ggagggggag	gaggaggcgg	60
cggcggcgaa	gggaggcggt	tggggccgcc	tcagggtcc	gctctgccat	tctgtaactg	120
gtccctcgte	cccgtgacac	tgccatcagg	gaagcgaaat	gttagcgag	agcaggaggc	180
agccagaaact	atatccccc	cttccctggg	gcggggggcg	ggccaggccg	gctgagccgg	240
gggaggggctc	ggggaggggg	tgccctggcca	ggccggccctg	ctgcgcgcga	tggatgacag	300
taaggtgtgtt	ggaggcgaag	taaaagaagcc	cggtaaacgt	ggtcggaagc	cagccaaaaa	360
tgactttgaaa	gcgaacacttg	agaggagccg	gcagagtgtca	agagaaatgcc	gagcccgaaa	420
aaagcttgaga	tatcagttat	tggaaagatt	ggtatccagt	cgagaagaag	ctatatgtgc	480
cctcagagag	gaacttgga	tgtacaagca	gtggtgcagt	gcaatggacc	agggaaaaat	540
ccctcttgaaa	ataaaggccc	tactcactgg	agaagagcag	aacaaatctc	agcagaactc	600
aagcaggcat	accaaggctg	ggaagacaga	tgctaatagc	aattccttgt	gaagattata	660
taaaagctgat	tcagtgattg	aagccaatat	tctgattccc	atggaagatg	gatggggca	720
agtgtactctc	ttggctccat	ttactacctg	ctgctcagta	aatctctctg	taaatctgtca	780
atctctacca	aaatgtgtga	tcgtagatct	caaaaggtat	tgctttaact	ttcaacactt	840
agaaaattcta	caaacattca	gacctgtctg	ggtttgtatt	gccaccocat	acatttaaca	900
tgtgtgtgatg	cttgaaaaca	caggagttaga	gaaaatcgat	gaagattgtg	tttttgcacc	960
ttaaactccac	atgtctttat	tggttaattt	atattcttct	catgtaattc	atgtaattgt	1020
atgtctgtgtg	gtgttttatg	tgtcaccacc	tttcatgttt	ttga		1064

<210> 88  
 <211> 1398  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 1456684CB1

<400> 88  
 agacccaaaa gcattaagga gaaaaagaaa actacatcac ataccagggg agaaataacc 60



```

gctgcagccca tagctacgtg cgttcgctac gaggattgag cgtctccacc cagtaagtgg 180
gcaagaggcgc gcagggaagtgc ggtacgcagg ggcgcgaaggc gcacagctgc tagacgactc 240
gctttccctctc cggcccaacctc ctgaagccgc gtctctacttt gacagactgca gggccgcggc 300
ctgtgtctctct ctgcttccaacc atctacataa tgaatccacc tatgaagcag aaacaagaag 360
aaatcaaaaga gaataataaag aatagttctg tcccaagaag aactctgaag atgattcagc 420
ctctcgcatc tggatctctt tttggaagag aaaaatgagc gtccgcagcg ttgtccaaaa 480
ggaaacatcg gaatgaccac ttaacatcta caacttccag cccctggggtt attgtccagc 540
aatctagtga aaataaaaaa ctctggaggag tcaccacgga gtcatttggat cttatgatta 600
aagaaatccc atctctctcag tattggaagc aagtggcaga aaaacggaga aaggcgctgt 660
atgaagacatc taagggaatat gagaaacttc ataaagaaat tgaacaaaag gacaatgaaa 720
ttgccccctc gaaaaaggag aataaagaac tggcagaagt agcagaacat gtacagtata 780
tggcagagctc aatagagaga ctgaatgggt aacctctgga taattttgaa tcactggata 840
atcagggaatt tgattctgaa gaagaaactg ttgaggattc tctagtggaa gactcagaaa 900
ttggcacgtg tgcctgaagg actgtatctt cctctacgga tgcacaaagg cactatgaa 960
atgcattaatc atttgactgt tgagaatttt actgcggaag ttaccctcca ctagtctctt 1020
gtacagcagtg acataactac ataattgcaa ctctggaatc aaatttccct gtttgaatcc 1080
ggggaccctca ttctcataaa gtacaaatc tatgtatctt taactatgaa tgggttatgt 1140
gaatagagatt ttctcagttg tcagccatga cttatgttta ttactaaata aacttcaaac 1200
tctctgtgaa cattgtgtat aacttagaat aatgaaatat aaggagtatg tgtagaaaaa 1260
aaaaaiaaaa

```

```

<210> 91
<211> 943
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 1968191CB1

```

```

<400> 91
gtcccatcttc cagaaatcac aaggatgtta ggcaatgaat ggagtaaaact gctcctctgag 60
gaaaaacagc gctacacctga tgaagcagac agagataagg agcgttacat gaaggaactg 120
gaacagtatc agaaaacaga ggcctacagc gtctctcagta ggaacaccca ggcacgctcag 180
aaaggcaaat ctcataggca agatgcagcc cggcaggcca ctcatgatac tgagaaagaa 240
acagaggtaa aggaacggctc tggttttgac atccctatat ttacagagata attcttgaac 300
catagcaaaag ctcgggaagc agagctccgc cagctctgca aatccacatc ggaagtctgag 360
gagaggaatg cagccctgca aaagcacgtg gagagcatgc gcacagcagt ggagaagctg 420
gaggtggatg tgatccagga gccggagccgc aacacagctc tacagcagct cctggagacc 480
ctgcggcagg tgctgaccag cagctttgcc agcatgacct tgccctggaag tggagagaca 540
cctacagttg acaccattga ctcatatatg catagctaca gttcgagaag ttgtgaacag actcgatcgt 600
cccacagaca atgaaaaattc ctgataatga tttatatgcc agcacatcat caaaaaaag 720
taggggaatg atcttatcaa tactactaaa accctgggat tacatggat gaacaaagt 780
atgtcatcag gacagctggg taagattctc gttgcattgt tgttaaatgt agtaaaatat attagaaaag 840
agaatcactg tagtcccgac tacttgggag gctgaggttg gaggattgct tgagcccgac 900
agttcagtc cagagagatc ctgtctctaa aaataaaaga aaa

```

```

<210> 92
<211> 1997
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 2052061CB1

```

```

<400> 92
cgaccacgcg tccgctcgcg tggggcgaga ggtaccccg ggggcagcta ggtgctccca 60
agaaaccccg ccccagagga cccgcacgag ttgttgccat tttctgta agccctgcgc 120

```



WO 00/44900

PCT/US00/02237

ctgccagtg	cggtccacac	tcattgtacac	ttatgtcttc	gctttctacc	tcaaaaagaa	1200
taaccagtc	attatctttg	agaataacca	agcagatcta	gagaatgcga	cagaggtgct	1260
ctcgggctac	cttgaacgag	atatttccca	agattctctg	caggatataaa	agcgaaaggt	1320
acaaagacaag	tacagatact	gtgagagtcg	acgaagggtt	ttgttacagc	atgtgcacga	1380
agggtatgaa	aaagatctctg	gggagtagac	tgaggactga	gaatggccct	gcataaaaatg	1440
aaactctgaaa	actttaccat	ctagagtgc	catgcaattt	aaacaaaaca	aaacaaaaca	1500
aggaggacat	aagcctattc	tgacaccact	ggtctgtagt	accagaattg	ttttgtttaa	1560
ggaaagtctt	agttaattat	atgttaataa	aaaggttagat	aacacattgt	acaacagtat	1620
cttagcgccg	caacaaaagt	gtgacagaca	cacataaaagc	cctccaactt	taactgttaa	1680
cgtagcttca	ttctcaaaag	tgactccttt	ttttcttttt	ttcttttctt	gagtgtagta	1740
cagttaaaa	ttcaaacagc	tccttgacac	tgcttttcat	gttcaaacca	gccattttgt	1800
tgtagctttg	taaaggacct	cttccccttc	ctcccctaca	catcacagata	cacccacaca	1860
cagactgact	ctctttctct	cataccacca	ggtcatgagt	gaatgatgct	taagtctctt	1920
taaaagaaat	cttgggagtg	ggaaaggggt	aggcagcaag	aggattcaac	cacccagaaa	1980
catataaac	ttgtatatga	cttttaaaac	aagaggacaa	cacagtat	ttcaaaaattg	2040
tatatagcgc	atatactggg	acaaagcaag	cggtggcagct	gtttgcataa	tggtttactt	2100
caaaaaatga	tttatctctt	aaaaatcttc	aagattatgt	ctatttgcct	tgcatctttt	2160
ttcagtttgc	ttatctttcc	cggtttgggg	ttgggataaa	gggtgtctgg	tttagcagct	2220
ctggaagctc	tatctagagc	tccttcaact	tcctgaggtt	attttgccct	ttctgggtgt	2280
ggtagtgctg	tggtggccca	gcctctctct	gccttgaatt	cctgctcttg	atcaggggaca	2340
aggggagctc	agctctgact	aatgccaatga	cctgattaa	gggtgacagca	gggagttttt	2400
tgctacagtc	tcatgaatga	acctgtccca	acctaatccc	ctccatggc	atcatgcctc	2460
taccacaagcc	tttgtgtgcc	catgttatgc	acacagctgc	aggcattctt	aagtcacctt	2520
tcgcataccg	tggaagcatt	ttaaaatttc	ttttactttt	tggttttccc	ttaattgtctg	2580
cttttccagat	tttagttatg	gctcgtctgc	tcaccccttc	tcacatttag	gggtgtcaag	2640
agaaagtgtt	gctttaaata	taaatagcca	ttcatttagt	ctcagttagt	tgctgttaaa	2700
tggttggtac	cgaattgtct	tgtgtgtgtt	gctgtgggtt	tggtttgaag	gcacacaccc	2760
ctagaacatg	atatctccat	ctagtgcatt	taaatagaaa	tgctgtgatt	aactatatac	2820
ttattgtcag	cagataggag	aatataaata	gcattttagc	tgtgatgtcc	atttttatga	2880
aatctctact	aagagctatg	ttaaaagtaa	aggatggtgg	tggttgatgt	aactatatac	2940
ctgttttagc	cattctgggt	gtggtatttt	tcaataggct	agcatctgta	aactctgcag	3000
ttttatagc	gagtgccagc	tgaactaggc	aactagatta	agaggtctaa	atatgaaaaa	3060
ccagttgagc	ctggggacct	ctctgctctc	cttttagatg	cttttgccca	ggggagtttt	3120
accattttgtg	aggcagcttt	gtctgctctt	acacgttaca	tccttattact	ccatggggaa	3180
gtagagttcag	tttctctctg	ctctttgcct	aagttaggct	ttgctgtaac	aacctcattt	3240
ttctctttac	aaaaggttgt	tacaggagat	ttactggcaa	ctgtttctttt	cccatcaaaa	3300
atcagttgat	gtttgtctgag	tataaaatgct	gcttcccttaa	accacttgtc	gctttaggat	3360
caacttttacc	tgtaactttt	ctctcttctc	cccttgccac	ctcaggtgca	aatctgaact	3420
cagttgtctgc	ttcttccatt	ttctctcttc	ttctccctct	ttccccaata	ttccattgac	3480
attattttac	ttcaaatgac	agcatcaatc	ttaaaaagat	atacattaaa	actaaggagt	3540
tttttttaag	aaagcctgaa	taagttctct	ttccctggtaa	ctttgaaaaag	cagtcagagt	3600
tgctatatag	atatatgtgg	ctcttttaaa	atgcttttgtg	tatgtgttgtg	gttttaaaaa	3660
aaaaatctta	ccagtttaact	ttgcagtgct	taggtttttag	tgctataaat	ccacgtgttc	3720
ctgttgcaac	aaatctggtg	aaattgtgtg	tgacacttgc	ataaccagtc	ttccacctg	3780
gaggaaacag	gcttttttaga	gatgctttct	atttcaatgt	tggaactactg	cttgagggta	3840
ttgcagttgt	gggtgcaact	cctaatttgt	atttcaatgt	cttgaggtg	ccttgcactg	3900
ttcaagctct	tcaacagatac	caccataatt	gacagaaatc	ccagagtgaag	ttctgttgtgt	3960
tattagttaga	taagctgccc	ccagctggga	taggcaagcc	atgacagctt	cctgttttca	4020
cttacagaa	tcttatctga	gggatctatt	cacagtaagc	accaaggttc	ccatgttttg	4080
aggtcagttg	cattgtcttt	tgaaaaagtc	atgcttctat	tgaaacaattc	attcagcagc	4140
agatggactt	tcagtgattt	aaaaataaat	tttgatccaa	agctcaggac	acaaacccca	4200
gtgttgtaaat	tgagtgcatt	ataataatcg	actaaatat	ctgtaaat	ccacacacca	4260
gattgtatgt	gtttttatgt	tgtttaataa	aatatgttag	atacacagctg	atacatcac	4320
ccatatagcag	caga					4334

<210> 94  
 <211> 1706  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc-feature

WO 00/44900

PCT/US00/02237

<223> Incyte ID No.: 2101803CB1

<400> 94  
 ccgcgcgccg ccgccaccat gagggccgag ggcctcgccg gcttggagcg cttctgcagc 60  
 cggggcaaaag gccgggggct cggggctctg cagcccttcc aggtggggga cttgctgttc 120  
 tccctgcccg cctatgccta cgtgctcacg gtcaacgagc ggggcaacca ctgcgagtac 180  
 gtcttcacca ggaagaagag attgtccaaa tgtggaagat gcaagcagcg attttactgc 240  
 aatgtggagt tcagagaaga agattggccc atgcacaagc tggaaatgttc tcccatggtt 300  
 gtttttgggg aaaaactggaa tccctcgagg actgtaagac taacagcaag gattctgtgcc 360  
 aaacagaaaaa tccaccacga gagaacacct tcggaaaaaa tgttagctgt gaaggagtgt 420  
 gaatcacatc tggataagtt agacaatgag aagaaggatt tgaattcagag tgacatagct 480  
 gctctccatc acttttact caagcatctc gaattccctg acaatgatag cctcgtagta 540  
 ctctttgcac aggtttaact taatggcttc acaattgaag atgaagaact tctctatttg 600  
 ggatcagcga tatttctctga tgttgcatgt atgaatcata gctgttgccc caatgtcatt 660  
 gtgacctaca aaggggacct ggcagaagtc agagctgtac aggaatacaa gccgggagag 720  
 gagggttttta ccagctatat tcatctcctg taccacaagg aagatagaaa tgaccggtta 780  
 agagattctt atttctttac ctgtgagtgc caggagtgtg ccaccaagga caaggataag 840  
 gccaaagttg aaatccggaa gctcagcgat cccccaagg cagaagccat ccgagacatg 900  
 gtcagatagt cagcgaacct cattgaagag ttccggaggg ccaagcacta taaatccctc 960  
 agtgagatctc tggagatctg cgagctcagc caggagaaga tgagctctgt gtttgaggac 1020  
 agtaacctgt acatgttgca catgatgtac caggccatgg gtgtctgctt gtacatgagc 1080  
 gactgggaag gagccctgca atattggacag aaaaatcata agccctacag taaccgctat 1140  
 ccttctgact cctcacaact ggctctcatc tggttgaagc tagggagact ctacatgggg 1200  
 ctggaacaca aagccgaggg ggagaagaag ctgaagaagg ccatgtgcaat ctggaagta 1260  
 gctcaccgca aagatcatcc atatatattc gagatcaaac aggaataatga aagcactga 1320  
 aacctgtcag catttcatgt ttcatttaaa cacttagttc agaaaactta aaggaattga 1380  
 atatttcaaa ttgcacacct cactccagca tctctgtaaa ataatitgaa tgaataact 1440  
 tcttgcactt aaacactgca catgcgctac tttagaggtt tttgtaactc tgactcttaa 1500  
 taccaaatct attttgaatg cttttgtttc ctaagagata atggcattgt ttcattatgt 1560  
 atacttttga cagacagagt tttaaaaagt gaattatttt ttctttctac tctcttgtaa 1620  
 tgttctgaac aaacttgaat gatgaagata ttaagagat atcagtaaaa agaacaaaaa 1680  
 ataaagatcc agaaagaaaa aaaggg 1706

<210> 95

<211> 1602

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 2112362CB1

<400> 95  
 ccgtggggca gtcgaggat tgcgtgaatt acgcggcggg gctgtcgccg tacgcggaca 60  
 agggcaaggt cggcctcccg gagatcttcg acccccggga ggagctggag cggaaaggtgt 120  
 ggggaactggc gaggtgtgtc tggcagttct ccaatgttgt gtccacacag ggtgccggga 180  
 tcagactgac cctctgcata ccgaacttca ggggttccca cggagtctgg acctggaggt 240  
 agcgaggtct ggcccccaag ttgcacacca cctttgagag cgcgcggccc accgagaccc 300  
 acatggcctg ggtgcagctg gagcgctgg gctcttcccg cctcttcccg agccagaagc 360  
 tggcagggct catgtgtgct tcaaggtctc ccagggacaa actggcagag ctccacggga 420  
 acatgtttgt ggaagaatgt gccaaagtga agacgcagta cgtccgagac acagtctgtg 480  
 gcaccattgg cctgaaggcc accggccggc tctgacacct ggctaaggca agggggctgc 540  
 gagctgcagc gggagagctg agggacacca tcttagactg ggaggactcc ctgcccgacc 600  
 gggacactggc actgcgcact gagggcacga ggaacgcga cctgttccat acgttgggtt 660  
 catcgctgca gatccggccc atgcgggaac tgcgctggc tccagagcgc cggggagggc 720  
 gctgtgtcact cgtcaacctg cagccacaca agcacgacgc ccatgtctgac ctccgatccg 780  
 atgtgtcact tgcagaggtc atgacccggt tcatgaagca tctgtggctg gagatccccg 840  
 cctgggaagct cccccgtgtg ctggagaggg cgctgccacc cctgcccctg ccgccccacc 900  
 ccaagctgga ccccaaggag gaattctcca ccgggataca cggctctatc cgcgcggccc 960  
 ccaagcagga gccctgcgcc cagcacaacg gctcagagcc cgccagcccc aaacggggagc 1020  
 ggcccacagg cctgcgcccc cacagacccc ccaaaaggtt gaaggccaag cgggtcccca 1080  
 gctgaccagg gtgcctgggg aggggtggggc tttttgtaga aactgtggat tctttttctc 1140

```

tcgtgggtctc actttgtttac ttgttttctgt ccccgaggagc ctcagggtctc tgagagctgt 1200
gctccagggcc aggggtttaca cctgcccctcc gtggtccctcc cctgggctccc aggggcccctc 1260
gggtgcgggttc cgggaagaag ccacaccccca gaggtgcagag ctagagcccctc gccacacccc 1320
agccctctgctc ttgctgtgttt gtccagaggtt gaggtctgggc cctccctggt cctcagctta 1380
aacaggaggtg aactccctctc gtccccagggt cctccctctct gggccccctca cagccccacc 1440
taccctctctc ccatgggtccc tgaggagggt gagaccacc cttaagttggg ggaatcagtag 1500
aggcttgacac tgcctttggg gctggaggga gacgtgggtc caccaggctt ctggaagaaat 1560
cctcaatgca ataaaaacaa tttctttctt gcaaaaaaaa aa 1602

```

&lt;210&gt; 96

&lt;211&gt; 1951

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 2117346CB1

&lt;400&gt; 96

```

gaggcgccgc cgaccggcgg cctcttggcg cggattaggg ggtctcggcg agggagctcat 60
caagctttgg tctatgtgtt ggccggttct agaagctctga ctagggaaga 120
ccaaagcagc actcgtttgcc aattaggga tggaccgttt ggtgtccctt agcaatgac 180
cctctgataa ccacaccttc cgaggctgct cctccctacc catggagcct tataatcaag 240
gtgctgaatg tgggccacct ccttttttcc tctgcttgca gtgtttcact cgaggctttg 300
agtacaagaa acatcaaagc gatcatactt atgaaataat gacttcagat ttctctgtcc 360
ttgatccccg ctggactgct caagaagaaa tggccctttt agaagctgtg atggactgtg 420
gcttttgaaa ttggcaggat gttagccaat aaatgtgcac caagaccaag gaggagtgtg 480
agaagcacta tatgaagcat ttcacataa accctctggt tgcactacc ctgctgaacc 540
tgaacaacgc agaggaagca aaaactgtcg acacagccat tccatttacc tctacagatg 600
accctccccg acctaccctt gactccttgc tttctcggga catggccggg tacatgccag 660
ctcgacgaga ttctcattgag gaatttgaca attatgcaga atgggacttg agagacattg 720
atctttgttg agatgactcg gacattttac atgctctgaa gatggctgtg gtatgatattc 780
atcatctccg gtttaaaggag agacaaagac gaaaaaaaat tataagagac atggatttaa 840
tcaaccttag aaagtttcaa ttaattggaac ggcggtatcc caaggaggtc caggacctgt 900
atgaacaacat gaggcgattt gcaagaattg tggggccagt ggaacatgac aaattcattg 960
aaagccatgc atttggaaattt gaaactccga gggaaatcaa gaggctccaa gaatacagga 1020
cagcagctatc taccaatttt gttagtgcca gaacctcaag tcacctcaag aagacacggg 1080
aggaagagcg ccttaaacgc actatgctct cagaagtctc ccagtatatc caggacagta 1140
gtgcttgcca cgcgtggctc ctgacattga ttcggccctg agtccctcca 1200
ttccaatggc ttcgaattca ggtagacgga gtgcaccacc cttgaacctc actgcccctc 1260
ctggcagcga gaaaaagaaa aggaagctctg tcaagctgtg aggttgggtc ttgtattatg 1320
ctggagccta tttagaatac aaatctgctc tattgaacga atgttaacaa caaggaggct 1380
taagacttgc gcaggcaaga cgaactcatc agatagatgt gaacaaacc ccgaaaatct 1440
atgatttctc catcagagaa ggtacatcac ctaaaaggcta aggcctcaag agcttgggat 1500
cagaagtctg aagtcttgaa tgtgggtggg taagggtggg gaaaccttaa 1560
agtgtgtttt cagctgaaat ctcatggtga aaacagggga aaggacaaag gaaaccttaa 1620
gtgttattgt cttctttctt cttactctct ctcttaaaaca cttcttgggt gttgtattatg 1680
ctgcagagtt gtgtgtcata taagctatta ttaaatgtga gtgggcatc attctcaaca 1740
cctcttgttaa ctaaaagacc catagtaccc tcacatcaca gtctctggga gaaattggag 1800
ctaaacccac agtctgtaaa tccaggaggt tcaagcctag aatccttaat atgtccggg 1860
gaaaggtttt tctgaataaa acacggcttt ggccttaaaa aaaactttgg ggaacagaa 1920
tttaaaaatt aaaagggaaa aggggttttt a 1951

```

&lt;210&gt; 97

&lt;211&gt; 854

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

<221> misc-feature  
 <223> Incyte ID No.: 2119917CB1

<400> 97  
 cccacgcgtc cgcggacggt gggcgcgccg tatgacggcc agcgacacgc cgcgcgggag 60  
 cgggcccagga gtccgagtcg gaggctcggt gaccacgtgc aagcatccgc gttgcgtctc 120  
 cctggggaag aggaaggcgt cgggtggagc gacacccctg gaggccatct 180  
 ggagttcggg gaaacccctgg aagaattgct tcaaggggaa acctgggaag aagcagctct 240  
 tcacctgaaa aatgttcaact ttgcctcagt tgtgaattct atcttgaga aggagaatta 300  
 ccattatggt actatatata tgaaggagaa agtggatgtg actcatgagt cagaacaaaa 360  
 gaattgtagc cctgaaaaaa atgaagggtt ggagtggtgt ccttggtggg aactacctcc 420  
 cctggaccag cttttctcgg gactgcgttg tttaaaagaa caagcgtatg atccatttaa 480  
 agaagatctt aacctctgg tgggatacaa aggaatcat ctctagggtg ccgagaagat 540  
 ttgattttct taaaaagac aagaataaagg tctggttagg gaatgaaaaa tgtatacatt 600  
 tcgggaacca tccattttat ctaaaaaagt tcttggtatt gccagtttat ttgcagcttc 660  
 ttaattgtat cccactctt tcagccagta cttgagaaaa tttttctgaa atatgtcatt 720  
 gaattgtatt ccagacacag aatacatgat aaatactgat attatgggta atctgctttc 780  
 catattttacc tatggatatg tacgtgcaat gtgccataac tagttgagag ggtgggtgag 840  
 gcagattttc tctt

<210> 98  
 <211> 1581  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc-feature  
 <223> Incyte ID No.: 2123456CB1

<400> 98  
 tccgggaaag tttctttgga ggtccggccc ggagcggcca tgtcccacgg ccccaagcag 60  
 cccggcgccg ccgcgctgcc ggcggggcgg ggcagctcgg gccagcatgg gggtctcgtg 120  
 gtgactgtca agcaagacgc cggcgagggt ccacgcgcgg gcgagagagg gtcccacagag 180  
 gaggagcccg gttctgcagc acggtgaagaa acgcggtcgg ccaaggggca agaagcggaa 240  
 gaagattctg ccgaatggcg ccaaggcacc ggtcacgggg tacgtgcgct tcttgaacga 300  
 gcggcgccag cagatccgca cgcgccacc ggatctgccc tttcccgaga tcaccaagat 360  
 gctgggcggc gaggtaggca agctgcagcc aacggaaaaa cagcggtagc tggatgaggc 420  
 cgagagagag aagcagcagt acatgaagga gctgcggggc taccagcagt ctgaagccta 480  
 taagatgtgt cgcgagaaaga tccagagaaa gaagatcaag aaagaagact cgagctctgt 540  
 gctcatgaac actctcctga atggacacaa ggggtggggg tgcgatggct tctccacctt 600  
 cgaattgtcc atcttcaact aagagttctt ggacccaaaac aaagcgcgtg aggcggagct 660  
 tcggcgcttg cggaaagatga atgtggcctt cgagagagcag tctggagcag gactggtgag 720  
 caccgagagc atgagcagcg ccgcgcgagcg tctggagcag gactggtgag tggagagagc 780  
 gaggagctgt gcgctgcagc agcagctcca ggccgctgac caggcgctca cgcgccagct 840  
 gcctcactg ccggtgcggg gcacggggcga aacgcccacg cttgggcactc tggacttcta 900  
 catggcccgg cttcacggag ccacgcagcg gcaccccgcc cagcacgaga agctcatcgt 960  
 ccgatcaag gaaatcctgt cccaggtcgc cagcgagcac cgttgaggag tgggcggggc 1020  
 cagcatgcag aggagaagct gtggggcgcg cccctgccca cccaccccgc tggacgagag 1080  
 gctgggggtg tttctgcagc ggcctgggtc atctctgcac cttgggggct ccagccccc 1140  
 taaaattaaa tttctgcagc atccctttag ctttcaatct ccccagcccc tggacccggg 1200  
 aaaaagcact cgctgcgcga tacaccacga agaacctcac agccgaggtg gccctctc 1260  
 ggagagcagc cagcgctcac actggtctct cggggcaccc ccagacaca gtcgagacga 1320  
 aaccaccccc cagcacacgc caggaccccc caaatctact actacggggg gctgtgccat 1380  
 aggcacacac ggaagctgcc ttgtggggag ttacotgggg tgtccccgcg atgctgtgac 1440  
 ccagatgggg tggggcgccg ctttgcctac cctgctctcc tcagcccgag ggacccctgt 1500  
 ggggggtgct ccttctcact gctggatcgg gactttttta ataaaaacaa gtaaaatttg 1560  
 tgtttttaaa aaaaaaaaaa a

<210> 99  
 <211> 2150





```

tacctcgttta ggcacagctc gcacagaggt tgtctgcaag cagtgtgaag ctcatctagg 480
tcacgtgtttt cctgattggac ctgggcccaa tggtcagagg ttttgcatca acagtgtggc 540
tttgaaagttc aaaccaagga aacactgacc atcttcaaga gtcccggttc ctggccacc 600
cttcagctgc accctcaatt cccactgaat acttgaatga cttgttttat ttgcaataaa 660
actgggctga atttgcaaaa aaaaaaaaaa a 691

```

&lt;210&gt; 101

&lt;211&gt; 2101

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 3128913CB1

&lt;400&gt; 101

```

gtggccttga gctcgggggt ggtggctcat tctctggcgg cctcctgggt tcgcggaag 60
aagagattac tcacactcct tctcaagcac agaaccagtt gtactgagct ttttgctaa 120
ctgttttcac caagaaatgc ttgggaatct ggagtgattt caacctgat acctcaggat 180
cctccgggaa aagaactaat actagtgaat gtagaagata acttttctct ggtatgagaa 240
tttaagcaga atcgagatgc tcaactcctg cagaatttgt ttcgtcagca atcagaaaaa 300
ttttgctacc agggagaccc tggggccccc gaggctctgt gccgactcca ggaactttgc 360
tatcagtggc taatgccaga gttgcacaca aaggagcaga tcttagaact ctggttactg 420
gagcagttcc tgagcattct gccctgagg gctgcagatc ggggtcagca acataatcca 480
gaaagcggcg aggaagctgt gaccctgttg gaggatttag agagggagtt tgatgacca 540
gggcagcagg tccagctag tccacaggga ccagcagtcg catggaaagg ttaacatgt 600
ctcagagcat cccaagagtc aacagacatc caccctcagc ccttaaaagc acagctgaaa 660
tcttgaaacc catgccttcc ccttaaaagt gatttgtgag acagtgaacc agcaacaaaa 720
gagggcatct cagaagaaaa atcacaggga atgtgtggaag caaggaaagt ctacagggga gaaactaaga 840
agtgagcatc aaagcaatgc ttttagtcaa gtgatcttca caaacaatct ctaggaaagg 900
agagaccttt atgatgaggc tgaagatgac ttgattctaa ctacagactc tataatgtgt 960
cagaaaagtc ctcagaaga gagaccttat agatgtgatg tatgtggggc tatgttcagg 1020
cagcatttct ctctaacaca acatcagaga atccatactg gatgaaaagcc ctataaatgt 1080
aaccagtgtg ggaaggcctt atgtagttaa tgggtgggaa tcttatctta ttattcatca gagaattcat 1140
agtggttgaga aagcatatga cctactgggt tttagccaga gataaagcct ctttcaatca gactcagcc 1200
ctcattagac atcggaataa atatctcatt tttagccaga gataaagcct ctttcaatca gactcagcc 1260
aaagcattca gtcaaatgct gagacacctat gaatgtaatg aatgtggaaa agctttcagg 1320
ccttagagtg gtaatgaaatg tcatcagaga atacatagtg gagagaaaac ctatgacat 1380
cagcagctcag agctgtattac cagtttgagc tcaaacctta tcagacatca gagaattcat 1560
agtgaatgtg gaaaagcttt gtgtaatgaa tggggcaaaa ctttcaaaag gagatcagcc 1620
agtggtgggg aacctttatca tcaattctgg gatgaaagct atatatgtaa tgaattgtgg 1680
cttgttcacg atcagagaa ggtccttatg gcccatcaaa gacttcacac tataaagttaa 1740
aaggctttca ggcacagatc tgttaataat tcaagtcagat ttttaagttt gttagtcaaa 1800
agagttttact ttggagcaaaa atccataaaa ggttataaaa tactagttct tgactctagc 1860
ttgctttgtg cagcattttcc cagtgcataa gttaaagtgt ccttgaaagc ttttctgtgt 1920
actaatcaga acagaataca gaagaatcat tacttccagc tcttctctta tttagaatca 1980
tcaggaaata cgaaaagttg gaatgtaaca ttgaaacctc attttgtatg aaagtgtcat 2040
gaatataaga acccaggctc tgtcactgca tctgattgtt agtgtgccag gttatggggg 2100
g 2101

```

&lt;210&gt; 102

&lt;211&gt; 2196

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 3282941CB1

<400> 102  
 cctactttct ctagaccaga aattatccaa aattgtgcac attagaatca cgtgggtagc 60  
 ccacaaaaat cccagtgcca aattataacc tgcctttcct gtggcccagc ggtttccaac 120  
 gacatttgggc cctcggctctg ataactactgg ctacatcgct cctcaggtcg 180  
 aaccttggacg gtggcgagtg ttgcaaatat ttccgcggt gcgcgcggat tctggacttg 240  
 ggccgccaact cgtagtccac gctccccggg gtcacgagag gggcgtaacg cctctgccac 300  
 ccacctcgct ttctcaccgc gcgcttcccc gcttgggttt ttagtcttcc ttggagcgct 360  
 ctctgtcgct cgctctcgcc agggagcgga aggcggagac agcgcagact gccagggggg 420  
 aggaaagagc acgcgtgtgg gcaaggggga caacgggatg tccacgggct cggtagtgta 480  
 tccggaggag atggagcttc gggggctgca cggggagtac ccggtccccg cctccaagag 540  
 gccgcccctc cgccgcttag agccgcagct cgctctgccc agtgacaact cgtcggcaga 600  
 ggaggaggac ccgcacgccc agggaggagc ctgcgctctg ggcacagccg gcagcgcgga 660  
 aggcctgcaag aggaagcggc cccgtgtggc tggggcgggc ggccgaggtg gtagcgcggg 720  
 cggtgtgtggc aagaagcccc tcccggccaa gggctcagcc gcagagtga cagcagtcga 780  
 gcgggaacgcy gccaacgccc gtgagctgct cccgtagcgc gtgctgagca aagcctctct 840  
 caggctcaag acagcctgc ctctgggtgccc ccccgacact aagctctcca agctggacac 900  
 gctccgctg tcttccagtt acatcgctca ctctggcgag ctgttgcagg agggaccgcta 960  
 tgagaacggc tacgtgcacc cagtgaacct gtacatggca tctgttgtct cgggaagacc 1020  
 ggactctgac accaaagaag ttctccgagc caacagacta tgttggaacca cgccttaaat 1080  
 cggagtggaa ctactctgat gggatatttc gttaaatgcy agtgtttggg aatggtagag 1140  
 agaaggagaga ctctgtgaga tgggaagaag ttcccgctgg attctccttg acccttcccc 1200  
 ttctcttggc actgtgatcg tgacaggtgg cgggtgtggc tgtcaactga cagcggccac 1260  
 gggtacagct gcgcgggatc tggggcgacca cgttttgcct ctccaaaagag agctctcttt 1320  
 cgtgacgaga cgcggcgaga ggtccaccct cgggcccctag ctctgtagac taactctcgg 1380  
 ctgctgcccc agcccgcgc agacagcccc cggatccgtt ctacagcgga cagagttcat 1440  
 cgcacacgtg gggcagcggt ccacacagcc cggcccttct cgggttgacac aatggttagg 1500  
 gaacggttag aacgcgctct acatccgctt taaagacaga ggtctagacy ttagatccgc 1560  
 gtcgggacag ggttttaagt gacaaaagaag ggcgagtggc ttctcttgca cgcgttcgta 1620  
 ctccagcaca gcgcccttct aacggggcggg aggaaggcgc ctgctcgcag ggttaggtgg 1680  
 agacacactt cccagactac ggcagcgagg tttaccggg agagcttcgg cccgttcgcy 1740  
 ctccctgcgc ggttgcttct tcaatccccg ctcttccca agctccccg tttttttaat 1800  
 caggcagcgct tctgtcaacc ctctccactt ctgggctgaa gcttccccaa gccccgctgc 1860  
 gccaacctgt gttgggttct ctctggggct cctctccgc ccgcgtctct ctctcaactg 1920  
 cagcaccccc agctccgact ccagactctc tgcatacagt ctcccactc cagcgtccgg 1980  
 gcgcccacaac tccaacacca cgtctctgcc cgcaggttct tcccgcgcgc gaggagcgcg 2040  
 caggggtgggc agctattgag agcaagtgat cctgcgtag ggaacgcgc cttgccccga 2100  
 gggtcgacta ccacaggaaa taacatatgt aaataaaatt atttttttat gaataataaa 2160  
 acgctgtgta aaaaacctga aaaaaaaa aaagag 2196

<210> 103  
 <211> 749  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> 735, 736, 741  
 <223> a or g or c or t, unknown, or other  
 <223> Incyte ID No.: 3286656CB1

<400> 103  
 gctgagactgt gctgcggcggc gcggcgcggc gcggctcgcc acggcgacgt gggagtgctt 60  
 ccggctggctg tgcaggggaga acaccgcagc agacctcaaa ccttggtctc agtgtcatgg 120  
 aatccgtcac ctttgaggat gtggccgctga agttcatcca ggaagtgggca ttgctggaga 180  
 gcgcacggag gagcctgtgc aaatacagga tcttgacaca gtgcagacc cttgcctcca 240  
 ggggaactcc accatgcaaa cccagtttgt tctccagctt ggggcaaaag gcagagccaa 300  
 aggcacagga ctcagggatt ctccgtgcca cagggtgtgc tgggaaatg caacttaaac 360  
 ccgaagagtt gccttctatg caggatcttt tgggaagaag atcctccagg gacatgcaaa 420  
 tggggccggg gctgttctct aggatgcagc tgggtccctc catagaaagc agggagacac 480  
 cattgactgc agaggacggc ccagctctcc agggagccgc ttggtctctg ggtatgacgg 540  
 gactgaagcg cgtattgcag attcagaggg tgggtatacc agtgcctact ctggggccac 600  
 gcaacccatg ggtggccagg gattctgcca tgtagcagct gctgtcttcc cctttgcctt 660

ccgcatgat tgtaagtttc tcgagtcctc ccagccatg cgtcctgtac aacctgtgga 720  
accaggcagc caaannncagt natagttgt 749

<210> 104

<211> 1311

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> 1294

<223> a or g or c or t, unknown, or other

<223> Incyte ID No.: 3490802CB1

<400> 104

gggcctttgt	ttctcgtgc	agcgggagct	ccagggttat	cctctgtgtt	ctgtgtcctg	60
tgcttataga	ggcccgctct	ctgtggcctg	gtgacctgca	agtattggga	gagccacagc	120
taaaccccg	gaccctctgga	agcctagaaa	tgggaccatt	gcaatttaga	gatgtggcca	180
tagaattctc	ttgggaggag	tggcattgcc	tggacactgc	acagcggaat	ttatataggg	240
atgtgatgtt	agagaactac	agaaacttgg	tcttccttgg	tattgtgtgc	tctaagccag	300
acctgtgtac	ctgtctggag	caaggaaaaa	aacctttaac	tatggaaa	catgagatga	360
tgcccaaac	cccagttatg	agttctcatt	ttgcccaaga	cctttggcca	gagaacatac	420
aaaattcttt	ccaaataggg	atgctgagaa	gatatgaaga	atgcagacat	gacaatttac	480
agttaaaaaa	aggctgtaaa	agcgtgggtg	agcataaggt	gcacaaagga	gggtataatg	540
gacttaacca	atgtttgaca	actaccacga	aagaaatatt	tcaatgtgat	aaatatggaa	600
aagtccttca	taagttttca	aattcaaca	catataagac	aagacatact	ggaataaact	660
ttttcaaatg	tataatatgt	ggcaaaagctt	ttaaacggtc	ctcaaccctt	actacacata	720
agaaaattct	tactggagag	aaacctttaca	aatgtgaaga	atgtggcaaa	gcttttaacc	780
aatctcaaaa	ccttactaca	cataagagaa	ttcatactgg	agagaaacct	tacaatgtg	840
aagaatgtgg	caaaagctttt	aactggtcct	cagaccttaa	taaacataag	aaaattcata	900
tggaacgaaa	accttaccata	gtgaagaatg	tgacagatct	tttaaatgtt	cttccacttt	960
taattagcat	aagataattc	atactggaga	gaaaccttat	gaatgtgatg	aatgtgggaa	1020
agcctttaac	cagccctcga	ctcttagtaa	atgttgaggt	ttatatggaa	acaaaacct	1080
acaaatataa	agaatgtgac	aaagcttttt	aagggaagttc	tcaaccttta	ttcacataaa	1140
ttcataccaa	acagaagccc	tacaagtgtg	aagaatgtgg	caaaacctat	aaacctataa	1200
caagttctca	atctcttttt	ttttgagatg	gagtttcaat	cttgtcacgc	aggcagaggt	1260
tgcagtgcac	actccagctc	aggcgacaga	gtangctctg	tcgcgacctc	a	1311

<210> 105

<211> 990

<212> DNA

<213> Homo sapiens

<220>

<221>

<223> Incyte ID No.: 3507366CB1

<400> 105

caaaaaaggaa	agaataaggc	aaattcttag	aatgtatgca	caacttaata	tccttgctcg	60
tttgatgtgc	attgatctca	tttaattagg	tttttgaaat	gaaaaattcaa	gaagggtgaag	120
ctcacacaat	tttttgcctt	gcataatgat	gcttccaact	tgtaacctgtg	gatcatcatag	180
aaagtgttagt	ttcaaaaggag	atggacaacac	gataactaca	gtttgatatt	caaggcctttg	240
ttgaaaaataa	tcctgcacat	aaatggtgtc	ctactccagg	ctgtgacaga	gcagtaagac	300
taacgacaaca	agggtcaaat	acatctggat	ctgataacct	cagcttccca	ttgtcgagag	360
ctctctctgt	tgattgtgga	aaaggacacc	tcttctgctg	ggagtgcctt	gggtgaagcac	420
atgagccttg	tgactgtccaa	acatggaaga	atgtgctgca	aaaaataaac	gaaagtgaac	480
cagaagaact	tgtgtggagt	agtgaagcct	acgaggatgc	cgccaattgt	ctctgggttat	540
taactaaact	caagccttgt	gccaaactgt	agtcctcaat	acagaagaat	gaaggctgtca	600
atcacatgca	gtgtgtctaa	tgcaagtatg	acttttctgt	gatttgcctt	gaagagtgtga	660
aaaaacatag	ttcgtccact	ggaggttatt	acagatgtac	tcgctatgaa	gcattccaac	720
acgtggaggga	gcaatccaag	gaaatgactg	tggaggctga	gaaaaaacac	aaacgatttc	780

```

aggaacttga cagatttatg cactattata caagatttaa aaacatgag catagttatc 840
agctagaaca acgcctctctt aaaacagcca aagaaaagat ggagcaaatg acgagagtct 900
caaaagaactg aaggaggctg tccagatacc acttttcattg gagatgagtt catgtgctct 960
aaaaatcgcc gcctctcaag tgtcttatca

```

&lt;210&gt; 106

&lt;211&gt; 1048

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 3573060CB1

&lt;400&gt; 106

```

ccgctgcagc tctccgcggg acatctcacc gttctggaga cagggctcgc tgcgtctcac 60
ggcttcttag gccgggggtg gacagccgcc ttccggccag aggggatgag gttgcgtcgc 120
gtctccgggag cgccgatggc gtgactggcc ccgcgcggag cagcgacact gcccgccgag 180
cccgtctctc tgcgccggag catgaatctc agtagcgcca gttagcacgga ggaaaaaggca 240
gtgacgacgc tgcctctggg ctgcgagctc agtcaggaga ggcggacttg gaccttcaga 300
ccccagcttg aggggaagca gagctgcagg ctgttgcttc atacgatttg cttggggggag 360
aaagccaaag agggagatgca tgcgctggag atcctgcccc cagcaaaacca ggaggacaag 420
aagatgcagc cggtcaccat tgcctcactc caggcctcag tctctcccat ggtctccatg 480
ttaggagtg cagctttctcc cccagttact ttccagctcc gggctggctg agggaccgtg 540
ttctcagtg gccaggaacg ttatgaagca tcagacctaa cctggggagg ggaggagtaa 600
gaagaagggg agggagaggga agagggaag gaagatgatg aggatgagga tgcagatata 660
tctctggagg agcaaaagcc tgtcaaaaca gtcaaaaggg tgggtcccca gaagcaggcg 720
agcgtgggcta agaaaaaaaaa gctggaaaaa gaagaagagg aataaagagg cagcgctaga 780
gacaagagcc ctgtgaaaaaa gcccaaaagc acagccagag ccaagaagcc aggatccaag 840
aaatgaggag ccacgccttg gggggcaccg tgcaaaagtg gcttccctg ggctgtgctg 900
cagggcacagg gtgcccctgt ccagccccca cactgtgtgc tgaatgcaac aggggtgttg 960
cgggggcaac atgagagccc ctaacccccca actctccact ttcaggaggc ccccatgtaa 1020
gagccccacc tccgggtcaca ataagtgt

```

&lt;210&gt; 107

&lt;211&gt; 1349

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc-feature

&lt;223&gt; Incyte ID No.: 3573661CB1

&lt;400&gt; 107

```

gagaaagcag agctatttca agagtgaagc acagaaggga atccagaggc catctaagcg 60
aggaagggtg tacaggcgagtg gagtgaaggg caggagcagg gccacgacca ggcacgacca 120
ccgaggggatg gaacttcaca gtggggtttca agccgctgct aggggatgca cacacgatgg 180
acaaacctgga gaagcagctc atctgcecca tctgctctgga gatgtctcc aaaccagtgg 240
tgatctctgcc ctgccaacac aacctgtgtg gcaaatgtgc caacgagctc ttccaggcct 300
cgaatctctct atggcagtc cggggctcca ccactgtgtc ttcaggaggc cgtttccgct 360
gccactcgtg caggcatgag gttgtctctg acagacagc gtctacagg ctgacggcaa 420
acgtgtctagt ggagaacatt atcgacattt acaagcagga gtcatccaag cgcgtgact 480
cgaaggctga atgtgcgagg agcatgaaag agcatgaaag cttcgggtgc cacaaggact 540
gacctgagct tgagggtccc acctgctctc tctgcaaggt cttcgggtgc agcgatggca 600
gtgaggtggc cccactttaca accatggcga aagcgcagaa gaggtagctc atggagaggg 720
ctcgctgtct ggtggcaggc aatgaccgag tgcaagcagt gatcacacag atggaggttg 780
tgtgtccagag tatcgaggac aatagccgga ggcagaagca gtgtgttaac cagaggtttg 840
agagcctgtg ccgagtgctg gaggagcgca aggggtgagct gctgcaggcg cttggccggg 840
agcaagaggga gaagctgcag cgcgtcccg gctcatccg tcagtatggc gaccacctgg 900
agggctctct taagtgtgtg gagtctgcca tccagtccat ggaagaggcca caaatggcgc 960

```



gagctctgtcc tgcaagataa cagctttgac ctgaataatg ttagtgacgc tgaacaggaa 300  
 gaaatggaaa ctcaatcttc agacttccca ccactccctga cccagccagc tcttgatcag 360  
 tcatccacta tctccttaca tccagcaacc tccaccagcag tctcgccaac aaactcccca 420  
 cgagctctccc tagtggtttc tccagcagcc tcccccagaaa tctctccaga agtttgtccc 480  
 cgagcttcta cagttgtctc tccagcagtc tctctcagtg tctctccagc 540  
 tctctccccc cagttctctt agaagtcctc ttagcggctt cagtgacatc cccaaaaagcc 600  
 tctcccgttaa ctctcccagc agctgccttt ccaacagcct cccagcaaaa taaggatgct 660  
 agcagctttc tagaaaccac tcttgacgtg gaagagatca ctggagaagg actcactgct 720  
 tctggtagtg gtgatgtcat gaggagacgt attgtaccc cagaagaagt tctgtctccc 780  
 tctccaaatg ggtggcggag agaggtgctc atcaagaaca gcagccacag atggcagggg 840  
 gagacctgtg attatggccc ctgtgggaag aggatgaagc aatttccaga agtgatcaag 900  
 tacctgagcc gcaacgtggt acacagtgtc cgcgcagagc acttcagctt cagtcctcgt 960  
 atgctgtgtg gagatttctt tgaagaaaga gacacgccaag agggcttgca gtgggtgcag 1020  
 ctctcagcag agggagatccc gtccaggatt caggcaatta ctggcaaacg ggtcgacct 1080  
 cgaaacactg agaaggctaa gactaaggaa gtccccagg tgaaacgggg tcgaggtcgg 1140  
 ccacctaagg tcaaaatcac tgagctattg aacaagacag acaaccgccc cctaagaaa 1200  
 ctggaggccc aagaaacatt gaatgaggag gtcaccaagg agatttgtca aagcaagaag 1260  
 aagatgaggg agaaggttca acggggagag tgtcagacta ctatccaagg gcaggccaga 1320  
 aataagcgga aacaagagac caagagctta aagcagaagg aagctaagaa gaaatccaag 1380  
 cgtgagaaag aaaaaggaaa gacaaagcag gaaaaactga aggaaaaagt caagagggaa 1440  
 aagaaggaga aggttaaaat gaaggaaaaag gaggaggtga ccaagcccaa ccagcctgt 1500  
 aaagcagata agaccctgtgc cacacagagg cgcttgagg aacggcagag gcagcagatg 1560  
 atcttgaggc acttgagaag gccgacagag gatattgtgtc tgactgcacca ccagccctgt 1620  
 cctgacttct cactgactccc tggcttgaca ttgcccagtg gagccttctc agactgctgt 1680  
 accattgttg agttctctga tagctttggc aaggtgtctgt gccttgatct tgcccaagtg 1740  
 tgtgcttagc ctggggggtc ctgcaggaag gggctctctgt gttcaagggt accactctgt 1800  
 tgaaggtgc aaaaaccg 1818

&lt;210&gt; 110

&lt;211&gt; 785

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 788

&lt;223&gt; a o r g c or t, unknown, or other

&lt;223&gt; Incyte ID No.: 4717936CB1

&lt;400&gt; 110

gtctctgtag tcatgaggct gaaggggggt gggacagtg tgaataaagg cactagaggg 60  
 agcccccaca cctctctctg gaactgttgc ccacatgcag ccccgagcac agccccttag 120  
 ccaaaccccta cctctctcc tccgaggggc cctctgagac cctcggtctgc ggtgtcctgt 180  
 cattaagatg ggcacaggggt gggagggtct ccagcggacc ctgaaggaag tcgctacat 240  
 ctctctctgc tctctgtgta tcaaggaact gctggattaa tggtagcagg gaactgcctc 300  
 ctctccccac cagtcacatt gctggcatcg ctacaggtggg cagggttagag taacaggagg 360  
 gcatagctgc agctctctgt gcagagcttg ccttagcttc tcattctctt ctttagcccc 420  
 cagcccgaatt gccatcaaga ctctgaagc cagctgtgct tgaccaagga tgggccaata 480  
 acaatgagta aacagtaaa gtgtgatcct gcttgagct gtgtcatcta gcagactgcg 540  
 ctcatctctt agcctctctc attcccaacc cctgctctgt gtggacctct gtggggcagg 600  
 agcttggaact ctctctgcta ggaccccagt aagattgtgg caagacctgg cactctcca 660  
 agcttgccac agtgagccca ccagctcaga tgggtgatct taccataccc tcatagtacc 720  
 aagaatggac tgccccnta agaaacctgt ttgagaatca ctgaattata aataaacc 780  
 cagat 785